



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2018 – 11:53 am GMT

PDB ID : 1JSC  
Title : Crystal Structure of the Catalytic Subunit of Yeast Acetohydroxyacid Synthase: A target for Herbicidal Inhibitors  
Authors : Pang, S.S.; Duggleby, R.G.; Guddat, L.W.  
Deposited on : 2001-08-17  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	630	<div> <div>3%</div> <div> <div></div> <div>68%</div> <div>17%</div> <div>•</div> <div>14%</div> </div> </div>
1	B	630	<div> <div>9%</div> <div> <div></div> <div>66%</div> <div>20%</div> <div>•</div> <div>13%</div> </div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8628 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETOHYDROXY-ACID SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	541	Total	C	N	O	S	0	0	0
			4071	2577	700	775	19			
1	B	550	Total	C	N	O	S	0	0	0
			4055	2560	692	784	19			


- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	K	0	0
			1	1		
2	A	1	Total	K	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is DIHYDROGENPHOSPHATE ION (three-letter code: 2HP) (formula: H<sub>2</sub>O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O<sub>7</sub>P<sub>2</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
5	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 6 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
6	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	203	Total	O	0	0
			203	203		
7	B	122	Total	O	0	0
			122	122		





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.55Å 109.40Å 178.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 2.60 41.93 – 2.59	Depositor EDS
% Data completeness (in resolution range)	94.1 (100.00-2.60) 93.2 (41.93-2.59)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.93 (at 2.58Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.188 , 0.219 0.188 , 0.219	Depositor DCC
$R_{free}$ test set	5807 reflections (10.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.2	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 54.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8628	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2HP, MG, K, FAD, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/4152	0.60	1/5639 (0.0%)
1	B	0.32	0/4131	0.57	0/5619
All	All	0.32	0/8283	0.58	1/11258 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	614	LEU	CA-CB-CG	5.05	126.91	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4071	0	4066	113	0
1	B	4055	0	3938	111	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	5	0	0	0	0
4	B	10	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	26	0	16	0	0
5	B	26	0	16	1	0
6	A	53	0	31	0	0
6	B	53	0	31	0	0
7	A	203	0	0	6	0
7	B	122	0	0	2	0
All	All	8628	0	8098	216	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 216 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:474:GLN:HE21	1:A:474:GLN:H	1.09	1.01
1:A:403:ILE:H	1:A:420:GLN:HE21	1.07	0.94
1:A:151:ARG:HH22	1:A:336:GLN:HE22	1.20	0.87
1:B:365:GLN:HA	1:B:390:PRO:HD2	1.56	0.84
1:B:499:GLN:HE22	1:B:580:GLN:NE2	1.77	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	535/630 (85%)	517 (97%)	15 (3%)	3 (1%)	27 51
1	B	536/630 (85%)	502 (94%)	29 (5%)	5 (1%)	19 38
All	All	1071/1260 (85%)	1019 (95%)	44 (4%)	8 (1%)	24 46

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	464	GLU
1	B	394	ARG
1	A	597	HIS
1	B	390	PRO
1	B	437	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/518 (84%)	428 (99%)	6 (1%)	69	87
1	B	422/518 (82%)	415 (98%)	7 (2%)	63	83
All	All	856/1036 (83%)	843 (98%)	13 (2%)	67	86

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	614	LEU
1	B	439	PHE
1	B	560	LEU
1	A	560	LEU
1	B	557	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	554	ASN
1	A	622	GLN
1	B	554	ASN
1	A	600	GLN
1	B	169	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	2HP	A	698	-	4,4,4	1.32	0	6,6,6	0.39	0
5	TPP	A	700	3	21,27,27	1.59	4 (19%)	26,40,40	1.67	7 (26%)
6	FAD	A	701	-	51,58,58	2.58	24 (47%)	57,89,89	1.84	6 (10%)
4	2HP	B	1697	-	4,4,4	1.34	0	6,6,6	0.39	0
4	2HP	B	1698	-	4,4,4	1.33	0	6,6,6	0.38	0
5	TPP	B	1700	3	21,27,27	1.55	4 (19%)	26,40,40	1.73	7 (26%)
6	FAD	B	1701	-	51,58,58	2.55	23 (45%)	57,89,89	1.75	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	2HP	A	698	-	-	0/0/0/0	0/0/0/0
5	TPP	A	700	3	-	0/16/17/17	0/2/2/2
6	FAD	A	701	-	-	0/28/50/50	0/6/6/6
4	2HP	B	1697	-	-	0/0/0/0	0/0/0/0
4	2HP	B	1698	-	-	0/0/0/0	0/0/0/0
5	TPP	B	1700	3	-	0/16/17/17	0/2/2/2
6	FAD	B	1701	-	-	0/28/50/50	0/6/6/6

The worst 5 of 55 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1700	TPP	C4-N3	-3.45	1.36	1.39
5	A	700	TPP	C4-N3	-2.70	1.37	1.39
5	A	700	TPP	PB-O2B	-2.11	1.46	1.54
5	B	1700	TPP	PB-O2B	-2.10	1.46	1.54
6	B	1701	FAD	C2-N3	2.10	1.42	1.38

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	701	FAD	C4X-C4-N3	-5.19	116.09	123.47
6	B	1701	FAD	C4X-C4-N3	-5.14	116.16	123.47
5	B	1700	TPP	N1'-C2'-N3'	-2.89	120.45	125.55
5	A	700	TPP	N1'-C2'-N3'	-2.72	120.76	125.55
5	B	1700	TPP	CM4-C4-C5	-2.60	121.92	127.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1700	TPP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	541/630 (85%)	-0.17	21 (3%) 39 31	33, 48, 90, 140	0
1	B	550/630 (87%)	0.31	58 (10%) 6 3	33, 56, 136, 147	0
All	All	1091/1260 (86%)	0.07	79 (7%) 15 11	33, 50, 125, 147	0

The worst 5 of 79 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	649	VAL	8.0
1	B	590	PHE	6.6
1	B	82	GLU	6.3
1	B	396	ALA	6.2
1	B	593	HIS	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	2HP	B	1698	5/5	0.83	0.26	133,133,133,133	0
4	2HP	A	698	5/5	0.85	0.24	120,120,121,121	0
4	2HP	B	1697	5/5	0.92	0.12	115,115,115,115	0
6	FAD	B	1701	53/53	0.95	0.19	54,60,86,87	0
5	TPP	A	700	26/26	0.98	0.13	36,46,49,50	0
6	FAD	A	701	53/53	0.98	0.16	40,49,73,74	0
3	MG	A	699	1/1	0.98	0.06	47,47,47,47	0
2	K	B	1696	1/1	0.98	0.20	43,43,43,43	0
2	K	A	696	1/1	0.99	0.19	32,32,32,32	0
3	MG	B	1699	1/1	0.99	0.06	42,42,42,42	0
5	TPP	B	1700	26/26	0.99	0.13	35,42,45,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.