



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 04:37 am GMT

PDB ID : 4JSR  
Title : Crystal Structure of human SIRT3 with ELT inhibitor 11c [N-{2-[1-(6-carbamoylthieno[3,2-d]pyrimidin-4-yl)piperidin-4-yl]ethyl}-N'-ethylthiophene-2,5-dicarboxamide]  
Authors : Dai, H.  
Deposited on : 2013-03-22  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	3793 (1.70-1.70)
Clashscore	122126	4167 (1.70-1.70)
Ramachandran outliers	120053	4100 (1.70-1.70)
Sidechain outliers	120020	4100 (1.70-1.70)
RSRZ outliers	108989	3718 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-3, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	2	0
			2153	1386	371	387	9			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	115	SER	-	EXPRESSION TAG	UNP Q9NTG7
A	116	ASN	-	EXPRESSION TAG	UNP Q9NTG7
A	117	ALA	-	EXPRESSION TAG	UNP Q9NTG7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is N-{2-[1-(6-carbamoylthieno[3,2-d]pyrimidin-4-yl)piperidin-4-yl]ethyl}-N'-ethyl thiophene-2,5-dicarboxamide (three-letter code: 1NQ) (formula: C<sub>22</sub>H<sub>26</sub>N<sub>6</sub>O<sub>3</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			33	22	6	3	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	306	Total O 306 306	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.32Å 119.32Å 44.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 1.70 35.69 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.00-1.70) 100.0 (35.69-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.99 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.198 , 0.228 0.189 , 0.214	Depositor DCC
$R_{free}$ test set	2006 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.028 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2493	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 1NQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.42	10/2218 (0.5%)	1.26	7/3021 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	259	CYS	CB-SG	-8.26	1.68	1.82
1	A	185	PHE	CE1-CZ	7.07	1.50	1.37
1	A	251	PHE	CE2-CZ	6.76	1.50	1.37
1	A	379	TRP	CG-CD1	6.29	1.45	1.36
1	A	251	PHE	CG-CD1	5.90	1.47	1.38
1	A	277	VAL	CB-CG2	5.87	1.65	1.52
1	A	283	CYS	CB-SG	5.85	1.92	1.82
1	A	293	PHE	CE1-CZ	5.84	1.48	1.37
1	A	282	VAL	CB-CG1	5.67	1.64	1.52
1	A	180	PHE	CE1-CZ	5.12	1.47	1.37

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	269	ARG	NE-CZ-NH2	-9.31	115.64	120.30
1	A	340	ARG	NE-CZ-NH1	-7.89	116.35	120.30
1	A	347	LEU	CB-CG-CD2	7.51	123.77	111.00
1	A	269	ARG	NE-CZ-NH1	7.42	124.01	120.30
1	A	351	LEU	CB-CG-CD2	6.75	122.47	111.00
1	A	179	ILE	CG1-CB-CG2	5.37	123.22	111.40
1	A	243	LYS	CD-CE-NZ	-5.34	99.42	111.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2168	32	0
2	A	1	0	0	0	0
3	A	33	0	26	6	0
4	A	306	0	0	9	1
All	All	2493	0	2194	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:GLU:HG3	4:A:693:HOH:O	1.63	0.96
3:A:402:1NQ:SAV	3:A:402:1NQ:H17	2.12	0.89
1:A:207:ASN:H	1:A:210:HIS:HD2	1.19	0.87
1:A:215:LEU:HD22	1:A:383:MET:HE2	1.58	0.85
1:A:387:VAL:O	1:A:391:THR:HG23	1.78	0.84
1:A:135:ARG:HA	4:A:659:HOH:O	1.81	0.80
3:A:402:1NQ:CAA	3:A:402:1NQ:OAD	2.34	0.75
1:A:205:LYS:HE2	1:A:394:LEU:HD12	1.70	0.73
1:A:205:LYS:NZ	1:A:391:THR:HB	2.05	0.72
1:A:228:GLN:O	1:A:248:HIS:HD2	1.73	0.71
1:A:203:ASN:HB3	4:A:667:HOH:O	1.94	0.65
1:A:207:ASN:H	1:A:210:HIS:CD2	2.11	0.64
1:A:207:ASN:HD22	1:A:209:THR:H	1.45	0.64
3:A:402:1NQ:H3	3:A:402:1NQ:OAD	1.98	0.63
1:A:235:ARG:HD3	4:A:716:HOH:O	1.99	0.61
1:A:388:GLN:HG3	4:A:669:HOH:O	2.00	0.60
1:A:215:LEU:HD13	1:A:386:LEU:HD22	1.84	0.59
1:A:354:HIS:HD2	4:A:796:HOH:O	1.85	0.58
1:A:215:LEU:HD22	1:A:383:MET:CE	2.35	0.53
1:A:205:LYS:HZ3	1:A:391:THR:HB	1.73	0.53
1:A:205:LYS:HZ1	1:A:391:THR:HB	1.72	0.52
1:A:266:GLU:CG	4:A:693:HOH:O	2.39	0.52
1:A:177:GLU:O	1:A:181:GLU:HG3	2.11	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:ASN:ND2	1:A:209:THR:H	2.08	0.51
1:A:219:LYS:HE3	1:A:379:TRP:CE2	2.47	0.49
1:A:296:GLU:HB3	1:A:297:PRO:HD2	1.95	0.48
3:A:402:1NQ:H10	3:A:402:1NQ:H19	1.76	0.47
3:A:402:1NQ:SAV	3:A:402:1NQ:CAP	2.95	0.47
1:A:261:ARG:HH21	1:A:282:VAL:HG23	1.80	0.46
3:A:402:1NQ:H2	3:A:402:1NQ:OAD	2.14	0.46
1:A:318:LEU:HD23	1:A:343:ILE:HB	1.98	0.46
1:A:345:ARG:HG2	4:A:605:HOH:O	2.16	0.45
1:A:175:TYR:HB2	1:A:176:PRO:HD2	1.99	0.45
1:A:145:GLY:HA3	1:A:320:THR:HB	2.00	0.44
1:A:324:VAL:HG13	4:A:788:HOH:O	2.18	0.44
1:A:150:THR:OG1	1:A:151:PRO:HD3	2.18	0.43
1:A:259:CYS:O	1:A:260[B]:GLN:HB2	2.18	0.43
1:A:215:LEU:CD2	1:A:383:MET:HE2	2.41	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:695:HOH:O	4:A:802:HOH:O[4_665]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	274/285 (96%)	265 (97%)	9 (3%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	239/246 (97%)	227 (95%)	12 (5%)	27 9

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	126	GLN
1	A	138	GLN
1	A	200	TYR
1	A	207	ASN
1	A	225	LEU
1	A	300	GLN
1	A	307	VAL
1	A	311	MET
1	A	324	VAL
1	A	336	SER
1	A	347	LEU
1	A	351	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	ASN
1	A	207	ASN
1	A	210	HIS
1	A	248	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	1NQ	A	402	-	27,36,36	1.01	1 (3%)	31,50,50	2.32	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1NQ	A	402	-	-	1/17/35/35	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	1NQ	CBB-SAV	-3.51	1.67	1.72

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	1NQ	N3-C2-N1	-8.04	121.98	128.86
3	A	402	1NQ	C4-C5-SAV	-3.93	107.16	112.00
3	A	402	1NQ	C2-N1-C6	3.52	120.36	111.81
3	A	402	1NQ	CAO-NBG-CAP	3.92	119.97	111.54
3	A	402	1NQ	C2-N3-C4	6.44	121.70	115.24

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	1NQ	CAA-CAJ-NAS-CAX

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	1NQ	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	274/285 (96%)	-0.10	8 (2%) 51 56	16, 24, 39, 59	5 (1%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	392	GLY	6.1
1	A	394	LEU	4.5
1	A	121	GLY	4.3
1	A	306	VAL	3.5
1	A	393	LYS	3.0
1	A	388	GLN	2.4
1	A	391	THR	2.3
1	A	384	ARG	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	1NQ	A	402	33/33	0.95	0.09	14,19,32,44	0
2	ZN	A	401	1/1	0.99	0.07	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.