



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 20, 2019 – 10:14 AM EDT

PDB ID : 1K3U
Title : CRYSTAL STRUCTURE OF WILD-TYPE TRYPTOPHAN SYNTHASE
COMPLEXED WITH N-[1H-INDOL-3-YL-ACETYL]ASPARTIC ACID
Authors : Weyand, M.; Schlichting, I.; Marabotti, A.; Mozzarelli, A.
Deposited on : 2001-10-04
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.3.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.3.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3793 (1.70-1.70)
Clashscore	122126	4167 (1.70-1.70)
Ramachandran outliers	120053	4100 (1.70-1.70)
Sidechain outliers	120020	4100 (1.70-1.70)
RSRZ outliers	108989	3718 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	<div> <div>5%</div> <div>83%</div> <div>16%</div> <div>.</div> </div>
2	B	396	<div> <div>2%</div> <div>87%</div> <div>12%</div> <div>..</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5783 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan Synthase Alpha Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	4	0
			2030	1286	353	383	8			


- Molecule 2 is a protein called Tryptophan Synthase Beta Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	394	Total	C	N	O	S	0	7	0
			3004	1886	526	572	20			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	34	SER	ARG	CLONING ARTIFACT	UNP P0A2K1

- Molecule 3 is N-[1H-INDOL-3-YL-ACETYL]ASPARTIC ACID (three-letter code: IAD) (formula: $C_{14}H_{14}N_2O_5$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			21	14	2	5		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	261	Total	O	0	0
			261	261		
6	B	451	Total	O	0	2
			451	451		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan Synthase Alpha Chain



• Molecule 2: Tryptophan Synthase Beta Chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	182.41 Å 59.72 Å 67.18 Å 90.00° 94.52° 90.00°	Depositor
Resolution (Å)	20.00 – 1.70 31.06 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.7 (20.00-1.70) 97.7 (31.06-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 1.70 Å)	Xtriage
Refinement program	REFMAC 5.0, CNS	Depositor
R, R_{free}	0.153 , 0.188 0.156 , 0.192	Depositor DCC
R_{free} test set	3939 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.456	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5783	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IAD, NA, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.22	4/2089 (0.2%)	1.16	10/2837 (0.4%)
2	B	1.34	12/3098 (0.4%)	1.27	24/4181 (0.6%)
All	All	1.30	16/5187 (0.3%)	1.23	34/7018 (0.5%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	291	ASP	CB-CG	-10.75	1.29	1.51
2	B	298	TYR	CE1-CZ	7.48	1.48	1.38
2	B	152[A]	MET	SD-CE	-7.26	1.37	1.77
2	B	152[B]	MET	SD-CE	-7.26	1.37	1.77
2	B	52	TYR	CD2-CE2	7.04	1.50	1.39
2	B	206	ARG	CG-CD	6.57	1.68	1.51
1	A	82	PHE	CD2-CE2	6.55	1.52	1.39
2	B	364	GLU	CD-OE1	-6.30	1.18	1.25
2	B	210	GLU	CD-OE2	6.11	1.32	1.25
2	B	143	SER	CA-CB	6.08	1.62	1.52
2	B	385	PHE	CB-CG	-5.96	1.41	1.51
1	A	123	VAL	CB-CG2	5.54	1.64	1.52
2	B	45	PHE	CE1-CZ	5.49	1.47	1.37
1	A	140	ARG	CG-CD	5.30	1.65	1.51
1	A	18	ALA	CA-CB	5.29	1.63	1.52
2	B	364	GLU	CD-OE2	5.14	1.31	1.25

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	291	ASP	CB-CG-OD1	-13.04	106.56	118.30
2	B	275	ARG	NE-CZ-NH2	9.38	124.99	120.30
2	B	286	MET	CA-CB-CG	9.04	128.68	113.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	175	ARG	NE-CZ-NH1	8.75	124.68	120.30
2	B	175	ARG	NE-CZ-NH2	-7.45	116.57	120.30
1	A	93	PRO	N-CD-CG	-7.05	92.62	103.20
1	A	42	ASP	CB-CG-OD2	7.00	124.60	118.30
2	B	394	ARG	NE-CZ-NH2	-7.00	116.80	120.30
2	B	225	ASP	CB-CG-OD2	6.75	124.37	118.30
2	B	381	ASP	CB-CG-OD2	6.53	124.17	118.30
2	B	206	ARG	NE-CZ-NH1	6.50	123.55	120.30
1	A	225	ARG	NE-CZ-NH2	-6.31	117.15	120.30
1	A	159	ASP	CB-CG-OD2	6.24	123.91	118.30
1	A	130	ASP	CB-CG-OD1	6.21	123.89	118.30
2	B	150	ARG	NE-CZ-NH1	-6.12	117.24	120.30
2	B	131	ARG	NE-CZ-NH1	6.07	123.33	120.30
1	A	42	ASP	CB-CG-OD1	-6.06	112.85	118.30
2	B	38	ASP	CB-CG-OD2	6.03	123.72	118.30
2	B	385	PHE	CB-CG-CD2	-5.99	116.61	120.80
2	B	202	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	70	ARG	NE-CZ-NH2	-5.73	117.44	120.30
2	B	176	ASP	CB-CG-OD2	5.61	123.35	118.30
2	B	329	ASP	CB-CG-OD1	5.55	123.30	118.30
2	B	222	ARG	CG-CD-NE	-5.52	100.21	111.80
2	B	315	TYR	CB-CG-CD2	-5.44	117.73	121.00
2	B	364	GLU	CG-CD-OE1	5.42	129.14	118.30
2	B	372	LEU	CB-CG-CD1	-5.40	101.82	111.00
1	A	157	ASN	N-CA-CB	-5.36	100.95	110.60
1	A	175	TYR	CG-CD2-CE2	-5.32	117.04	121.30
2	B	298	TYR	CA-CB-CG	-5.27	103.38	113.40
2	B	41	PHE	CB-CG-CD1	-5.24	117.13	120.80
2	B	121	LEU	CB-CG-CD1	-5.21	102.15	111.00
1	A	11	LEU	CB-CG-CD1	-5.20	102.17	111.00
2	B	100	ARG	NE-CZ-NH2	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2030	0	2030	28	0
2	B	3004	0	2979	22	0
3	A	21	0	12	0	0
4	B	1	0	0	0	0
5	B	15	0	6	0	0
6	A	261	0	0	7	1
6	B	451	0	0	8	1
All	All	5783	0	5027	50	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:219[B]:LYS:HE3	6:B:1253:HOH:O	1.09	1.23
1:A:108:ASN:ND2	1:A:109:ASN:OD1	2.00	0.93
1:A:69:LEU:HD23	6:A:991:HOH:O	1.78	0.81
1:A:117[B]:ARG:NH1	6:A:1041:HOH:O	2.25	0.69
2:B:219[B]:LYS:CE	6:B:1253:HOH:O	1.88	0.64
1:A:188:ARG:CB	6:A:1051:HOH:O	2.46	0.64
1:A:30:ILE:HD12	1:A:84:MET:SD	2.38	0.64
2:B:337:LYS:HE2	2:B:391:LEU:HD21	1.82	0.62
1:A:108:ASN:HA	6:A:889:HOH:O	2.01	0.60
1:A:113:ALA:O	1:A:117[B]:ARG:HG3	2.01	0.59
2:B:44:GLN:HG3	6:B:1236:HOH:O	2.02	0.58
2:B:379:ARG:NH1	6:B:1190:HOH:O	2.40	0.55
1:A:112:ASP:OD1	1:A:146:HIS:HE1	1.90	0.54
2:B:341:ARG:HD3	6:B:1061:HOH:O	2.08	0.54
2:B:337:LYS:CE	2:B:391:LEU:HD21	2.39	0.53
1:A:194:HIS:O	1:A:198:GLU:HG2	2.09	0.51
2:B:301:SER:OG	2:B:350:GLU:HG3	2.12	0.50
1:A:100:LEU:C	1:A:100:LEU:HD13	2.32	0.49
2:B:115:HIS:CE1	2:B:189:GLY:HA2	2.47	0.49
2:B:385:PHE:HB2	6:B:984:HOH:O	2.13	0.48
2:B:129:LYS:HE2	6:B:1065:HOH:O	2.12	0.48
1:A:171:ARG:HH11	1:A:171:ARG:HG2	1.80	0.47
1:A:22:PHE:HA	1:A:49:GLU:O	2.16	0.46
2:B:276:VAL:HA	2:B:285:PRO:HA	1.97	0.46
1:A:156:PRO:HB2	1:A:191:LEU:CD2	2.45	0.46
1:A:30:ILE:HD11	1:A:76:VAL:HG22	1.97	0.45
2:B:2:THR:N	6:B:1054:HOH:O	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:LEU:HB2	6:A:961:HOH:O	2.17	0.44
2:B:86:HIS:CE1	2:B:236:ASN:HB3	2.53	0.44
1:A:53:PRO:HA	1:A:68:ASN:OD1	2.18	0.44
1:A:191:LEU:HA	1:A:192:PRO:HD2	1.65	0.43
1:A:156:PRO:O	1:A:196:LEU:HD21	2.17	0.43
2:B:337:LYS:HE2	2:B:391:LEU:CD2	2.46	0.43
2:B:391:LEU:O	2:B:395:GLY:N	2.49	0.43
1:A:157:ASN:O	1:A:158:ALA:C	2.57	0.43
1:A:66[B]:ASN:ND2	6:A:881:HOH:O	2.46	0.42
2:B:38:ASP:OD1	2:B:38:ASP:C	2.57	0.42
1:A:22:PHE:CD1	1:A:22:PHE:C	2.92	0.42
1:A:186:GLU:OE2	1:A:239:LYS:HE2	2.19	0.42
2:B:337:LYS:HE2	2:B:391:LEU:HD11	1.99	0.42
1:A:199:LYS:HD3	1:A:199:LYS:HA	1.76	0.42
1:A:156:PRO:HB2	1:A:191:LEU:HD23	2.02	0.42
2:B:195:HIS:CG	2:B:196:PRO:HA	2.54	0.41
2:B:263:GLU:CD	2:B:263:GLU:H	2.23	0.41
1:A:156:PRO:HA	1:A:196:LEU:HD21	2.04	0.41
1:A:36:ILE:O	1:A:40:LEU:HG	2.21	0.40
1:A:95:ILE:O	1:A:95:ILE:HG23	2.21	0.40
1:A:10:GLN:HB3	6:A:979:HOH:O	2.22	0.40
2:B:376:LEU:HD12	2:B:376:LEU:HA	1.95	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:939:HOH:O	6:B:1239:HOH:O[1_565]	2.14	0.06

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/268 (101%)	266 (98%)	3 (1%)	1 (0%)	36	19
2	B	399/396 (101%)	389 (98%)	10 (2%)	0	100	100
All	All	669/664 (101%)	655 (98%)	13 (2%)	1 (0%)	49	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/208 (101%)	209 (99%)	2 (1%)	81	73
2	B	315/310 (102%)	314 (100%)	1 (0%)	93	90
All	All	526/518 (102%)	523 (99%)	3 (1%)	87	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	LEU
1	A	239	LYS
2	B	384	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	HIS
2	B	44	GLN
2	B	236	ASN
2	B	246	ASN
2	B	375	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	IAD	A	801	-	15,22,22	1.37	2 (13%)	16,30,30	1.17	3 (18%)
5	PLP	B	802	2	15,15,16	1.21	2 (13%)	20,22,23	2.24	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IAD	A	801	-	-	6/10/16/16	0/2/2/2
5	PLP	B	802	2	-	0/6/6/8	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	801	IAD	C11-N	3.27	1.40	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	802	PLP	C5-C4	-2.65	1.37	1.40
3	A	801	IAD	CB-CA	2.53	1.57	1.53
5	B	802	PLP	C5A-C5	2.42	1.57	1.50

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	802	PLP	C4A-C4-C5	-6.06	114.98	120.91
5	B	802	PLP	O4P-C5A-C5	4.29	117.53	109.35
5	B	802	PLP	C3-C2-N1	-3.33	116.38	120.75
5	B	802	PLP	C4A-C4-C3	2.94	125.45	120.51
3	A	801	IAD	C-CA-N	-2.81	106.31	112.39
5	B	802	PLP	O3P-P-O4P	2.47	113.31	106.73
3	A	801	IAD	C5-C4-C9	-2.15	117.91	120.89
3	A	801	IAD	C4-C9-C8	2.13	120.99	118.17
5	B	802	PLP	C2A-C2-C3	2.06	123.42	120.96
5	B	802	PLP	O4P-P-O1P	-2.03	100.79	106.47

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	801	IAD	C-CA-CB-CG
3	A	801	IAD	CB-CA-N-C11
3	A	801	IAD	C-CA-N-C11
3	A	801	IAD	N-CA-CB-CG
3	A	801	IAD	C3-C10-C11-N
3	A	801	IAD	C3-C10-C11-O11

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/268 (100%)	0.02	14 (5%) 27 30	11, 18, 36, 55	0
2	B	394/396 (99%)	-0.20	7 (1%) 68 73	9, 14, 25, 64	0
All	All	662/664 (99%)	-0.11	21 (3%) 47 52	9, 15, 31, 64	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	191	LEU	6.2
1	A	190	ALA	5.8
2	B	393	ALA	5.3
2	B	395	GLY	4.8
1	A	192	PRO	4.8
2	B	385	PHE	4.4
1	A	189	GLY	4.3
2	B	394	ARG	3.4
1	A	194	HIS	3.0
2	B	2	THR	2.8
1	A	187	ASN	2.8
1	A	15	ARG	2.6
2	B	390	ILE	2.6
1	A	195	HIS	2.5
1	A	1	MET	2.5
1	A	186	GLU	2.3
2	B	392	LYS	2.3
1	A	249	LYS	2.3
1	A	268	ALA	2.2
1	A	157	ASN	2.1
1	A	2	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	IAD	A	801	21/21	0.94	0.09	14,17,19,20	0
5	PLP	B	802	15/16	0.98	0.10	9,11,13,14	0
4	NA	B	803	1/1	1.00	0.09	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around IAD A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.