



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 13, 2020 – 03:57 PM EST

PDB ID : 2K3U  
Title : Structure of the tyrosine-sulfated C5a receptor N-terminus in complex with the immune evasion protein CHIPS.  
Authors : Ippel, J.H.; Bunschoten, A.; Kemmink, J.; Liskamp, R.  
Deposited on : 2008-05-16

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.4  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 89%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136327	12091
Ramachandran outliers	132723	10835
Sidechain outliers	132532	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	91	 82% 8% 10%
2	B	24	 33% 8% 58%

## 2 Ensemble composition and analysis

This entry contains 25 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy and chemical shift difference*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:37-A:118, B:12-B:13, B:15-B:22 (92)	0.37	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 6, 8, 9, 11, 14, 16, 19, 20, 21, 22, 23
2	5, 12, 24
3	7, 10, 13
4	15, 17
Single-model clusters	18; 25

### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1839 atoms, of which 912 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Chemotaxis inhibitory protein.

Mol	Chain	Residues	Atoms						Trace
1	A	91	Total	C	H	N	O	S	0
			1489	473	751	122	141	2	

- Molecule 2 is a protein called C5aR(P7-28S) peptide.


Mol	Chain	Residues	Atoms						Trace
2	B	24	Total	C	H	N	O	S	1
			350	111	161	28	48	2	

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Chemotaxis inhibitory protein



- Molecule 2: C5aR(P7-28S) peptide




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: Chemotaxis inhibitory protein





- Molecule 2: C5aR(P7-28S) peptide



### 4.2.2 Score per residue for model 2


- Molecule 1: Chemotaxis inhibitory protein

Chain A:  84% 5% • 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  42% 58%



### 4.2.3 Score per residue for model 3

- Molecule 1: Chemotaxis inhibitory protein

Chain A:  78% 11% • 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  33% 58%



### 4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: Chemotaxis inhibitory protein

Chain A:  80% 9% • 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  29% 13% 58%



### 4.2.5 Score per residue for model 5


- Molecule 1: Chemotaxis inhibitory protein

Chain A:  82% 8% 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  33% 8% 58%



### 4.2.6 Score per residue for model 6


- Molecule 1: Chemotaxis inhibitory protein

Chain A:  81% 8% 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  25% 17% 58%



### 4.2.7 Score per residue for model 7


- Molecule 1: Chemotaxis inhibitory protein

Chain A:  77% 13% 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  33% 8% 58%




### 4.2.8 Score per residue for model 8

- Molecule 1: Chemotaxis inhibitory protein




- Molecule 2: C5aR(P7-28S) peptide




### 4.2.9 Score per residue for model 9

- Molecule 1: Chemotaxis inhibitory protein




- Molecule 2: C5aR(P7-28S) peptide




### 4.2.10 Score per residue for model 10

- Molecule 1: Chemotaxis inhibitory protein




- Molecule 2: C5aR(P7-28S) peptide





### 4.2.11 Score per residue for model 11

- Molecule 1: Chemotaxis inhibitory protein

Chain A:  81% 9% 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  25% 17% 58%



### 4.2.12 Score per residue for model 12

- Molecule 1: Chemotaxis inhibitory protein

Chain A:  82% 7% 10%




- Molecule 2: C5aR(P7-28S) peptide


Chain B:  38% 58%



### 4.2.13 Score per residue for model 13


- Molecule 1: Chemotaxis inhibitory protein

Chain A:  84% 5% 10%




- Molecule 2: C5aR(P7-28S) peptide

Chain B:  33% 8% 58%




#### 4.2.14 Score per residue for model 14

- Molecule 1: Chemotaxis inhibitory protein




- Molecule 2: C5aR(P7-28S) peptide




#### 4.2.15 Score per residue for model 15

- Molecule 1: Chemotaxis inhibitory protein




- Molecule 2: C5aR(P7-28S) peptide




#### 4.2.16 Score per residue for model 16

- Molecule 1: Chemotaxis inhibitory protein



- Molecule 2: C5aR(P7-28S) peptide



### 4.2.17 Score per residue for model 17


- Molecule 1: Chemotaxis inhibitory protein

Chain A: 81% 8% 10%



- Molecule 2: C5aR(P7-28S) peptide


Chain B: 25% 17% 58%



### 4.2.18 Score per residue for model 18


- Molecule 1: Chemotaxis inhibitory protein

Chain A: 81% 8% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 25% 17% 58%



### 4.2.19 Score per residue for model 19

- Molecule 1: Chemotaxis inhibitory protein

Chain A: 85% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 21% 21% 58%



#### 4.2.20 Score per residue for model 20

- Molecule 1: Chemotaxis inhibitory protein

Chain A: 81% 5% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 21% 21% 58%



#### 4.2.21 Score per residue for model 21


- Molecule 1: Chemotaxis inhibitory protein

Chain A: 79% 11% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 38% 58% 58%



#### 4.2.22 Score per residue for model 22


- Molecule 1: Chemotaxis inhibitory protein

Chain A: 84% 7% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 38% 58% 58%



#### 4.2.23 Score per residue for model 23


- Molecule 1: Chemotaxis inhibitory protein

Chain A: 79% 10% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 25% 17% 58%



#### 4.2.24 Score per residue for model 24


- Molecule 1: Chemotaxis inhibitory protein

Chain A: 84% 5% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 38% . 58%



#### 4.2.25 Score per residue for model 25

- Molecule 1: Chemotaxis inhibitory protein

Chain A: 82% 7% 10%



- Molecule 2: C5aR(P7-28S) peptide

Chain B: 33% 8% 58%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, torsion angle dynamics, simulated annealing docking, molecular dynamics*.

Of the 130 calculated structures, 25 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	refinement	1.2
ARIA	structure solution	1.2
CNSSOLVE	structure solution	1.1
TALOS	structure solution	
TALOS	geometry optimization	
YASARA	structure solution	Yasara Structure 8.3.3/WHATIF
YASARA	refinement	Yasara Structure 8.3.3/WHATIF

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	2k3u_cs.cif
Number of chemical shift lists	1
Total number of shifts	1348
Number of shifts mapped to atoms	1348
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	89%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

### 5.1 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	668	687	687	1±1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes
2	B	78	69	68	1±0
All	All	18650	18900	18875	29

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:63:VAL:HG11	2:B:20:LEU:HD23	0.54	1.79	25	18
1:A:109:VAL:HG23	2:B:13:HIS:CD2	0.52	2.40	12	2
1:A:99:PHE:CZ	2:B:22:LEU:HD13	0.47	2.45	21	2
1:A:109:VAL:HG22	2:B:15:ASP:OD1	0.45	2.11	6	1
1:A:109:VAL:HG23	2:B:13:HIS:NE2	0.43	2.29	17	2
1:A:97:TYR:CD2	1:A:109:VAL:CG1	0.42	3.02	20	2
1:A:105:LYS:CE	2:B:19:THR:HG21	0.42	2.44	3	1
1:A:95:LYS:HE2	1:A:97:TYR:CE2	0.41	2.49	9	1

## 5.2 Torsion angles ⓘ

### 5.2.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	82/91 (90%)	75±1 (92±2%)	5±1 (7±2%)	1±1 (1±1%)	17	62
2	B	10/24 (42%)	7±1 (74±13%)	2±1 (21±13%)	1±1 (5±9%)	4	24
All	All	2300/2875 (80%)	2069 (90%)	189 (8%)	42 (2%)	13	56

All 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	101	LYS	11
1	A	115	LYS	9
1	A	102	GLY	6
2	B	17	LYS	5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
2	B	19	THR	5
2	B	12	GLY	2
1	A	93	MET	2
1	A	118	GLU	1
2	B	22	LEU	1

## 5.2.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	73/80 (91%)	71±1 (97±1%)	2±1 (3±1%)	44 87
2	B	9/19 (47%)	8±1 (88±9%)	1±1 (12±9%)	9 51
All	All	2050/2475 (83%)	1960 (96%)	90 (4%)	35 82

All 21 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	109	VAL	13
1	A	100	LYS	13
1	A	38	LEU	13
2	B	15	ASP	11
2	B	21	ASP	8
2	B	17	LYS	8
1	A	49	LEU	4
1	A	55	ASN	3
1	A	106	SER	2
1	A	94	TYR	2
1	A	95	LYS	2
1	A	92	LYS	2
1	A	82	GLU	1
1	A	80	LEU	1
1	A	83	ASP	1
1	A	42	ASP	1
1	A	44	ARG	1
2	B	18	ASP	1
1	A	71	TYR	1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	78	THR	1
1	A	48	TYR	1

### 5.2.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.3 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	TYS	B	11	2	15,16,17	0.92±0.06	0±0 (0±0%)
2	TYS	B	14	2	15,16,17	1.25±0.06	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	TYS	B	11	2	18,22,24	1.40±0.13	0±0 (0±0%)
2	TYS	B	14	2	18,22,24	0.87±0.05	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYS	B	11	2	-	0±0,9,11,13	0±0,1,1,1
2	TYS	B	14	2	-	0±0,9,11,13	0±0,1,1,1

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.5 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.6 Other polymers [i](#)

There are no such molecules in this entry.

## 5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 89% for the well-defined parts and 86% for the entire structure.

### 6.1 Chemical shift list 1

File name: 2k3u\_cs.cif

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 6.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1348
Number of shifts mapped to atoms	1348
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	5

#### 6.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	91	$-0.11 \pm 0.08$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	83	$-0.03 \pm 0.11$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	91	$0.03 \pm 0.14$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	86	$0.12 \pm 0.47$	None needed ( $< 0.5$ ppm)

#### 6.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 89%, i.e. 1044 atoms were assigned a chemical shift out of a possible 1171. 9 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	426/456 (93%)	182/182 (100%)	164/184 (89%)	80/90 (89%)
Sidechain	553/624 (89%)	353/366 (96%)	185/227 (81%)	15/31 (48%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	65/91 (71%)	45/47 (96%)	20/42 (48%)	0/2 (0%)
Overall	1044/1171 (89%)	580/595 (97%)	369/453 (81%)	95/123 (77%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 86%, i.e. 1185 atoms were assigned a chemical shift out of a possible 1380. 10 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	483/545 (89%)	215/217 (99%)	182/222 (82%)	86/106 (81%)
Sidechain	625/728 (86%)	410/427 (96%)	199/267 (75%)	16/34 (47%)
Aromatic	77/107 (72%)	53/55 (96%)	24/50 (48%)	0/2 (0%)
Overall	1185/1380 (86%)	678/699 (97%)	405/539 (75%)	102/142 (72%)

#### 6.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	49	LEU	HD23	-1.14	2.14 – -0.66	-6.7
1	A	49	LEU	HD22	-1.14	2.14 – -0.66	-6.7
1	A	49	LEU	HD21	-1.14	2.14 – -0.66	-6.7
1	A	108	TYR	HD1	5.27	8.44 – 5.44	-5.6
1	A	108	TYR	HD2	5.27	8.44 – 5.44	-5.6

#### 6.1.5 Random Coil Index (RCI) plots ⓘ

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

