



Full wwPDB NMR Structure Validation Report ⓘ

Feb 16, 2018 – 01:06 am GMT

PDB ID : 2KDW
Title : Solution structure of RppH mutant E53A from Escherichia coli
Authors : Li, H.; Bi, Y.; Jin, C.
Deposited on : 2009-01-19

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk30686
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136279	12091
Ramachandran outliers	132675	10835
Sidechain outliers	132484	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	164	 91% 6% •

2 Ensemble composition and analysis

This entry contains 21 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:160 (159)	0.38	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2717 atoms, of which 1352 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called RNA pyrophosphohydrolase.

Mol	Chain	Residues	Atoms						Trace
1	A	164	Total	C	H	N	O	S	0
			2717	871	1352	249	237	8	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ALA	GLU	ENGINEERED	UNP P0A776

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA pyrophosphohydrolase




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: RNA pyrophosphohydrolase




4.2.2 Score per residue for model 2

- Molecule 1: RNA pyrophosphohydrolase




4.2.3 Score per residue for model 3

- Molecule 1: RNA pyrophosphohydrolase




4.2.4 Score per residue for model 4

- Molecule 1: RNA pyrophosphohydrolase




4.2.5 Score per residue for model 5

- Molecule 1: RNA pyrophosphohydrolase




4.2.6 Score per residue for model 6

- Molecule 1: RNA pyrophosphohydrolase




4.2.7 Score per residue for model 7

- Molecule 1: RNA pyrophosphohydrolase




4.2.8 Score per residue for model 8

- Molecule 1: RNA pyrophosphohydrolase




4.2.9 Score per residue for model 9

- Molecule 1: RNA pyrophosphohydrolase




4.2.10 Score per residue for model 10

- Molecule 1: RNA pyrophosphohydrolase




4.2.11 Score per residue for model 11

- Molecule 1: RNA pyrophosphohydrolase




4.2.12 Score per residue for model 12

- Molecule 1: RNA pyrophosphohydrolase




4.2.13 Score per residue for model 13

- Molecule 1: RNA pyrophosphohydrolase




4.2.14 Score per residue for model 14

- Molecule 1: RNA pyrophosphohydrolase




4.2.15 Score per residue for model 15

- Molecule 1: RNA pyrophosphohydrolase




4.2.16 Score per residue for model 16

- Molecule 1: RNA pyrophosphohydrolase




4.2.17 Score per residue for model 17

- Molecule 1: RNA pyrophosphohydrolase




4.2.18 Score per residue for model 18

- Molecule 1: RNA pyrophosphohydrolase




4.2.19 Score per residue for model 19

- Molecule 1: RNA pyrophosphohydrolase




4.2.20 Score per residue for model 20

- Molecule 1: RNA pyrophosphohydrolase



4.2.21 Score per residue for model 21

- Molecule 1: RNA pyrophosphohydrolase



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 21 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	
CYANA	structure solution	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.69±0.02	0±0/1362 (0.0±0.0%)	1.12±0.21	4±11/1843 (0.2±0.6%)
All	All	0.69	2/28602 (0.0%)	1.14	83/38703 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.2±1.5
All	All	0	26

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	160	GLU	CA-C	6.04	1.68	1.52	1	1
1	A	124	TRP	CG-CD2	5.36	1.52	1.43	1	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	76	ARG	NE-CZ-NH2	-19.73	110.44	120.30	1	1
1	A	141	ARG	NE-CZ-NH2	-18.93	110.83	120.30	1	1
1	A	125	ARG	NE-CZ-NH2	-17.52	111.54	120.30	1	1
1	A	8	ARG	NE-CZ-NH2	-17.05	111.78	120.30	1	1
1	A	124	TRP	NE1-CE2-CD2	-16.67	90.63	107.30	1	1
1	A	144	TYR	CB-CG-CD2	-14.17	112.50	121.00	1	7
1	A	85	ARG	NE-CZ-NH2	-13.72	113.44	120.30	1	1
1	A	144	TYR	CB-CG-CD1	11.30	127.78	121.00	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	18	ARG	NE-CZ-NH2	-11.15	114.73	120.30	1	1
1	A	62	ARG	NE-CZ-NH2	-10.51	115.05	120.30	1	1
1	A	124	TRP	CE2-CD2-CG	-9.98	99.31	107.30	1	1
1	A	125	ARG	NH1-CZ-NH2	9.75	130.12	119.40	1	1
1	A	77	TYR	CB-CG-CD2	-9.67	115.20	121.00	1	2
1	A	124	TRP	CE3-CZ3-CH2	-9.01	111.29	121.20	1	1
1	A	76	ARG	NH1-CZ-NH2	8.84	129.13	119.40	1	1
1	A	124	TRP	N-CA-CB	-8.74	94.87	110.60	1	1
1	A	21	GLN	CB-CG-CD	8.71	134.26	111.60	1	1
1	A	8	ARG	NE-CZ-NH1	8.66	124.63	120.30	1	1
1	A	160	GLU	C-N-CA	8.63	143.27	121.70	1	1
1	A	124	TRP	CZ3-CH2-CZ2	-8.28	111.66	121.60	1	1
1	A	124	TRP	CD1-CG-CD2	-7.99	99.91	106.30	1	1
1	A	134	ARG	NE-CZ-NH2	-7.84	116.38	120.30	1	1
1	A	26	ARG	NE-CZ-NH2	-7.83	116.39	120.30	1	1
1	A	125	ARG	CG-CD-NE	-7.63	95.78	111.80	1	1
1	A	91	VAL	CA-CB-CG2	7.62	122.33	110.90	1	1
1	A	124	TRP	CG-CD2-CE3	7.48	140.63	133.90	1	1
1	A	76	ARG	CG-CD-NE	-7.31	96.45	111.80	1	1
1	A	145	ARG	NE-CZ-NH2	-7.20	116.70	120.30	1	1
1	A	27	ARG	NE-CZ-NH2	-7.00	116.80	120.30	6	6
1	A	85	ARG	NH1-CZ-NH2	6.99	127.09	119.40	1	1
1	A	7	TYR	CB-CG-CD2	-6.78	116.93	121.00	1	1
1	A	52	ARG	NE-CZ-NH2	-6.69	116.95	120.30	16	5
1	A	159	GLN	C-N-CA	6.22	137.26	121.70	1	1
1	A	141	ARG	NE-CZ-NH1	6.19	123.40	120.30	1	1
1	A	72	ARG	NE-CZ-NH2	-6.05	117.27	120.30	3	9
1	A	124	TRP	CG-CD1-NE1	-5.97	104.12	110.10	1	1
1	A	124	TRP	CH2-CZ2-CE2	-5.97	111.43	117.40	1	1
1	A	21	GLN	CG-CD-NE2	5.91	130.87	116.70	1	1
1	A	16	CYS	CA-CB-SG	-5.83	103.50	114.00	1	1
1	A	35	PHE	CB-CG-CD2	-5.83	116.72	120.80	11	1
1	A	146	ARG	NE-CZ-NH2	-5.82	117.39	120.30	12	2
1	A	141	ARG	CD-NE-CZ	5.81	131.74	123.60	1	1
1	A	124	TRP	CB-CA-C	-5.79	98.82	110.40	1	1
1	A	124	TRP	N-CA-C	5.75	126.54	111.00	1	1
1	A	13	ILE	CA-CB-CG1	5.66	121.76	111.00	1	1
1	A	18	ARG	NH1-CZ-NH2	5.53	125.49	119.40	1	1
1	A	129	TYR	CB-CG-CD2	-5.52	117.69	121.00	20	1
1	A	77	TYR	CD1-CE1-CZ	-5.45	114.90	119.80	1	1
1	A	44	GLU	OE1-CD-OE2	-5.43	116.78	123.30	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	35	PHE	CB-CG-CD1	5.42	124.60	120.80	11	1
1	A	160	GLU	N-CA-C	5.38	125.53	111.00	1	1
1	A	141	ARG	NH1-CZ-NH2	5.38	125.32	119.40	1	1
1	A	75	LEU	CB-CG-CD2	5.34	120.07	111.00	1	1
1	A	124	TRP	CB-CG-CD2	5.31	133.50	126.60	1	1
1	A	89	LYS	CG-CD-CE	-5.24	96.19	111.90	1	1
1	A	125	ARG	CB-CG-CD	-5.22	98.04	111.60	1	1
1	A	160	GLU	CA-C-N	5.17	128.58	117.20	1	1
1	A	123	GLY	C-N-CA	5.04	134.29	121.70	1	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	85	ARG	Sidechain	7
1	A	144	TYR	Sidechain	4
1	A	129	TYR	Sidechain	4
1	A	77	TYR	Sidechain	3
1	A	141	ARG	Sidechain	1
1	A	121	PHE	Sidechain	1
1	A	160	GLU	Peptide	1
1	A	52	ARG	Sidechain	1
1	A	134	ARG	Sidechain	1
1	A	8	ARG	Sidechain	1
1	A	35	PHE	Sidechain	1
1	A	26	ARG	Sidechain	1

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1325	1308	1308	1±2
All	All	27825	27468	27468	23

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:124:TRP:CE3	1:A:124:TRP:O	0.67	2.48	1	1
1:A:124:TRP:CD2	1:A:124:TRP:O	0.62	2.52	1	1
1:A:86:TRP:HA	1:A:91:VAL:CG1	0.61	2.26	1	1
1:A:124:TRP:O	1:A:125:ARG:HB2	0.54	2.03	1	1
1:A:124:TRP:CE3	1:A:124:TRP:N	0.53	2.75	1	1
1:A:65:VAL:CG2	1:A:102:LEU:HD12	0.49	2.37	1	1
1:A:135:GLN:OE1	1:A:135:GLN:HA	0.49	2.07	1	1
1:A:104:LEU:HD22	1:A:107:GLY:HA2	0.47	1.87	21	1
1:A:24:TRP:CD1	1:A:33:TRP:CD1	0.45	3.05	17	2
1:A:134:ARG:HA	1:A:134:ARG:HE	0.45	1.72	1	1
1:A:124:TRP:O	1:A:125:ARG:CB	0.44	2.59	1	1
1:A:134:ARG:HH21	1:A:141:ARG:CZ	0.44	2.25	1	1
1:A:124:TRP:CG	1:A:125:ARG:N	0.43	2.86	7	2
1:A:24:TRP:CD1	1:A:33:TRP:CE3	0.43	3.07	18	2
1:A:25:ALA:HB3	1:A:121:PHE:CD1	0.42	2.49	3	1
1:A:51:TYR:CD1	1:A:62:ARG:HA	0.41	2.50	2	1
1:A:134:ARG:NH2	1:A:141:ARG:CZ	0.41	2.84	1	1
1:A:34:GLN:NE2	1:A:37:GLN:HE22	0.41	2.14	5	1
1:A:124:TRP:C	1:A:124:TRP:CD2	0.41	2.92	1	1
1:A:74:TRP:CG	1:A:96:LYS:HE3	0.41	2.51	6	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/164 (97%)	145±2 (91±1%)	11±2 (7±1%)	3±2 (2±1%)	11	50
All	All	3339/3444 (97%)	3035 (91%)	231 (7%)	73 (2%)	11	50

All 22 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	63	LYS	13
1	A	62	ARG	9
1	A	113	MET	7
1	A	6	GLY	6
1	A	18	ARG	6
1	A	37	GLN	5
1	A	124	TRP	4
1	A	36	PRO	4
1	A	89	LYS	4
1	A	106	SER	3
1	A	76	ARG	1
1	A	2	ILE	1
1	A	130	TRP	1
1	A	72	ARG	1
1	A	108	ASP	1
1	A	30	GLN	1
1	A	117	SER	1
1	A	120	GLU	1
1	A	15	ILE	1
1	A	129	TYR	1
1	A	21	GLN	1
1	A	86	TRP	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/146 (97%)	131±3 (93±2%)	10±3 (7±2%)	21	68
All	All	2961/3066 (97%)	2753 (93%)	208 (7%)	21	68

All 58 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	97	GLN	14
1	A	45	SER	13
1	A	115	THR	12
1	A	116	SER	11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	144	TYR	10
1	A	68	LEU	10
1	A	76	ARG	9
1	A	55	PHE	9
1	A	133	VAL	6
1	A	85	ARG	6
1	A	149	LYS	5
1	A	24	TRP	5
1	A	129	TYR	5
1	A	78	LYS	4
1	A	75	LEU	4
1	A	81	LYS	4
1	A	23	MET	4
1	A	122	ASP	4
1	A	95	GLN	4
1	A	35	PHE	4
1	A	58	VAL	4
1	A	108	ASP	4
1	A	8	ARG	3
1	A	136	VAL	3
1	A	37	GLN	3
1	A	28	PHE	3
1	A	73	ASN	3
1	A	48	GLN	3
1	A	112	ASN	2
1	A	101	LEU	2
1	A	33	TRP	2
1	A	124	TRP	2
1	A	125	ARG	2
1	A	134	ARG	2
1	A	11	VAL	2
1	A	88	THR	2
1	A	21	GLN	2
1	A	62	ARG	1
1	A	148	MET	1
1	A	47	GLU	1
1	A	145	ARG	1
1	A	14	VAL	1
1	A	27	ARG	1
1	A	151	PHE	1
1	A	32	SER	1
1	A	41	ASN	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	22	VAL	1
1	A	105	VAL	1
1	A	10	ASN	1
1	A	135	GLN	1
1	A	106	SER	1
1	A	91	VAL	1
1	A	17	ASN	1
1	A	3	ASP	1
1	A	159	GLN	1
1	A	57	GLU	1
1	A	137	VAL	1
1	A	114	GLN	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided