



Full wwPDB NMR Structure Validation Report ⓘ

Feb 16, 2018 – 04:05 am GMT

PDB ID : 1KMF
Title : NMR STRUCTURE OF HUMAN INSULIN MUTANT ILE-A2-ALLO-ILE,
HIS-B10-ASP, PRO-B28-LYS, LYS-B29-PRO, 15 STRUCTURES
Authors : Xu, B.; Hua, Q.X.; Nakagawa, S.H.; Jia, W.; Chu, Y.C.; Katsoyannis, P.G.;
Weiss, M.A.
Deposited on : 2001-12-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.7.3 (157068), CSD as539be (2018)
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk30686
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 43%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136279	12091
Ramachandran outliers	132675	10835
Sidechain outliers	132484	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	21	
2	B	30	

2 Ensemble composition and analysis

This entry contains 15 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:21, B:5-B:26 (41)	0.28	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 4, 6, 7, 11, 13
2	2, 8, 9, 10, 14, 15
Single-model clusters	3; 5; 12

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 781 atoms, of which 378 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Insulin.

Mol	Chain	Residues	Atoms						Trace
1	A	21	Total	C	H	N	O	S	0
			312	99	149	25	35	4	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	IIL	ILE	ENGINEERED	UNP P01308

- Molecule 2 is a protein called Insulin.

Mol	Chain	Residues	Atoms						Trace
2	B	30	Total	C	H	N	O	S	0
			469	156	229	38	44	2	

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	ASP	HIS	ENGINEERED	UNP P01308
B	28	LYS	PRO	ENGINEERED	UNP P01308
B	29	PRO	LYS	ENGINEERED	UNP P01308

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Insulin



- Molecule 2: Insulin



4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Insulin



- Molecule 2: Insulin




4.2.2 Score per residue for model 2

• Molecule 1: Insulin



• Molecule 2: Insulin




4.2.3 Score per residue for model 3

• Molecule 1: Insulin



• Molecule 2: Insulin




4.2.4 Score per residue for model 4

• Molecule 1: Insulin




• Molecule 2: Insulin




4.2.5 Score per residue for model 5

- Molecule 1: Insulin




- Molecule 2: Insulin




4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: Insulin



- Molecule 2: Insulin




4.2.7 Score per residue for model 7

- Molecule 1: Insulin




- Molecule 2: Insulin




4.2.8 Score per residue for model 8

• Molecule 1: Insulin



• Molecule 2: Insulin




4.2.9 Score per residue for model 9

• Molecule 1: Insulin



• Molecule 2: Insulin




4.2.10 Score per residue for model 10

• Molecule 1: Insulin




• Molecule 2: Insulin




4.2.11 Score per residue for model 11

- Molecule 1: Insulin




- Molecule 2: Insulin



4.2.12 Score per residue for model 12

- Molecule 1: Insulin




- Molecule 2: Insulin




4.2.13 Score per residue for model 13

- Molecule 1: Insulin




- Molecule 2: Insulin




4.2.14 Score per residue for model 14

- Molecule 1: Insulin



- Molecule 2: Insulin




4.2.15 Score per residue for model 15

- Molecule 1: Insulin



- Molecule 2: Insulin



5 Refinement protocol and experimental data overview

The models were refined using the following method: *Distance geometry and simulated annealing*.

Of the 50 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DGII	refinement	InsightII 2000
X-PLOR	refinement	3.85

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5246
Number of chemical shift lists	4
Total number of shifts	569
Number of shifts mapped to atoms	565
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	4
Assignment completeness (well-defined parts)	43%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IIL

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0.0±0.0	1.0±0.0
All	All	0	15

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	22	ARG	Sidechain	15

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	151	133	133	10±4
2	B	174	161	161	26±4
All	All	4875	4410	4410	445

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 48.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:14:ALA:O	2:B:18:VAL:HG23	0.90	1.67	8	14
2:B:15:LEU:HD13	2:B:24:PHE:CE2	0.81	2.11	1	3
1:A:5:GLN:O	1:A:8:THR:HG22	0.81	1.73	5	6
2:B:6:LEU:HD12	2:B:11:LEU:HA	0.80	1.51	11	7
2:B:5:HIS:C	2:B:6:LEU:HD22	0.75	2.02	8	7
1:A:3:VAL:HG21	2:B:26:TYR:CE2	0.75	2.17	11	3
2:B:12:VAL:HG23	2:B:26:TYR:OH	0.74	1.83	15	1
2:B:6:LEU:HD12	2:B:10:ASP:C	0.72	2.05	6	6
1:A:16:LEU:HD13	2:B:14:ALA:HB1	0.71	1.61	6	8
1:A:16:LEU:CD1	2:B:14:ALA:HB1	0.68	2.18	6	7
2:B:6:LEU:HD12	2:B:11:LEU:CA	0.67	2.19	11	6
1:A:3:VAL:HG22	2:B:26:TYR:CE1	0.65	2.27	12	1
1:A:16:LEU:HD21	2:B:11:LEU:CD2	0.63	2.24	8	7
1:A:13:LEU:HD23	1:A:13:LEU:O	0.63	1.92	5	1
1:A:3:VAL:HG22	2:B:26:TYR:CZ	0.62	2.29	4	9
2:B:6:LEU:HD23	2:B:11:LEU:HD23	0.62	1.71	10	2
2:B:15:LEU:CD1	2:B:24:PHE:CE2	0.62	2.82	1	2
1:A:16:LEU:HD21	2:B:11:LEU:HD21	0.61	1.71	13	8
2:B:24:PHE:CD1	2:B:25:PHE:N	0.61	2.69	5	2
2:B:23:GLY:O	2:B:24:PHE:CD2	0.61	2.54	5	2
2:B:12:VAL:CG2	2:B:26:TYR:CE2	0.60	2.85	12	1
2:B:5:HIS:O	2:B:6:LEU:HD12	0.60	1.97	13	2
2:B:15:LEU:HD13	2:B:24:PHE:CD2	0.60	2.32	1	3
2:B:25:PHE:CD1	2:B:25:PHE:N	0.59	2.70	5	2
2:B:6:LEU:N	2:B:6:LEU:HD22	0.58	2.13	1	5
1:A:9:SER:O	2:B:5:HIS:CE1	0.58	2.56	10	3
2:B:25:PHE:CG	2:B:25:PHE:O	0.58	2.56	6	4
2:B:5:HIS:O	2:B:6:LEU:HD22	0.58	1.98	8	2
2:B:6:LEU:HD12	2:B:11:LEU:N	0.58	2.13	6	2
2:B:7:CYS:O	2:B:11:LEU:HB2	0.56	2.00	13	12
2:B:5:HIS:C	2:B:6:LEU:HD12	0.56	2.21	10	2
2:B:7:CYS:O	2:B:8:GLY:C	0.56	2.44	13	2
2:B:25:PHE:CD2	2:B:25:PHE:O	0.56	2.59	6	4
1:A:13:LEU:HA	1:A:16:LEU:HD12	0.56	1.76	15	2
1:A:16:LEU:HB3	2:B:18:VAL:HG21	0.55	1.77	7	2
2:B:6:LEU:N	2:B:6:LEU:CD2	0.55	2.68	1	4
2:B:6:LEU:HD13	2:B:10:ASP:HB3	0.55	1.78	6	3
2:B:6:LEU:N	2:B:6:LEU:CD1	0.55	2.69	10	2
1:A:10:ILE:O	1:A:11:CYS:CB	0.54	2.55	8	2
1:A:6:CYS:HA	1:A:11:CYS:CB	0.54	2.32	15	1
1:A:3:VAL:CG2	2:B:26:TYR:CE2	0.54	2.90	9	3
2:B:24:PHE:CD1	2:B:24:PHE:C	0.54	2.80	3	3

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:14:ALA:C	2:B:18:VAL:HG23	0.54	2.23	4	4
2:B:7:CYS:O	2:B:11:LEU:CB	0.54	2.56	7	9
2:B:25:PHE:CD1	2:B:25:PHE:O	0.54	2.61	5	1
2:B:14:ALA:O	2:B:18:VAL:N	0.53	2.41	5	13
2:B:15:LEU:HB3	2:B:24:PHE:CD2	0.53	2.38	15	4
2:B:9:SER:O	2:B:13:GLU:N	0.53	2.41	9	6
2:B:7:CYS:O	2:B:11:LEU:N	0.53	2.39	13	2
2:B:6:LEU:HD12	2:B:10:ASP:O	0.53	2.04	5	4
2:B:20:GLY:O	2:B:22:ARG:N	0.53	2.42	5	1
2:B:6:LEU:O	2:B:7:CYS:SG	0.52	2.67	3	11
2:B:19:CYS:SG	2:B:24:PHE:CB	0.52	2.98	3	1
2:B:19:CYS:CB	2:B:23:GLY:O	0.52	2.57	15	4
2:B:25:PHE:C	2:B:25:PHE:CD1	0.52	2.80	12	1
2:B:21:GLU:OE2	2:B:22:ARG:CD	0.52	2.57	8	1
1:A:16:LEU:HD13	2:B:14:ALA:CB	0.52	2.33	6	4
1:A:3:VAL:HG21	2:B:26:TYR:CD2	0.52	2.40	11	1
2:B:25:PHE:O	2:B:26:TYR:O	0.52	2.28	14	3
1:A:16:LEU:HD22	2:B:15:LEU:HG	0.52	1.81	12	1
1:A:21:ASN:OD1	2:B:22:ARG:CB	0.52	2.58	3	1
2:B:7:CYS:O	2:B:8:GLY:O	0.52	2.28	13	2
1:A:3:VAL:HG22	2:B:26:TYR:CE2	0.51	2.40	9	1
1:A:3:VAL:O	1:A:7:CYS:HB3	0.51	2.06	12	9
1:A:9:SER:O	2:B:5:HIS:NE2	0.51	2.43	15	1
2:B:23:GLY:O	2:B:24:PHE:O	0.51	2.29	5	2
1:A:7:CYS:O	2:B:5:HIS:NE2	0.50	2.43	13	2
2:B:19:CYS:O	2:B:21:GLU:N	0.50	2.44	1	2
2:B:18:VAL:HG12	2:B:19:CYS:SG	0.50	2.46	6	1
1:A:3:VAL:HG13	2:B:26:TYR:OH	0.49	2.07	1	1
1:A:7:CYS:O	2:B:5:HIS:CE1	0.49	2.66	3	1
1:A:16:LEU:CD1	2:B:14:ALA:CB	0.49	2.90	10	3
1:A:6:CYS:O	2:B:5:HIS:CE1	0.49	2.65	10	2
2:B:24:PHE:CG	2:B:24:PHE:O	0.49	2.64	3	1
1:A:7:CYS:HA	2:B:6:LEU:O	0.49	2.08	5	11
2:B:12:VAL:CG2	2:B:26:TYR:CE1	0.48	2.96	7	1
2:B:24:PHE:C	2:B:24:PHE:CD1	0.48	2.86	4	1
1:A:10:ILE:O	1:A:11:CYS:SG	0.48	2.72	3	1
1:A:6:CYS:O	1:A:9:SER:O	0.48	2.32	9	3
1:A:3:VAL:O	1:A:7:CYS:CB	0.48	2.62	12	6
2:B:9:SER:O	2:B:12:VAL:N	0.47	2.47	13	1
1:A:20:CYS:O	1:A:21:ASN:OD1	0.47	2.33	3	2
2:B:8:GLY:O	2:B:9:SER:C	0.47	2.53	2	8

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:8:THR:HG23	1:A:9:SER:OG	0.47	2.09	2	1
2:B:15:LEU:HB3	2:B:24:PHE:CG	0.47	2.45	15	3
1:A:10:ILE:HG23	1:A:11:CYS:N	0.47	2.25	13	1
1:A:6:CYS:O	2:B:6:LEU:O	0.47	2.33	8	4
2:B:19:CYS:SG	2:B:23:GLY:O	0.46	2.74	2	2
1:A:21:ASN:ND2	2:B:22:ARG:O	0.46	2.49	8	2
1:A:10:ILE:CG2	1:A:11:CYS:N	0.46	2.78	6	1
2:B:24:PHE:CD1	2:B:24:PHE:O	0.46	2.68	3	1
2:B:7:CYS:O	2:B:26:TYR:OH	0.46	2.33	4	1
2:B:8:GLY:O	2:B:10:ASP:N	0.46	2.49	2	6
1:A:6:CYS:HB3	2:B:6:LEU:HB2	0.45	1.86	6	2
2:B:6:LEU:O	2:B:7:CYS:CB	0.45	2.64	3	3
2:B:6:LEU:CD1	2:B:10:ASP:C	0.45	2.84	4	3
1:A:10:ILE:HG22	1:A:11:CYS:H	0.45	1.70	4	1
1:A:10:ILE:O	1:A:11:CYS:HB2	0.45	2.11	8	2
2:B:6:LEU:HD22	2:B:6:LEU:N	0.45	2.24	12	1
2:B:22:ARG:O	2:B:23:GLY:C	0.45	2.53	9	1
2:B:12:VAL:HG23	2:B:26:TYR:CE1	0.45	2.47	4	1
1:A:5:GLN:O	1:A:5:GLN:OE1	0.45	2.34	9	1
2:B:6:LEU:CD2	2:B:6:LEU:N	0.45	2.79	12	1
2:B:25:PHE:O	2:B:26:TYR:C	0.45	2.55	3	4
1:A:13:LEU:HD23	1:A:13:LEU:N	0.45	2.27	4	1
2:B:25:PHE:O	2:B:25:PHE:CG	0.45	2.69	2	2
2:B:23:GLY:O	2:B:24:PHE:HB3	0.45	2.10	6	13
2:B:12:VAL:HG22	2:B:26:TYR:CE1	0.45	2.46	7	1
2:B:21:GLU:O	2:B:23:GLY:N	0.45	2.49	6	3
2:B:12:VAL:HG23	2:B:26:TYR:CZ	0.45	2.46	6	2
2:B:6:LEU:HB3	2:B:11:LEU:CG	0.45	2.41	8	1
1:A:3:VAL:CG2	2:B:26:TYR:CZ	0.45	3.00	8	2
2:B:19:CYS:HB2	2:B:23:GLY:O	0.45	2.12	9	3
2:B:6:LEU:N	2:B:6:LEU:HD12	0.45	2.25	10	1
1:A:10:ILE:O	1:A:11:CYS:C	0.44	2.55	3	1
2:B:21:GLU:O	2:B:22:ARG:C	0.44	2.54	14	6
2:B:14:ALA:O	2:B:18:VAL:CG2	0.44	2.57	14	1
2:B:5:HIS:CE1	2:B:7:CYS:SG	0.44	3.10	13	1
2:B:6:LEU:CD1	2:B:10:ASP:O	0.44	2.66	11	4
2:B:12:VAL:O	2:B:13:GLU:C	0.44	2.55	1	4
2:B:19:CYS:SG	2:B:24:PHE:HB3	0.44	2.53	3	2
1:A:3:VAL:HA	2:B:11:LEU:HD12	0.44	1.89	11	2
1:A:15:GLN:O	1:A:19:TYR:CD2	0.44	2.71	11	1
2:B:19:CYS:SG	2:B:24:PHE:HB2	0.43	2.52	3	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:15:LEU:CB	2:B:24:PHE:CE2	0.43	3.01	1	1
1:A:4:GLU:O	1:A:8:THR:CG2	0.43	2.67	11	3
1:A:13:LEU:HA	1:A:16:LEU:HB2	0.43	1.91	4	6
2:B:25:PHE:N	2:B:25:PHE:CD1	0.43	2.85	1	1
2:B:12:VAL:HG23	2:B:26:TYR:CE2	0.43	2.49	12	1
2:B:6:LEU:HB3	2:B:11:LEU:HG	0.42	1.90	8	1
1:A:8:THR:O	1:A:9:SER:OG	0.42	2.36	5	1
2:B:9:SER:O	2:B:10:ASP:C	0.42	2.57	13	1
2:B:8:GLY:C	2:B:10:ASP:N	0.42	2.72	2	3
2:B:24:PHE:O	2:B:24:PHE:CD1	0.42	2.73	14	1
2:B:11:LEU:HA	2:B:11:LEU:HD23	0.42	1.75	2	1
1:A:10:ILE:O	1:A:11:CYS:O	0.42	2.37	15	1
2:B:12:VAL:CG2	2:B:26:TYR:OH	0.42	2.61	15	1
2:B:15:LEU:CB	2:B:24:PHE:CD2	0.42	3.03	9	2
1:A:9:SER:O	2:B:5:HIS:CD2	0.42	2.73	3	1
1:A:7:CYS:O	2:B:7:CYS:SG	0.42	2.78	3	1
2:B:10:ASP:OD1	2:B:10:ASP:N	0.42	2.52	12	1
2:B:14:ALA:O	2:B:17:LEU:N	0.42	2.50	9	1
1:A:10:ILE:O	1:A:11:CYS:HB3	0.42	2.15	15	1
1:A:9:SER:O	2:B:5:HIS:ND1	0.42	2.53	10	1
2:B:14:ALA:O	2:B:15:LEU:C	0.41	2.57	4	4
1:A:7:CYS:C	2:B:5:HIS:CE1	0.41	2.93	1	1
2:B:15:LEU:HB2	2:B:24:PHE:CZ	0.41	2.50	1	1
2:B:20:GLY:O	2:B:21:GLU:HB2	0.41	2.16	5	1
2:B:7:CYS:O	2:B:11:LEU:HB3	0.41	2.15	7	1
1:A:9:SER:C	2:B:5:HIS:NE2	0.41	2.74	15	1
2:B:12:VAL:O	2:B:14:ALA:N	0.41	2.53	1	2
2:B:12:VAL:C	2:B:14:ALA:N	0.41	2.74	1	2
2:B:9:SER:O	2:B:13:GLU:HB2	0.41	2.16	2	1
2:B:20:GLY:O	2:B:21:GLU:C	0.41	2.57	8	1
1:A:20:CYS:O	1:A:21:ASN:ND2	0.41	2.53	12	1
1:A:21:ASN:ND2	2:B:23:GLY:C	0.40	2.74	14	1
1:A:21:ASN:OD1	2:B:22:ARG:HB3	0.40	2.16	3	1
2:B:6:LEU:HD13	2:B:6:LEU:HA	0.40	1.74	11	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	18/21 (86%)	15±1 (86±6%)	2±1 (12±6%)	0±1 (2±3%)	11	50
2	B	22/30 (73%)	14±2 (62±7%)	7±2 (30±8%)	2±1 (8±4%)	2	13
All	All	600/765 (78%)	435 (72%)	131 (22%)	34 (6%)	4	22

All 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	7	CYS	12
2	B	8	GLY	7
2	B	26	TYR	4
1	A	11	CYS	3
1	A	20	CYS	2
2	B	21	GLU	2
2	B	24	PHE	2
2	B	20	GLY	1
1	A	10	ILE	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	19/19 (100%)	15±1 (81±8%)	4±1 (19±8%)	5	37
2	B	18/26 (69%)	12±1 (67±7%)	6±1 (33±7%)	1	12
All	All	555/675 (82%)	414 (75%)	141 (25%)	2	24

All 23 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	11	LEU	15
2	B	16	TYR	15
1	A	13	LEU	13
2	B	26	TYR	12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	12	SER	11
2	B	9	SER	10
2	B	17	LEU	10
1	A	11	CYS	8
2	B	22	ARG	7
1	A	14	TYR	6
2	B	24	PHE	5
2	B	25	PHE	4
1	A	17	GLU	4
2	B	7	CYS	4
2	B	21	GLU	3
1	A	18	ASN	3
1	A	5	GLN	3
2	B	10	ASP	2
1	A	10	ILE	2
1	A	6	CYS	1
1	A	4	GLU	1
2	B	13	GLU	1
1	A	9	SER	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	IIL	A	2	1	7,7,8	0.99±0.03	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles

that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	IIL	A	2	1	6,8,10	1.29±0.03	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	IIL	A	2	1	-	0±0,6,8,10	0±0,0,0,0

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 43% for the well-defined parts and 42% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5246

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	123
Number of shifts mapped to atoms	121
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	2
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atoms found in structure. All 2 occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	2	IIL	HG2	0.64	0.02	1
A	2	IIL	HD1	0.51	0.02	1

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 20%, i.e. 97 atoms were assigned a chemical shift out of a possible 481. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	37/205 (18%)	37/82 (45%)	0/82 (0%)	0/41 (0%)
Sidechain	56/219 (26%)	56/128 (44%)	0/84 (0%)	0/7 (0%)
Aromatic	4/57 (7%)	4/30 (13%)	0/26 (0%)	0/1 (0%)
Overall	97/481 (20%)	97/240 (40%)	0/192 (0%)	0/49 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 16%, i.e. 97 atoms were assigned a chemical shift out of a possible 589. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	37/248 (15%)	37/99 (37%)	0/100 (0%)	0/49 (0%)
Sidechain	56/275 (20%)	56/161 (35%)	0/104 (0%)	0/10 (0%)
Aromatic	4/66 (6%)	4/35 (11%)	0/30 (0%)	0/1 (0%)
Overall	97/589 (16%)	97/295 (33%)	0/234 (0%)	0/60 (0%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: BMRB entry 5246

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	181
Number of shifts mapped to atoms	181
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.2.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 22%, i.e. 108 atoms were assigned a chemical shift out of a possible 481. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	44/205 (21%)	44/82 (54%)	0/82 (0%)	0/41 (0%)
Sidechain	56/219 (26%)	56/128 (44%)	0/84 (0%)	0/7 (0%)
Aromatic	8/57 (14%)	8/30 (27%)	0/26 (0%)	0/1 (0%)
Overall	108/481 (22%)	108/240 (45%)	0/192 (0%)	0/49 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 25%, i.e. 146 atoms were assigned a chemical shift out of a possible 589. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	58/248 (23%)	58/99 (59%)	0/100 (0%)	0/49 (0%)
Sidechain	78/275 (28%)	78/161 (48%)	0/104 (0%)	0/10 (0%)
Aromatic	10/66 (15%)	10/35 (29%)	0/30 (0%)	0/1 (0%)
Overall	146/589 (25%)	146/295 (49%)	0/234 (0%)	0/60 (0%)


7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:



7.3 Chemical shift list 3

File name: BMRB entry 5246

Chemical shift list name: *assigned_chem_shift_list_3*

7.3.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	104
Number of shifts mapped to atoms	102
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	2
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atoms found in structure. All 2 occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	2	IIL	HG2	0.68	0.02	1
A	2	IIL	HD1	0.5	0.02	1

7.3.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.3.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 16%, i.e. 79 atoms were assigned a chemical shift out of a possible 481. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	27/205 (13%)	27/82 (33%)	0/82 (0%)	0/41 (0%)
Sidechain	48/219 (22%)	48/128 (38%)	0/84 (0%)	0/7 (0%)
Aromatic	4/57 (7%)	4/30 (13%)	0/26 (0%)	0/1 (0%)
Overall	79/481 (16%)	79/240 (33%)	0/192 (0%)	0/49 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 13%, i.e. 79 atoms were assigned a chemical shift out of a possible 589. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	27/248 (11%)	27/99 (27%)	0/100 (0%)	0/49 (0%)
Sidechain	48/275 (17%)	48/161 (30%)	0/104 (0%)	0/10 (0%)
Aromatic	4/66 (6%)	4/35 (11%)	0/30 (0%)	0/1 (0%)
Overall	79/589 (13%)	79/295 (27%)	0/234 (0%)	0/60 (0%)

7.3.4 Statistically unusual chemical shifts [i](#)


There are no statistically unusual chemical shifts.

7.3.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication

of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.4 Chemical shift list 4

File name: BMRB entry 5246

Chemical shift list name: *assigned_chem_shift_list_4*

7.4.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	161
Number of shifts mapped to atoms	161
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.4.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.4.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 21%, i.e. 102 atoms were assigned a chemical shift out of a possible 481. 0 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	39/205 (19%)	39/82 (48%)	0/82 (0%)	0/41 (0%)
Sidechain	55/219 (25%)	55/128 (43%)	0/84 (0%)	0/7 (0%)
Aromatic	8/57 (14%)	8/30 (27%)	0/26 (0%)	0/1 (0%)
Overall	102/481 (21%)	102/240 (42%)	0/192 (0%)	0/49 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 22%, i.e. 129 atoms were assigned a chemical shift out of a possible 589. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	48/248 (19%)	48/99 (48%)	0/100 (0%)	0/49 (0%)
Sidechain	73/275 (27%)	73/161 (45%)	0/104 (0%)	0/10 (0%)
Aromatic	8/66 (12%)	8/35 (23%)	0/30 (0%)	0/1 (0%)
Overall	129/589 (22%)	129/295 (44%)	0/234 (0%)	0/60 (0%)

7.4.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.4.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:

