



wwPDB EM Model Validation Summary Report ⓘ

Apr 9, 2020 – 06:43 AM EDT

PDB ID : 6KMF
EMDB ID : EMD-0724
Title : FimA type V pilus from P.gingivalis
Authors : Shibata, S.; Shoji, M.; Matsunami, H.; Matthews, M.; Imada, K.; Nakayama, K.; Wolf, M.
Deposited on : 2019-07-31
Resolution : 3.60 Å(reported)
Based on initial model : 6JZJ

This is a wwPDB EM Model Validation Summary Report for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.10.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	136327	1886
Ramachandran outliers	132723	1663
Sidechain outliers	132532	1531

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	337	 78% 15% 6%
1	B	337	 90% 10%
1	C	337	 86% 14%
1	D	337	 5% • 94%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 7704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Major fimbrium subunit FimA type-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	316	Total	C	N	O	S	0	0
			2403	1515	402	478	8		
1	B	337	Total	C	N	O	S	0	0
			2568	1620	431	508	9		
1	C	337	Total	C	N	O	S	0	0
			2568	1620	431	508	9		
1	D	21	Total	C	N	O	S	0	0
			165	105	29	30	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major fimbrium subunit FimA type-1



- Molecule 1: Major fimbrium subunit FimA type-1



- Molecule 1: Major fimbrium subunit FimA type-1



- Molecule 1: Major fimbrium subunit FimA type-1



ILE	THR	GLY	THR	LYS
THR	GLY	ASP	ASN	ARG
PRO	LEU	ALA	GLY	HIS
GLY	ALA	ALA	ALA	ALA
THR	GLY	GLY	TYR	ARG
ASN	ASN	ALA	THR	MET
ASN	ASN	ASP	PRO	ALA
PRO	LEU	LEU	ALA	PHE
GLY	ALA	ALA	ASN	THR
ASN	ALA	ALA	TYR	GLU
PRO	ALA	ALA	ALA	ILE
ILE	GLN	GLN	ASN	LYS
THR	ALA	ALA	VAL	GLN
GLY	ALA	ALA	PRO	GLN
SER	ASN	TRP	TRP	MET
A363	VAL	VAL	LEU	SER
V367	ASP	ASP	ARG	ALA
Q368	ALA	ALA	ASN	ALA
	GLU	GLU	TYR	ASP
Q379	GLY	GLY	VAL	ASN
	LYS	THR	ALA	ILE
W383	THR	TYR	PRO	TYR
	TYR	ALA	ALA	PHE
	PRO	VAL	ASP	VAL
	VAL	ALA	ALA	PRO
	LEU	LEU	PRO	GLU
	VAL	VAL	GLN	LYS
	ASN	ASN	GLY	ILE
	THR	PHE	PHE	TYR
	ASN	ASN	TYR	GLY
	SER	SER	VAL	LEU
	ASN	SER	ASN	ASN
	ASN	ASN	ALA	LEU
	TYR	TYR	GLY	PHE
	THR	THR	GLY	GLY
	PRO	LYS	ILE	THR
	ASN	LYS	HIS	LEU
	ASN	ASN	PRO	VAL
	ILE	ILE	THR	ASN
	GLU	ARG	LEU	ALA
	ASN	ASN	VAL	ASP
	HIS	HIS	VAL	ALA
	LYS	LYS	VAL	ASN
	TYR	TYR	TYR	TYR
	ASP	ASP	GLY	THR
	ILE	ILE	LEU	GLY
	LYS	LYS	GLN	SER
	LEU	LEU	LYS	LEU
	THR	THR	ASN	THR

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	61728	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TALOS ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	46	Depositor
Minimum defocus (nm)	-1500	Depositor
Maximum defocus (nm)	-2500	Depositor
Magnification	125000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2449	0.58	0/3332
1	B	0.34	0/2619	0.55	0/3567
1	C	0.36	0/2619	0.58	0/3567
1	D	0.34	0/169	0.49	0/232
All	All	0.35	0/7856	0.57	0/10698

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2403	0	2360	34	0
1	B	2568	0	2516	23	0
1	C	2568	0	2516	29	0
1	D	165	0	155	4	0
All	All	7704	0	7547	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:ARG:HE	1:B:340:LYS:HG2	1.58	0.69
1:A:62:VAL:HG21	1:B:367:VAL:HG21	1.77	0.65
1:B:222:THR:HG22	1:B:224:SER:H	1.62	0.64
1:A:277:LEU:HD23	1:A:343:ILE:HD11	1.80	0.64
1:A:146:ASN:ND2	1:A:157:ASN:OD1	2.32	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/337 (93%)	271 (86%)	43 (14%)	0	100	100
1	B	335/337 (99%)	291 (87%)	44 (13%)	0	100	100
1	C	335/337 (99%)	288 (86%)	47 (14%)	0	100	100
1	D	19/337 (6%)	19 (100%)	0	0	100	100
All	All	1003/1348 (74%)	869 (87%)	134 (13%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/268 (94%)	250 (100%)	1 (0%)	92	97
1	B	268/268 (100%)	268 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	268/268 (100%)	267 (100%)	1 (0%)	92	97
1	D	17/268 (6%)	17 (100%)	0	100	100
All	All	804/1072 (75%)	802 (100%)	2 (0%)	94	98

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	VAL
1	C	176	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	157	ASN
1	C	91	GLN
1	C	236	ASN
1	B	146	ASN
1	C	205	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.