



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 26, 2018 – 06:50 AM EDT

PDB ID : 3KPJ  
Title : Crystal Structure of hPNMT in Complex AdoHcy and Bound Phosphate  
Authors : Drinkwater, N.; Martin, J.L.  
Deposited on : 2009-11-16  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031021  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031021




# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4155 (2.50-2.50)
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div>3%</div> <div>69%</div> <div>20%</div> <div>11%</div> </div>
1	B	289	<div> <div>7%</div> <div>75%</div> <div>17%</div> <div>7%</div> </div>



## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4255 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylethanolamine N-methyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	2	0	0
			2016	1280	358	369	9			
1	B	268	Total	C	N	O	S	0	0	0
			2083	1320	370	384	9			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	283	GLU	-	EXPRESSION TAG	UNP P11086
A	284	HIS	-	EXPRESSION TAG	UNP P11086
A	285	HIS	-	EXPRESSION TAG	UNP P11086
A	286	HIS	-	EXPRESSION TAG	UNP P11086
A	287	HIS	-	EXPRESSION TAG	UNP P11086
A	288	HIS	-	EXPRESSION TAG	UNP P11086
A	289	HIS	-	EXPRESSION TAG	UNP P11086
B	283	GLU	-	EXPRESSION TAG	UNP P11086
B	284	HIS	-	EXPRESSION TAG	UNP P11086
B	285	HIS	-	EXPRESSION TAG	UNP P11086
B	286	HIS	-	EXPRESSION TAG	UNP P11086
B	287	HIS	-	EXPRESSION TAG	UNP P11086
B	288	HIS	-	EXPRESSION TAG	UNP P11086
B	289	HIS	-	EXPRESSION TAG	UNP P11086

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>5</sub>S).





- 
- [O-]P(=O)([O-])[O-]

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 5	O 4	P 1	0	0
3	B	1	Total 5	O 4	P 1	0	0





- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	47	Total 47	O 47	0	0
4	B	47	Total 47	O 47	0	0





- Molecule 1: Phenylethanolamine N-methyltransferase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.42Å 94.42Å 186.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.38 – 2.50 33.38 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.4 (33.38-2.50) 96.5 (33.38-2.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.51Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.211 , 0.267 0.205 , 0.262	Depositor DCC
$R_{free}$ test set	1472 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	66.0	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4255	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/2067	0.54	0/2813
1	B	0.42	0/2136	0.58	0/2908
All	All	0.40	0/4203	0.56	0/5721

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2016	0	1983	39	0
1	B	2083	0	2045	32	0
2	A	26	0	19	2	0
2	B	26	0	19	1	0
3	A	5	0	0	1	0
3	B	5	0	0	1	0
4	A	47	0	0	3	0
4	B	47	0	0	2	0
All	All	4255	0	4066	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.



All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:261:HIS:O	1:B:262:LEU:HB2	1.83	0.78
1:B:94:PHE:O	1:B:150:ARG:HD3	1.93	0.69
1:B:195:GLN:HE22	1:B:242:ALA:HA	1.61	0.65
1:B:75:LEU:HD11	1:B:179:VAL:HG23	1.84	0.58
1:A:91:CYS:O	1:A:150:ARG:HD2	2.02	0.58
1:A:95:GLU:HA	1:A:150:ARG:HD3	1.86	0.58
1:B:83:THR:HA	1:B:123:TRP:CZ2	2.38	0.58
1:B:207:PRO:HB3	1:B:281:GLY:HA3	1.85	0.57
1:B:267:ASP:OD2	1:B:269:VAL:HG12	2.05	0.57
1:A:263:GLN:HG2	4:A:300:HOH:O	2.04	0.57
1:A:56:TRP:CE2	1:A:256:TYR:HB2	2.40	0.57
1:A:182:PHE:CZ	3:A:290:PO4:O2	2.58	0.56
1:A:178:LEU:HD11	1:A:205:LEU:HB2	1.88	0.55
1:B:220:SER:HA	1:B:231:VAL:O	2.07	0.55
1:B:221:TRP:HA	1:B:229:LEU:O	2.07	0.55
1:A:237:GLU:O	1:A:241:GLU:HG3	2.08	0.54
1:A:125:MET:HB2	4:A:296:HOH:O	2.08	0.53
1:A:115:GLN:HB2	1:A:117:GLU:HG3	1.92	0.52
1:A:74:THR:HG22	1:A:96:ASP:HB3	1.92	0.52
1:A:130:ALA:O	1:A:134:GLU:HG3	2.10	0.52
1:B:216:ALA:HB1	1:B:269:VAL:HG23	1.92	0.52
1:A:141:GLN:O	1:A:145:ARG:HB2	2.10	0.52
1:A:162:PRO:HA	1:A:200:HIS:CD2	2.45	0.50
1:A:261:HIS:HD2	4:A:298:HOH:O	1.93	0.50
1:A:53:VAL:HG12	1:A:262:LEU:HD13	1.95	0.49
1:B:59:ARG:O	1:B:63:GLN:HG3	2.13	0.49
1:A:100:THR:HA	1:A:155:LEU:O	2.12	0.49
1:A:182:PHE:HA	1:A:185:GLU:OE2	2.13	0.48
1:B:219:GLU:HB2	1:B:269:VAL:HA	1.96	0.48
1:B:261:HIS:O	1:B:262:LEU:CB	2.59	0.47
1:A:117:GLU:HB3	1:A:118:PRO:HD2	1.97	0.47
1:B:76:ILE:HG12	1:B:98:THR:HB	1.97	0.47
1:B:182:PHE:CZ	3:B:290:PO4:O3	2.68	0.46
1:A:91:CYS:HA	1:A:97:ILE:HD11	1.98	0.46
1:B:261:HIS:ND1	4:B:311:HOH:O	2.36	0.46
1:B:118:PRO:HD2	4:B:323:HOH:O	2.15	0.46
1:B:44:ARG:HG2	1:B:262:LEU:HD22	1.98	0.46
1:A:102:PHE:HB2	2:A:3001:SAH:C4	2.46	0.46
1:A:49:ASN:HA	1:A:50:PRO:HD3	1.70	0.46
1:B:14:PRO:HB2	1:B:15:ASP:H	1.53	0.45

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:LEU:HD12	1:B:177:ALA:HB3	1.99	0.45
1:B:127:SER:HB3	1:B:140:TRP:HB2	1.98	0.45
1:A:178:LEU:CD1	1:A:205:LEU:HB2	2.46	0.45
1:B:159:VAL:HG22	2:B:3002:SAH:N1	2.32	0.44
1:A:188:SER:HA	1:A:189:PRO:HD2	1.84	0.44
1:B:49:ASN:OD1	1:B:50:PRO:HD2	2.18	0.44
1:A:187:VAL:HG11	2:A:3001:SAH:C5	2.48	0.43
1:B:217:LEU:HD21	1:B:273:PHE:HB3	2.00	0.43
1:A:267:ASP:OD1	1:A:269:VAL:HG12	2.17	0.43
1:B:218:GLU:OE1	1:B:218:GLU:HA	2.18	0.43
1:A:101:ASP:O	1:A:156:PRO:HA	2.17	0.43
1:A:82:PRO:HG3	1:A:106:ASN:ND2	2.34	0.42
1:B:220:SER:OG	1:B:268:ASP:OD1	2.34	0.42
1:A:73:ARG:N	1:A:176:ASP:OD1	2.48	0.42
1:A:216:ALA:HB1	1:A:269:VAL:HG23	2.01	0.42
1:A:75:LEU:HD23	1:A:76:ILE:N	2.35	0.42
1:B:188:SER:HA	1:B:189:PRO:HD3	1.85	0.42
1:A:206:ARG:HG2	1:A:207:PRO:N	2.33	0.42
1:B:189:PRO:HD2	1:B:193:SER:CB	2.50	0.42
1:A:169:SER:HA	1:A:170:PRO:HD3	1.89	0.42
1:A:195:GLN:HE22	1:A:242:ALA:HA	1.85	0.41
1:A:222:TYR:CB	1:A:267:ASP:HB3	2.50	0.41
1:B:254:ARG:HE	1:B:254:ARG:HB3	1.65	0.41
1:B:262:LEU:HD23	1:B:262:LEU:HA	1.89	0.41
1:A:115:GLN:HB2	1:A:117:GLU:CG	2.51	0.41
1:B:152:LYS:O	1:B:153:ARG:HB3	2.20	0.41
1:A:53:VAL:HG12	1:A:262:LEU:CD1	2.51	0.41
1:A:64:THR:O	1:A:67:THR:HG23	2.20	0.41
1:B:17:ALA:HB3	1:B:18:PRO:HD3	2.04	0.40
1:A:191:LEU:HD23	1:A:191:LEU:HA	1.72	0.40
1:A:161:GLN:HA	1:A:162:PRO:HD3	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/289 (88%)	246 (96%)	9 (4%)	0	100	100
1	B	266/289 (92%)	251 (94%)	15 (6%)	0	100	100
All	All	521/578 (90%)	497 (95%)	24 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/233 (91%)	209 (99%)	3 (1%)	69	88
1	B	218/233 (94%)	214 (98%)	4 (2%)	62	84
All	All	430/466 (92%)	423 (98%)	7 (2%)	65	86

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	VAL
1	A	191	LEU
1	A	206	ARG
1	B	29	ARG
1	B	128	GLN
1	B	229	LEU
1	B	245	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	106	ASN
1	A	195	GLN

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	39	ASN
1	B	93	HIS
1	B	106	ASN
1	B	128	GLN
1	B	195	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	A	290	-	4,4,4	0.65	0	6,6,6	0.59	0
2	SAH	A	3001	-	20,28,28	1.17	2 (10%)	19,40,40	2.53	3 (15%)
3	PO4	B	290	-	4,4,4	0.80	0	6,6,6	0.74	0
2	SAH	B	3002	-	20,28,28	1.17	2 (10%)	19,40,40	2.37	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	290	-	-	0/0/0/0	0/0/0/0
2	SAH	A	3001	-	-	0/7/31/31	0/3/3/3
3	PO4	B	290	-	-	0/0/0/0	0/0/0/0
2	SAH	B	3002	-	-	0/7/31/31	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3002	SAH	C2-N1	2.36	1.38	1.33
2	A	3001	SAH	C2-N1	2.43	1.38	1.33
2	A	3001	SAH	C2-N3	3.85	1.38	1.32
2	B	3002	SAH	C2-N3	3.97	1.38	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3001	SAH	N3-C2-N1	-10.02	120.29	128.86
2	B	3002	SAH	N3-C2-N1	-9.20	120.99	128.86
2	B	3002	SAH	C4-C5-N7	-2.61	106.88	109.41
2	A	3001	SAH	C5'-SD-CG	-2.44	94.94	102.27
2	A	3001	SAH	C4'-O4'-C1'	-2.29	107.44	109.83
2	B	3002	SAH	C5'-SD-CG	-2.18	95.73	102.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	290	PO4	1	0
2	A	3001	SAH	2	0
3	B	290	PO4	1	0
2	B	3002	SAH	1	0

## 5.7 Other polymers

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	257/289 (88%)	-0.08	9 (3%) 44 47	54, 79, 108, 132	4 (1%)
1	B	268/289 (92%)	-0.12	19 (7%) 16 16	49, 69, 100, 124	0
All	All	525/578 (90%)	-0.10	28 (5%) 26 28	49, 73, 106, 132	4 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	180	SER	4.4
1	B	14	PRO	3.7
1	B	79	GLY	3.5
1	B	227	ALA	3.4
1	A	179	VAL	3.3
1	B	180	SER	3.0
1	A	214	ILE	2.8
1	A	166	GLY	2.7
1	B	181	ALA	2.7
1	B	83	THR	2.6
1	A	183	CYS	2.5
1	B	281	GLY	2.5
1	B	78	ILE	2.4
1	A	24	ALA	2.4
1	A	79	GLY	2.3
1	B	179	VAL	2.3
1	B	85	TYR	2.3
1	B	213	LEU	2.3
1	B	86	GLN	2.3
1	A	213	LEU	2.2
1	B	212	LEU	2.2
1	B	183	CYS	2.1
1	A	86	GLN	2.1
1	B	29	ARG	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	167	ALA	2.0
1	B	214	ILE	2.0
1	B	15	ASP	2.0
1	B	80	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	A	290	5/5	0.88	0.35	120,120,124,129	0
3	PO4	B	290	5/5	0.92	0.24	100,105,111,112	0
2	SAH	A	3001	26/26	0.94	0.21	67,86,91,97	0
2	SAH	B	3002	26/26	0.95	0.19	54,66,72,75	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.