



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 11, 2018 – 05:19 am GMT

PDB ID : 5KPK
Title : Glycogen Synthase Kinase 3 beta Complexed with BRD0209
Authors : Lakshminarasimhan, D.; White, A.; Nadupalli, A.; Suto, R.K.
Deposited on : 2016-07-04
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3481 (2.40-2.40)
Clashscore	122126	3956 (2.40-2.40)
Ramachandran outliers	120053	3897 (2.40-2.40)
Sidechain outliers	120020	3898 (2.40-2.40)
RSRZ outliers	108989	3386 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	424	<div> <div> <div></div> <div>75%</div> <div>6%</div> <div>18%</div> </div> </div>
1	B	424	<div> <div> <div></div> <div>74%</div> <div>6%</div> <div>19%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycogen synthase kinase-3 beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	347	Total	C	N	O	P	S	0	0	0
			2775	1784	475	504	1	11			
1	B	345	Total	C	N	O	P	S	0	0	0
			2769	1779	474	504	1	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P49841
A	-2	SER	-	expression tag	UNP P49841
A	-1	PRO	-	expression tag	UNP P49841
A	0	GLY	-	expression tag	UNP P49841
B	-3	GLY	-	expression tag	UNP P49841
B	-2	SER	-	expression tag	UNP P49841
B	-1	PRO	-	expression tag	UNP P49841
B	0	GLY	-	expression tag	UNP P49841

- Molecule 2 is (4 {S})-3-cyclopropyl-4,7,7-trimethyl-4-phenyl-2,6,8,9-tetrahydropyrazolo[3,4-b]quinolin-5-one (three-letter code: 6VK) (formula: C₂₂H₂₅N₃O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	22	3	1		
2	B	1	Total	C	N	O	0	0
			26	22	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	120	Total	O	0	0
			120	120		
3	B	120	Total	O	0	0
			120	120		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycogen synthase kinase-3 beta



- Molecule 1: Glycogen synthase kinase-3 beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	64.37Å 67.71Å 67.56Å 79.56° 88.18° 76.50°	Depositor
Resolution (Å)	50.00 – 2.40 36.61 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.5 (50.00-2.40) 97.6 (36.61-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.167 , 0.224 0.174 , 0.223	Depositor DCC
R_{free} test set	2099 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	46.8	Xtriage
Anisotropy	0.245	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5836	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 6VK, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/2827	0.94	3/3847 (0.1%)
1	B	0.74	0/2821	0.96	11/3837 (0.3%)
All	All	0.74	0/5648	0.95	14/7684 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	180	ARG	NE-CZ-NH1	10.14	125.37	120.30
1	B	144	ARG	NE-CZ-NH1	9.80	125.20	120.30
1	B	180	ARG	NE-CZ-NH1	8.33	124.47	120.30
1	A	180	ARG	NE-CZ-NH2	-7.50	116.55	120.30
1	B	144	ARG	NE-CZ-NH2	-6.34	117.13	120.30
1	B	223	ARG	NE-CZ-NH2	6.32	123.46	120.30
1	B	180	ARG	NE-CZ-NH2	-6.14	117.23	120.30
1	B	328	ARG	NE-CZ-NH1	5.85	123.22	120.30
1	B	67	PHE	CA-C-N	5.81	127.82	116.20
1	B	113	ARG	NE-CZ-NH2	-5.74	117.43	120.30
1	B	113	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	B	220	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	A	239	ASP	CB-CG-OD1	5.38	123.14	118.30
1	B	328	ARG	NE-CZ-NH2	-5.24	117.68	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	67	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2775	0	2784	3	0
1	B	2769	0	2784	6	0
2	A	26	0	0	0	0
2	B	26	0	0	0	0
3	A	120	0	0	0	0
3	B	120	0	0	0	0
All	All	5836	0	5568	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:ARG:HD3	1:A:203:SER:O	2.06	0.56
1:B:180:ARG:HD3	1:B:203:SER:O	2.08	0.54
1:B:291:PHE:CZ	1:B:293:PHE:HB2	2.42	0.53
1:B:301:TRP:CD1	1:B:318:SER:HB3	2.48	0.48
1:B:80:GLU:CD	1:B:113:ARG:HH22	2.18	0.47
1:B:313:ALA:HB2	1:B:339:PHE:CE1	2.52	0.44
1:A:289:THR:OG1	1:A:290:GLU:N	2.51	0.44
1:B:48:PRO:O	1:B:50:ARG:N	2.53	0.41
1:A:188:LEU:O	1:A:196:LEU:HA	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/424 (81%)	322 (94%)	20 (6%)	0	100	100
1	B	340/424 (80%)	324 (95%)	14 (4%)	2 (1%)	27	39
All	All	682/848 (80%)	646 (95%)	34 (5%)	2 (0%)	43	58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	49	ASP
1	B	271	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/365 (84%)	286 (94%)	20 (6%)	19	29
1	B	307/365 (84%)	291 (95%)	16 (5%)	25	41
All	All	613/730 (84%)	577 (94%)	36 (6%)	21	34

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	THR
1	A	91	LYS
1	A	92	ARG
1	A	93	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	144	ARG
1	A	147	SER
1	A	169	LEU
1	A	198	LEU
1	A	206	GLN
1	A	209	ARG
1	A	232	THR
1	A	247	LEU
1	A	252	LEU
1	A	266	LEU
1	A	279	GLU
1	A	291	PHE
1	A	317	CYS
1	A	329	LEU
1	A	350	LEU
1	A	383	ARG
1	B	49	ASP
1	B	52	GLN
1	B	59	THR
1	B	87	VAL
1	B	91	LYS
1	B	113	ARG
1	B	125	GLU
1	B	144	ARG
1	B	169	LEU
1	B	198	LEU
1	B	206	GLN
1	B	261	SER
1	B	263	VAL
1	B	266	LEU
1	B	306	ARG
1	B	350	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PTR	A	216	1	16,16,17	0.85	0	21,22,24	1.36	4 (19%)
1	PTR	B	216	1	16,16,17	0.97	0	21,22,24	1.68	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	216	1	-	0/9/11/13	0/1/1/1
1	PTR	B	216	1	-	0/9/11/13	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	216	PTR	O2P-P-OH	-2.24	97.92	105.50
1	A	216	PTR	CG-CB-CA	-2.17	109.91	114.23
1	B	216	PTR	O-C-CA	-2.02	119.40	124.96
1	A	216	PTR	O3P-P-OH	-2.02	98.68	105.50
1	A	216	PTR	CD2-CE2-CZ	2.03	122.23	119.73
1	B	216	PTR	O3P-P-O2P	2.44	117.23	107.59
1	B	216	PTR	O3P-P-OH	2.68	114.54	105.50
1	B	216	PTR	O2P-P-O1P	2.86	121.75	110.60
1	A	216	PTR	O3P-P-O2P	2.92	119.14	107.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	6VK	A	500	-	26,30,30	2.30	5 (19%)	33,48,48	1.94	6 (18%)
2	6VK	B	500	-	26,30,30	2.62	6 (23%)	33,48,48	1.69	9 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6VK	A	500	-	-	0/6/46/46	0/4/5/5
2	6VK	B	500	-	-	0/6/46/46	0/4/5/5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	6VK	C14-C13	-5.97	1.47	1.54
2	A	500	6VK	C14-C13	-5.30	1.48	1.54
2	B	500	6VK	N2-N1	2.37	1.41	1.37
2	A	500	6VK	C6-N	2.62	1.38	1.35
2	A	500	6VK	N2-N1	2.76	1.42	1.37
2	B	500	6VK	C6-N	2.89	1.38	1.35
2	B	500	6VK	C7-C6	2.96	1.53	1.50
2	B	500	6VK	C8-N	3.18	1.42	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	6VK	C8-N	3.43	1.42	1.38
2	A	500	6VK	C9-N2	8.10	1.44	1.33
2	B	500	6VK	C9-N2	9.22	1.45	1.33

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	6VK	C7-C6-N	-5.35	111.14	115.79
2	A	500	6VK	C7-C6-N	-5.25	111.23	115.79
2	A	500	6VK	C21-C16-C14	-2.89	116.72	121.17
2	B	500	6VK	C12-C10-C9	-2.71	115.73	119.33
2	A	500	6VK	C15-C14-C13	-2.53	103.05	114.39
2	A	500	6VK	C12-C10-C9	-2.47	116.05	119.33
2	B	500	6VK	C4-C5-C6	-2.12	116.97	119.00
2	B	500	6VK	C21-C16-C14	-2.07	117.99	121.17
2	B	500	6VK	C2-C1-C7	-2.06	106.56	109.84
2	B	500	6VK	C15-C14-C13	-2.03	105.28	114.39
2	B	500	6VK	C13-C14-C5	2.27	107.69	103.96
2	B	500	6VK	C11-C10-C9	2.32	122.40	119.33
2	B	500	6VK	C5-C6-N	2.68	124.73	120.82
2	A	500	6VK	C11-C10-C9	4.25	124.96	119.33
2	A	500	6VK	C5-C6-N	4.59	127.52	120.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/424 (81%)	-0.23	6 (1%) 70 67	31, 51, 97, 133	0
1	B	344/424 (81%)	-0.25	8 (2%) 60 57	30, 50, 93, 128	0
All	All	690/848 (81%)	-0.24	14 (2%) 65 62	30, 51, 95, 133	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	384	ILE	7.4
1	B	93	PHE	6.7
1	B	293	PHE	5.7
1	A	93	PHE	5.6
1	A	35	SER	5.2
1	B	289	THR	5.1
1	B	92	ARG	4.1
1	A	92	ARG	3.7
1	A	293	PHE	3.6
1	B	35	SER	3.0
1	B	292	LYS	2.8
1	A	36	LYS	2.6
1	B	290	GLU	2.5
1	B	291	PHE	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	PTR	A	216	16/17	0.98	0.11	38,44,49,50	0
1	PTR	B	216	16/17	0.98	0.11	32,39,46,47	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	6VK	A	500	26/26	0.98	0.11	34,38,42,44	0
2	6VK	B	500	26/26	0.99	0.12	31,37,39,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.