



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 18, 2019 – 11:15 PM EST

PDB ID : 3L2L
Title : X-ray Crystallographic Analysis of Pig Pancreatic Alpha-Amylase with Limit Dextrin and Oligosaccharide
Authors : Larson, S.B.; Day, J.S.; McPherson, A.
Deposited on : 2009-12-15
Resolution : 2.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.4
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5449 (2.14-2.10)
Clashscore	122126	5972 (2.14-2.10)
Ramachandran outliers	120053	5914 (2.14-2.10)
Sidechain outliers	120020	5915 (2.14-2.10)
RSRZ outliers	108989	5321 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	705[A]	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	801	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9218 atoms, of which 4197 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pancreatic alpha-amylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	496	8372	2710	4084	747	805	26	0	56	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	411	ASP	ALA	SEE REMARK 999	UNP P00690

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	21	6	10	5	0	1
2	A	1	21	6	10	5	0	1
2	A	1	21	6	10	5	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	1
			23	6	11	6		
2	A	1	Total	C	H	O	0	1
			21	6	10	5		
2	A	1	Total	C	H	O	0	1
			21	6	10	5		
2	A	1	Total	C	H	O	0	1
			20	6	9	5		
2	A	1	Total	C	H	O	0	1
			21	6	10	5		
2	A	1	Total	C	H	O	0	1
			23	6	11	6		
2	A	1	Total	C	H	O	0	1
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			24	6	12	6		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	584	Total	O	0	76
			607	607		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pancreatic alpha-amylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.59Å 114.82Å 118.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.01 – 2.11 14.01 – 2.11	Depositor EDS
% Data completeness (in resolution range)	84.0 (14.01-2.11) 84.0 (14.01-2.11)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 2.12Å)	Xtriage
Refinement program	REFMAC 5.5.0089	Depositor
R, R_{free}	0.115 , 0.157 0.123 , 0.163	Depositor DCC
R_{free} test set	4749 reflections (10.07%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.020 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9218	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC, PCA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/4483	0.71	3/6085 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	195	ARG	NE-CZ-NH1	6.46	123.53	120.30
1	A	195	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	A	402	ASP	CB-CG-OD2	-5.05	113.76	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4288	4084	4075	32	0
2	A	124	113	108	5	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	607	0	0	14	0
All	All	5021	4197	4183	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178[A]:MET:HE1	5:A:928:HOH:O	1.75	0.85
2:A:601[A]:GLC:H3	5:A:1034:HOH:O	1.91	0.69
1:A:441[B]:GLN:OE1	1:A:441[B]:GLN:HA	1.92	0.69
1:A:153[A]:ASP:OD2	1:A:156[A]:GLN:HG3	1.95	0.66
1:A:460[B]:ASN:CG	5:A:525[B]:HOH:O	2.36	0.64
1:A:261[B]:LYS:HE3	5:A:574[B]:HOH:O	1.99	0.62
1:A:436[B]:LEU:HD23	5:A:748:HOH:O	2.02	0.59
1:A:470[A]:SER:OG	1:A:472:ASP:OD1	2.20	0.59
1:A:261[B]:LYS:CE	5:A:574[B]:HOH:O	2.53	0.57
1:A:123:ASN:HB3	5:A:533:HOH:O	2.04	0.56
1:A:220[A]:ASN:HB3	5:A:638:HOH:O	2.06	0.56
1:A:153[B]:ASP:HB3	1:A:156[B]:GLN:HE21	1.74	0.52
1:A:495[A]:LYS:NZ	5:A:990:HOH:O	2.41	0.52
1:A:424[B]:ARG:NE	5:A:830:HOH:O	2.44	0.51
1:A:340[B]:SER:OG	1:A:386:HIS:ND1	2.39	0.50
1:A:418:ALA:HB2	1:A:428[A]:VAL:HG13	1.94	0.49
1:A:123:ASN:CB	5:A:533:HOH:O	2.60	0.48
1:A:436[B]:LEU:HD22	1:A:437:SER:N	2.29	0.47
1:A:396[B]:TRP:HZ3	5:A:822:HOH:O	1.98	0.47
1:A:140:LYS:NZ	2:A:801:GLC:O3	2.49	0.46
1:A:9:GLY:HA2	5:A:956:HOH:O	2.16	0.45
1:A:53:ASN:HB2	2:A:602[A]:GLC:H3	1.99	0.44
1:A:449:TYR:CZ	1:A:495[A]:LYS:HD2	2.52	0.43
1:A:305[B]:HIS:NE2	1:A:352:GLN:OE1	2.51	0.43
1:A:53:ASN:OD1	2:A:602[A]:GLC:O4	2.37	0.42
1:A:203:TRP:CE2	2:A:801:GLC:H2	2.55	0.42
1:A:369[B]:GLU:HG2	1:A:370:VAL:N	2.35	0.42
1:A:442:THR:O	1:A:444[A]:LEU:HD13	2.18	0.42
1:A:436[B]:LEU:CD2	5:A:748:HOH:O	2.63	0.41
1:A:132[A]:SER:OG	1:A:134[A]:TRP:CD1	2.74	0.41
1:A:129:VAL:CG1	1:A:178[B]:MET:HG2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	551/496 (111%)	538 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	468/411 (114%)	460 (98%)	8 (2%)	63	69

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115[A]	CYS
1	A	115[B]	CYS
1	A	134[A]	TRP
1	A	134[B]	TRP
1	A	170	LEU
1	A	274[A]	MET
1	A	274[B]	MET
1	A	293	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	ASN
1	A	243	GLN
1	A	408	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PCA	A	1	1	8,8,9	1.62	2 (25%)	9,10,12	2.30	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	CD-N	3.60	1.44	1.34
1	A	1	PCA	CA-C	2.01	1.52	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CA-N-CD	-3.32	102.20	113.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CB-CA-N	3.25	112.63	103.30
1	A	1	PCA	CG-CD-N	3.03	116.64	108.35
1	A	1	PCA	O-C-CA	-2.65	119.06	125.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	A	601[A]	2	11,11,12	0.57	0	15,15,17	2.78	8 (53%)
2	GLC	A	602[A]	2	11,11,12	1.12	1 (9%)	15,15,17	2.27	4 (26%)
2	GLC	A	603[A]	2	11,11,12	0.73	0	15,15,17	1.58	3 (20%)
2	GLC	A	604[A]	2	12,12,12	0.51	0	17,17,17	1.31	1 (5%)
2	GLC	A	701[A]	2	11,11,12	0.60	0	15,15,17	1.68	2 (13%)
2	GLC	A	702[A]	2	11,11,12	0.70	0	15,15,17	1.82	2 (13%)
2	GLC	A	703[A]	2	11,11,12	0.75	0	15,15,17	3.01	4 (26%)
2	GLC	A	704[A]	2	11,11,12	0.66	0	15,15,17	1.43	1 (6%)
2	GLC	A	705[A]	2	12,12,12	0.65	0	17,17,17	1.90	3 (17%)
2	GLC	A	706[A]	2	11,11,12	0.71	0	15,15,17	2.23	4 (26%)
2	GLC	A	801	-	12,12,12	0.48	0	17,17,17	0.86	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	601[A]	2	-	2/2/19/22	0/1/1/1
2	GLC	A	602[A]	2	-	2/2/19/22	0/1/1/1
2	GLC	A	603[A]	2	-	0/2/19/22	0/1/1/1
2	GLC	A	604[A]	2	-	2/2/22/22	0/1/1/1
2	GLC	A	701[A]	2	-	2/2/19/22	0/1/1/1
2	GLC	A	702[A]	2	-	0/2/19/22	0/1/1/1
2	GLC	A	703[A]	2	-	2/2/19/22	0/1/1/1
2	GLC	A	704[A]	2	-	0/2/19/22	0/1/1/1
2	GLC	A	705[A]	2	1/1/5/5	2/2/22/22	0/1/1/1
2	GLC	A	706[A]	2	-	2/2/19/22	0/1/1/1
2	GLC	A	801	-	1/1/5/5	1/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	602[A]	GLC	C2-C3	3.01	1.57	1.52

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	703[A]	GLC	C1-O5-C5	9.84	125.58	112.20
2	A	601[A]	GLC	C1-O5-C5	7.35	122.19	112.20
2	A	602[A]	GLC	C1-O5-C5	6.34	120.82	112.20
2	A	706[A]	GLC	C1-C2-C3	5.45	116.36	109.66
2	A	701[A]	GLC	C1-O5-C5	5.25	119.34	112.20
2	A	705[A]	GLC	O5-C1-C2	5.05	119.37	110.31
2	A	702[A]	GLC	C1-C2-C3	-4.83	103.72	109.66
2	A	705[A]	GLC	O1-C1-C2	3.88	120.03	109.03
2	A	703[A]	GLC	C1-C2-C3	-3.70	105.10	109.66
2	A	604[A]	GLC	C1-O5-C5	3.70	120.76	113.68
2	A	601[A]	GLC	O5-C1-C2	-3.69	105.11	110.79
2	A	706[A]	GLC	C3-C4-C5	-3.67	103.63	110.23
2	A	603[A]	GLC	C1-C2-C3	3.60	114.08	109.66
2	A	601[A]	GLC	O5-C5-C4	3.45	119.22	110.83
2	A	704[A]	GLC	O5-C5-C6	3.37	112.48	107.15
2	A	602[A]	GLC	O2-C2-C3	3.35	116.70	110.16
2	A	602[A]	GLC	O3-C3-C2	3.34	116.22	110.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601[A]	GLC	C6-C5-C4	-3.26	105.33	113.00
2	A	706[A]	GLC	O5-C5-C6	3.22	112.25	107.15
2	A	706[A]	GLC	O5-C1-C2	3.20	115.72	110.79
2	A	705[A]	GLC	O1-C1-O5	3.07	119.64	110.41
2	A	702[A]	GLC	C2-C3-C4	-3.00	105.70	110.89
2	A	603[A]	GLC	O5-C5-C6	2.89	111.73	107.15
2	A	603[A]	GLC	C1-O5-C5	2.60	115.74	112.20
2	A	703[A]	GLC	O5-C5-C6	2.60	111.26	107.15
2	A	601[A]	GLC	C1-C2-C3	2.55	112.79	109.66
2	A	703[A]	GLC	C2-C3-C4	-2.43	106.68	110.89
2	A	601[A]	GLC	C2-C3-C4	-2.28	106.95	110.89
2	A	601[A]	GLC	O2-C2-C1	2.18	113.60	109.17
2	A	701[A]	GLC	C2-C3-C4	-2.14	107.19	110.89
2	A	601[A]	GLC	O5-C5-C6	2.12	110.51	107.15
2	A	602[A]	GLC	O5-C5-C4	2.05	115.80	110.83
2	A	801	GLC	C1-O5-C5	-2.04	109.78	113.68

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	705[A]	GLC	C1
2	A	801	GLC	C1

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	706[A]	GLC	O5-C5-C6-O6
2	A	703[A]	GLC	O5-C5-C6-O6
2	A	601[A]	GLC	O5-C5-C6-O6
2	A	602[A]	GLC	O5-C5-C6-O6
2	A	706[A]	GLC	C4-C5-C6-O6
2	A	604[A]	GLC	C4-C5-C6-O6
2	A	602[A]	GLC	C4-C5-C6-O6
2	A	705[A]	GLC	O5-C5-C6-O6
2	A	604[A]	GLC	O5-C5-C6-O6
2	A	703[A]	GLC	C4-C5-C6-O6
2	A	705[A]	GLC	C4-C5-C6-O6
2	A	701[A]	GLC	O5-C5-C6-O6
2	A	601[A]	GLC	C4-C5-C6-O6
2	A	801	GLC	O5-C5-C6-O6
2	A	701[A]	GLC	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601[A]	GLC	1	0
2	A	602[A]	GLC	2	0
2	A	801	GLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	495/496 (99%)	-1.02	2 (0%) 92 93	10, 16, 29, 61	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	134[A]	TRP	3.3
1	A	350	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	1	8/9	0.97	0.07	20,27,31,43	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	A	601[A]	11/12	0.50	0.39	35,39,43,44	21
2	GLC	A	701[A]	11/12	0.65	0.31	20,24,28,29	21
2	GLC	A	801	12/12	0.74	0.32	22,31,33,34	24
2	GLC	A	602[A]	11/12	0.78	0.17	12,31,33,37	21
2	GLC	A	706[A]	11/12	0.80	0.19	37,42,45,46	21
2	GLC	A	604[A]	12/12	0.83	0.23	21,27,30,32	23
2	GLC	A	703[A]	11/12	0.87	0.12	33,40,44,48	20
2	GLC	A	702[A]	11/12	0.88	0.24	36,44,50,53	21
2	GLC	A	705[A]	12/12	0.88	0.11	20,36,45,48	0
2	GLC	A	603[A]	11/12	0.92	0.10	22,29,31,32	21
2	GLC	A	704[A]	11/12	0.97	0.08	18,34,39,41	0
4	CL	A	502	1/1	1.00	0.06	15,15,15,15	0
3	CA	A	501	1/1	1.00	0.05	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.