



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 18, 2019 – 11:16 PM EST

PDB ID : 3L2M
Title : X-ray Crystallographic Analysis of Pig Pancreatic Alpha-Amylase with Alpha-cyclodextrin
Authors : Larson, S.B.; Day, J.S.; McPherson, A.
Deposited on : 2009-12-15
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.4
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	10189 (2.00-1.96)
Clashscore	122126	11405 (2.00-1.96)
Ramachandran outliers	120053	11281 (2.00-1.96)
Sidechain outliers	120020	11280 (2.00-1.96)
RSRZ outliers	108989	9953 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	801	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	802	-	-	-	X
2	GLC	A	803	-	-	-	X
2	GLC	A	804	-	-	-	X
2	GLC	A	806	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9278 atoms, of which 4234 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pancreatic alpha-amylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	496	8283	2676	4054	737	790	26	0	47	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	411	ASP	ALA	SEE REMARK 999	UNP P00690

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	21	6	10	5	0	0
2	A	1	21	6	10	5	0	0
2	A	1	21	6	10	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	592	Total 615	O 615	0	42

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pancreatic alpha-amylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.65Å 114.88Å 118.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.30 – 1.97 29.29 – 1.97	Depositor EDS
% Data completeness (in resolution range)	89.2 (29.30-1.97) 89.8 (29.29-1.97)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.96Å)	Xtriage
Refinement program	REFMAC 5.5.0089	Depositor
R, R_{free}	0.127 , 0.160 0.140 , 0.173	Depositor DCC
R_{free} test set	6279 reflections (10.11%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.021 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9278	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC, PCA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/4418	0.65	0/5992

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4229	4054	4051	20	0
2	A	198	180	162	6	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	615	0	0	7	0
All	All	5044	4234	4213	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:ASN:ND2	5:A:647:HOH:O	2.25	0.69
2:A:703:GLC:O2	2:A:704:GLC:O3	2.08	0.62
1:A:470[A]:SER:OG	1:A:472:ASP:OD1	2.17	0.62
1:A:178[B]:MET:HE2	1:A:178[B]:MET:HA	1.86	0.58
2:A:803:GLC:H2	2:A:803:GLC:C6	2.40	0.51
1:A:135:ASP:OD2	2:A:805:GLC:O6	2.29	0.49
1:A:129:VAL:CG1	1:A:178[B]:MET:HG2	2.44	0.47
1:A:135:ASP:OD2	2:A:805:GLC:C6	2.61	0.47
1:A:274[B]:MET:HG3	1:A:277:LEU:HD12	1.97	0.47
1:A:178[B]:MET:HA	1:A:178[B]:MET:CE	2.46	0.45
1:A:22[B]:VAL:HG23	5:A:939:HOH:O	2.18	0.44
1:A:244[A]:SER:HB2	1:A:253:VAL:HG11	1.99	0.44
2:A:803:GLC:C6	2:A:803:GLC:C2	2.94	0.44
1:A:343[A]:ARG:HD3	5:A:833[A]:HOH:O	2.18	0.43
1:A:15:HIS:HD2	5:A:765:HOH:O	2.03	0.42
1:A:436[A]:LEU:HD23	1:A:437:SER:N	2.35	0.42
1:A:162:LEU:HD11	2:A:703:GLC:H62	2.02	0.41
1:A:200:LYS:HE3	5:A:847:HOH:O	2.21	0.41
1:A:4:PRO:HA	1:A:229:PHE:CG	2.56	0.40
1:A:424[A]:ARG:NH1	5:A:1060:HOH:O	2.53	0.40
1:A:174:TYR:O	1:A:178[A]:MET:HG2	2.22	0.40
1:A:495[C]:LYS:NZ	5:A:1097:HOH:O	2.49	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	545/496 (110%)	535 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/411 (112%)	454 (98%)	8 (2%)	63	65

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115[A]	CYS
1	A	115[B]	CYS
1	A	134	TRP
1	A	170	LEU
1	A	274[A]	MET
1	A	274[B]	MET
1	A	343[A]	ARG
1	A	343[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	435	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	1	1	8,8,9	1.53	2 (25%)	9,10,12	2.24	6 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	CD-N	3.38	1.43	1.34
1	A	1	PCA	CA-C	2.20	1.53	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CA-N-CD	-2.93	103.54	113.58
1	A	1	PCA	CB-CA-C	-2.84	108.80	112.70
1	A	1	PCA	CG-CD-N	2.66	115.62	108.35
1	A	1	PCA	CB-CA-N	2.54	110.59	103.30
1	A	1	PCA	O-C-CA	-2.43	119.54	125.11
1	A	1	PCA	OE-CD-CG	-2.43	122.48	126.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 20 ligands modelled in this entry, 2 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	A	601	2	11,11,12	0.72	0	15,15,17	1.05	1 (6%)
2	GLC	A	602	2	11,11,12	0.65	0	15,15,17	1.69	3 (20%)
2	GLC	A	603	2	11,11,12	0.62	0	15,15,17	1.12	1 (6%)
2	GLC	A	604	2	11,11,12	0.57	0	15,15,17	1.38	1 (6%)
2	GLC	A	605	2	11,11,12	0.79	0	15,15,17	1.23	1 (6%)
2	GLC	A	606	2	11,11,12	0.81	0	15,15,17	1.26	2 (13%)
2	GLC	A	701	2	11,11,12	0.55	0	15,15,17	1.23	2 (13%)
2	GLC	A	702	2	11,11,12	0.60	0	15,15,17	1.45	2 (13%)
2	GLC	A	703	2	11,11,12	0.59	0	15,15,17	1.21	1 (6%)
2	GLC	A	704	2	11,11,12	0.65	0	15,15,17	1.06	1 (6%)
2	GLC	A	705	2	11,11,12	0.55	0	15,15,17	1.53	2 (13%)
2	GLC	A	706	2	11,11,12	0.60	0	15,15,17	1.45	2 (13%)
2	GLC	A	801	2	11,11,12	0.62	0	15,15,17	1.28	3 (20%)
2	GLC	A	802	2	11,11,12	0.58	0	15,15,17	1.27	1 (6%)
2	GLC	A	803	2	11,11,12	0.55	0	15,15,17	1.04	2 (13%)
2	GLC	A	804	2	11,11,12	0.58	0	15,15,17	1.22	2 (13%)
2	GLC	A	805	2	11,11,12	0.71	0	15,15,17	1.47	2 (13%)
2	GLC	A	806	2	11,11,12	0.62	0	15,15,17	1.48	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	601	2	-	0/2/19/22	0/1/1/1
2	GLC	A	602	2	-	1/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	603	2	-	0/2/19/22	0/1/1/1
2	GLC	A	604	2	-	0/2/19/22	0/1/1/1
2	GLC	A	605	2	-	0/2/19/22	0/1/1/1
2	GLC	A	606	2	-	0/2/19/22	0/1/1/1
2	GLC	A	701	2	-	0/2/19/22	0/1/1/1
2	GLC	A	702	2	-	2/2/19/22	0/1/1/1
2	GLC	A	703	2	-	2/2/19/22	0/1/1/1
2	GLC	A	704	2	-	1/2/19/22	0/1/1/1
2	GLC	A	705	2	-	2/2/19/22	0/1/1/1
2	GLC	A	706	2	-	1/2/19/22	0/1/1/1
2	GLC	A	801	2	-	2/2/19/22	0/1/1/1
2	GLC	A	802	2	-	2/2/19/22	0/1/1/1
2	GLC	A	803	2	-	2/2/19/22	0/1/1/1
2	GLC	A	804	2	-	2/2/19/22	0/1/1/1
2	GLC	A	805	2	-	2/2/19/22	0/1/1/1
2	GLC	A	806	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	705	GLC	C1-O5-C5	3.83	117.41	112.20
2	A	604	GLC	O4-C4-C3	-3.55	102.11	110.34
2	A	706	GLC	O5-C1-C2	-3.47	105.45	110.79
2	A	703	GLC	O5-C1-C2	-3.32	105.67	110.79
2	A	803	GLC	C1-O5-C5	3.27	116.65	112.20
2	A	702	GLC	O5-C1-C2	-3.26	105.77	110.79
2	A	805	GLC	O5-C1-C2	-3.16	105.91	110.79
2	A	704	GLC	O5-C1-C2	-3.06	106.08	110.79
2	A	603	GLC	O5-C1-C2	-3.05	106.08	110.79
2	A	706	GLC	C1-O5-C5	3.05	116.34	112.20
2	A	804	GLC	O5-C1-C2	-3.04	106.11	110.79
2	A	806	GLC	C1-O5-C5	3.02	116.31	112.20
2	A	701	GLC	C1-O5-C5	3.01	116.29	112.20
2	A	702	GLC	C3-C4-C5	2.96	115.54	110.23
2	A	602	GLC	C1-O5-C5	2.95	116.22	112.20
2	A	806	GLC	O5-C1-C2	-2.82	106.45	110.79
2	A	804	GLC	O5-C5-C6	2.75	111.50	107.15
2	A	705	GLC	O5-C1-C2	-2.62	106.75	110.79
2	A	802	GLC	O5-C1-C2	-2.55	106.86	110.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	606	GLC	O2-C2-C3	2.53	115.11	110.16
2	A	601	GLC	O4-C4-C3	-2.52	104.50	110.34
2	A	602	GLC	C3-C4-C5	2.51	114.73	110.23
2	A	605	GLC	O4-C4-C3	-2.44	104.68	110.34
2	A	801	GLC	O5-C1-C2	-2.34	107.19	110.79
2	A	801	GLC	C2-C3-C4	2.24	114.78	110.89
2	A	701	GLC	C1-C2-C3	2.23	112.40	109.66
2	A	801	GLC	C1-C2-C3	2.12	112.26	109.66
2	A	602	GLC	O5-C1-C2	-2.11	107.54	110.79
2	A	606	GLC	O3-C3-C2	2.11	113.94	110.02
2	A	803	GLC	O5-C5-C6	2.08	110.44	107.15
2	A	805	GLC	O5-C5-C4	2.08	115.88	110.83

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	803	GLC	O5-C5-C6-O6
2	A	805	GLC	O5-C5-C6-O6
2	A	803	GLC	C4-C5-C6-O6
2	A	802	GLC	O5-C5-C6-O6
2	A	802	GLC	C4-C5-C6-O6
2	A	805	GLC	C4-C5-C6-O6
2	A	801	GLC	C4-C5-C6-O6
2	A	705	GLC	C4-C5-C6-O6
2	A	702	GLC	C4-C5-C6-O6
2	A	703	GLC	O5-C5-C6-O6
2	A	703	GLC	C4-C5-C6-O6
2	A	706	GLC	O5-C5-C6-O6
2	A	801	GLC	O5-C5-C6-O6
2	A	804	GLC	O5-C5-C6-O6
2	A	806	GLC	O5-C5-C6-O6
2	A	705	GLC	O5-C5-C6-O6
2	A	702	GLC	O5-C5-C6-O6
2	A	806	GLC	C4-C5-C6-O6
2	A	704	GLC	C4-C5-C6-O6
2	A	804	GLC	C4-C5-C6-O6
2	A	602	GLC	C4-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	703	GLC	2	0
2	A	704	GLC	1	0
2	A	803	GLC	2	0
2	A	805	GLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	495/496 (99%)	-0.16	14 (2%) 53 55	7, 13, 24, 48	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	134	TRP	4.7
1	A	350	ASN	3.7
1	A	460[A]	ASN	3.5
1	A	349	VAL	3.4
1	A	461[A]	SER	3.1
1	A	374	ALA	3.1
1	A	327	PHE	3.0
1	A	348	PHE	2.9
1	A	9	GLY	2.6
1	A	142[A]	LYS	2.5
1	A	352	GLN	2.5
1	A	351	GLY	2.3
1	A	99	ILE	2.2
1	A	338	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	1	8/9	0.97	0.15	16,22,27,39	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	A	704	11/12	0.44	0.29	91,101,104,104	0
2	GLC	A	803	11/12	0.50	0.62	117,120,121,121	0
2	GLC	A	701	11/12	0.53	0.29	79,86,92,98	0
2	GLC	A	801	11/12	0.53	0.74	115,118,118,119	0
2	GLC	A	804	11/12	0.54	0.44	110,112,115,118	0
2	GLC	A	702	11/12	0.62	0.25	89,92,98,99	0
2	GLC	A	802	11/12	0.64	0.55	108,113,115,117	0
2	GLC	A	805	11/12	0.66	0.33	97,106,109,109	0
2	GLC	A	705	11/12	0.67	0.37	102,106,107,107	0
2	GLC	A	703	11/12	0.68	0.21	101,102,102,103	0
2	GLC	A	706	11/12	0.73	0.36	92,101,103,105	0
2	GLC	A	806	11/12	0.78	0.56	111,113,116,116	0
2	GLC	A	603	11/12	0.94	0.20	29,34,41,46	0
2	GLC	A	605	11/12	0.95	0.14	23,28,34,40	0
2	GLC	A	604	11/12	0.95	0.18	25,31,42,47	0
2	GLC	A	602	11/12	0.96	0.10	20,27,29,29	0
2	GLC	A	606	11/12	0.96	0.11	18,22,26,28	0
2	GLC	A	601	11/12	0.97	0.10	16,20,22,24	0
4	CL	A	502	1/1	1.00	0.08	15,15,15,15	0
3	CA	A	501	1/1	1.00	0.07	15,15,15,15	0

6.5 Other polymers ⓘ

There are no such residues in this entry.