



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 06:12 pm GMT

PDB ID : 4L2I  
Title : Electron transferring flavoprotein of *Acidaminococcus fermentans*: Towards a mechanism of flavin-based electron bifurcation  
Authors : Mowafy, A.M.; Chowdhury, N.P.; Demmer, J.; Upadhyay, V.; Kolzer, S.; Jayamani, E.; Kahnt, J.; Demmer, U.; Ermler, U.; Buckel, W.  
Deposited on : 2013-06-04  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1761 (1.48-1.44)
Clashscore	122126	1816 (1.48-1.44)
Ramachandran outliers	120053	1793 (1.48-1.44)
Sidechain outliers	120020	1793 (1.48-1.44)
RSRZ outliers	108989	1733 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div></div> </div> </div>
2	B	263	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>7%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Electron transfer flavoprotein alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	2	0
			2597	1628	445	505	19			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	341	HIS	-	EXPRESSION TAG	UNP D2RIQ3
A	342	HIS	-	EXPRESSION TAG	UNP D2RIQ3
A	343	HIS	-	EXPRESSION TAG	UNP D2RIQ3
A	344	HIS	-	EXPRESSION TAG	UNP D2RIQ3
A	345	HIS	-	EXPRESSION TAG	UNP D2RIQ3
A	346	HIS	-	EXPRESSION TAG	UNP D2RIQ3

- Molecule 2 is a protein called Electron transfer flavoprotein alpha/beta-subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	263	Total	C	N	O	S	0	3	0
			2006	1258	341	390	17			

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
3	B	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0

- Molecule 5 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	271	Total	O	0	0
			271	271		
6	B	244	Total	O	0	0
			244	244		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Electron transfer flavoprotein alpha subunit



- Molecule 2: Electron transfer flavoprotein alpha/beta-subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.52Å 84.79Å 106.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.07 – 1.45 45.07 – 1.45	Depositor EDS
% Data completeness (in resolution range)	95.1 (45.07-1.45) 95.1 (45.07-1.45)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.150 , 0.208 0.158 , 0.215	Depositor DCC
$R_{free}$ test set	6099 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.5	Xtriage
Anisotropy	1.467	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.12	1/2644 (0.0%)	1.20	9/3573 (0.3%)
2	B	1.13	2/2033 (0.1%)	1.26	13/2758 (0.5%)
All	All	1.12	3/4677 (0.1%)	1.23	22/6331 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	37	TYR	CE1-CZ	5.97	1.46	1.38
1	A	260	PHE	CG-CD1	5.36	1.46	1.38
2	B	99	ALA	C-O	5.13	1.33	1.23

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	179	ARG	NE-CZ-NH1	-14.01	113.29	120.30
1	A	118	ARG	NE-CZ-NH2	-13.75	113.42	120.30
1	A	118	ARG	NE-CZ-NH1	12.58	126.59	120.30
2	B	179	ARG	NE-CZ-NH2	10.57	125.58	120.30
2	B	72	ASP	CB-CG-OD1	7.62	125.15	118.30
1	A	122	ARG	NE-CZ-NH1	7.51	124.06	120.30
2	B	122	ASP	CB-CG-OD2	-7.29	111.74	118.30
2	B	158	ARG	NE-CZ-NH2	-7.15	116.72	120.30
2	B	202	TYR	CB-CG-CD2	-6.54	117.08	121.00
2	B	212	ASP	CB-CG-OD1	6.17	123.85	118.30
2	B	86	ASP	CB-CG-OD1	6.16	123.85	118.30
2	B	40	TYR	CB-CG-CD2	-5.96	117.42	121.00
2	B	202	TYR	CB-CG-CD1	5.80	124.48	121.00
2	B	158	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	38	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	135	ASP	CB-CG-OD2	-5.72	113.15	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	163	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	A	38	ARG	NE-CZ-NH2	-5.39	117.60	120.30
2	B	71	ARG	NE-CZ-NH2	-5.39	117.61	120.30
1	A	217	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	A	114	ASP	CB-CG-OD2	-5.05	113.76	118.30
2	B	158	ARG	N-CA-CB	-5.04	101.54	110.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2597	0	2571	11	0
2	B	2006	0	2055	4	0
3	A	53	0	31	0	0
3	B	53	0	31	2	0
4	A	1	0	0	0	0
5	B	27	0	12	0	0
6	A	271	0	0	0	0
6	B	244	0	0	2	0
All	All	5252	0	4700	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:ARG:HG2	2:B:163[B]:MET:HE2	1.24	1.13
1:A:209:VAL:HG21	1:A:215:ALA:HB2	1.50	0.91
1:A:209:VAL:CG2	1:A:215:ALA:HB2	2.22	0.68
1:A:54:THR:O	1:A:76:GLN:HA	2.09	0.53
2:B:65:HIS:HD2	6:B:585:HOH:O	1.95	0.49
1:A:106:ILE:O	1:A:168:THR:HA	2.13	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:TRP:CD1	1:A:49:ALA:HB3	2.49	0.47
2:B:47:GLN:HE22	2:B:181:LYS:HG2	1.80	0.47
1:A:310:PRO:O	1:A:313[A]:GLU:HB2	2.17	0.45
2:B:86:ASP:HB3	2:B:89:PHE:CD2	2.53	0.44
1:A:27:VAL:HG23	1:A:29:VAL:HG12	2.00	0.43
1:A:109:THR:O	1:A:113:ARG:HG3	2.20	0.42
3:B:301:FAD:HM82	6:B:644:HOH:O	2.18	0.42
1:A:316:LYS:HA	1:A:316:LYS:HD2	1.90	0.42
1:A:148:ALA:HB3	3:B:301:FAD:HM72	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/346 (98%)	329 (97%)	8 (2%)	1 (0%)	43	18
2	B	264/263 (100%)	256 (97%)	8 (3%)	0	100	100
All	All	602/609 (99%)	585 (97%)	16 (3%)	1 (0%)	49	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/283 (98%)	276 (99%)	2 (1%)	85	66
2	B	223/220 (101%)	220 (99%)	3 (1%)	71	40
All	All	501/503 (100%)	496 (99%)	5 (1%)	78	51

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	SER
1	A	137	GLU
2	B	153	THR
2	B	198	GLU
2	B	225	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	65	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FAD	A	401	-	51,58,58	2.19	17 (33%)	57,89,89	2.51	19 (33%)
3	FAD	B	301	-	51,58,58	2.39	9 (17%)	57,89,89	2.47	19 (33%)
5	NAD	B	302	-	25,29,48	1.74	4 (16%)	25,45,73	1.38	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	401	-	-	0/28/50/50	0/6/6/6
3	FAD	B	301	-	-	0/28/50/50	0/6/6/6
5	NAD	B	302	-	-	0/12/32/62	0/3/3/5

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	FAD	C2-N1	-2.99	1.32	1.38
3	A	401	FAD	C2-N3	-2.76	1.32	1.38
3	A	401	FAD	C9-C9A	-2.35	1.35	1.40
3	B	301	FAD	P-O1P	-2.21	1.42	1.50
3	A	401	FAD	C8M-C8	-2.18	1.46	1.51
3	A	401	FAD	C2A-N1A	2.34	1.38	1.33
3	A	401	FAD	C2A-N3A	2.39	1.36	1.32
3	B	301	FAD	O2'-C2'	2.47	1.48	1.43
3	B	301	FAD	O3'-C3'	2.79	1.49	1.43
3	A	401	FAD	C5A-C4A	2.87	1.47	1.40
3	A	401	FAD	C10-N1	2.94	1.37	1.33
3	A	401	FAD	C8-C7	3.01	1.48	1.40
5	B	302	NAD	C2A-N3A	3.15	1.37	1.32
5	B	302	NAD	O4B-C1B	3.16	1.45	1.41
3	A	401	FAD	O4B-C1B	3.50	1.46	1.41
3	A	401	FAD	C4X-C10	3.50	1.46	1.41
3	B	301	FAD	C4X-C10	3.56	1.46	1.41
3	B	301	FAD	C9A-C5X	3.59	1.49	1.42
5	B	302	NAD	C5A-C4A	3.61	1.48	1.40
3	A	401	FAD	C5'-C4'	3.68	1.57	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	FAD	C4X-N5	3.76	1.38	1.33
3	A	401	FAD	C9A-C5X	3.89	1.50	1.42
3	A	401	FAD	C9-C8	3.98	1.48	1.37
5	B	302	NAD	PN-O3	4.92	1.67	1.60
3	A	401	FAD	C4-N3	5.67	1.43	1.33
3	A	401	FAD	C9A-N10	6.24	1.46	1.38
3	B	301	FAD	C8A-N9A	6.51	1.44	1.36
3	B	301	FAD	C4X-N5	6.84	1.43	1.33
3	B	301	FAD	C9A-N10	6.93	1.47	1.38
3	B	301	FAD	C10-N1	7.44	1.43	1.33

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	FAD	C4B-O4B-C1B	-9.99	99.42	109.83
3	B	301	FAD	C10-C4X-N5	-5.11	114.71	120.59
3	A	401	FAD	C4X-C4-N3	-4.43	117.16	123.47
3	A	401	FAD	N3A-C2A-N1A	-4.07	125.38	128.86
5	B	302	NAD	C1B-N9A-C4A	-3.96	119.79	126.64
3	B	301	FAD	N3A-C2A-N1A	-3.83	125.59	128.86
3	A	401	FAD	C1B-N9A-C4A	-3.80	120.08	126.64
3	B	301	FAD	C6-C5X-C9A	-3.57	114.22	119.01
3	B	301	FAD	C7M-C7-C8	-3.42	113.58	120.72
3	A	401	FAD	C10-C4X-N5	-3.41	116.68	120.59
3	B	301	FAD	C4X-C4-N3	-3.36	118.69	123.47
3	B	301	FAD	C5X-C9A-N10	-3.17	115.30	117.71
3	B	301	FAD	C9A-N10-C10	-2.83	118.00	121.77
3	B	301	FAD	C9-C8-C7	-2.74	115.14	119.94
5	B	302	NAD	N3A-C2A-N1A	-2.67	126.58	128.86
3	A	401	FAD	C5X-C9A-N10	-2.51	115.80	117.71
3	B	301	FAD	C1B-N9A-C4A	-2.45	122.41	126.64
3	A	401	FAD	C9-C8-C7	-2.40	115.73	119.94
3	B	301	FAD	O4'-C4'-C5'	-2.22	105.08	110.02
3	A	401	FAD	C8M-C8-C9	-2.05	115.30	120.36
3	A	401	FAD	C6-C5X-C9A	-2.02	116.30	119.01
3	A	401	FAD	C6-C7-C8	2.09	123.60	119.94
3	A	401	FAD	C9-C9A-C5X	2.28	123.64	119.89
5	B	302	NAD	O5D-PN-O2N	2.51	117.51	107.59
3	B	301	FAD	C6-C5X-N5	2.56	121.88	118.95
3	B	301	FAD	C6-C7-C8	2.85	124.93	119.94
3	A	401	FAD	C4X-C10-N10	2.86	123.06	120.40
3	A	401	FAD	C1'-N10-C9A	2.92	120.90	118.31

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	FAD	O2'-C2'-C1'	2.97	116.93	109.61
3	B	301	FAD	O2P-P-O1P	3.20	128.38	112.14
3	A	401	FAD	C6-C5X-N5	3.40	122.85	118.95
3	A	401	FAD	O4B-C4B-C3B	3.49	112.06	105.15
3	A	401	FAD	C4-N3-C2	3.65	118.25	115.14
3	A	401	FAD	C8M-C8-C7	3.70	128.44	120.72
3	B	301	FAD	C4X-N5-C5X	3.82	120.76	116.76
3	B	301	FAD	O5'-C5'-C4'	4.44	121.20	109.36
3	A	401	FAD	C4-C4X-N5	4.73	124.02	118.70
3	B	301	FAD	C1'-N10-C9A	5.62	123.31	118.31
3	A	401	FAD	C4X-N5-C5X	6.05	123.10	116.76
3	B	301	FAD	C4-C4X-C10	6.27	124.64	119.95
3	B	301	FAD	C4X-C10-N10	7.39	127.28	120.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	FAD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	338/346 (97%)	-0.27	7 (2%) 63 64	12, 19, 39, 96	0
2	B	263/263 (100%)	-0.22	5 (1%) 66 67	12, 19, 38, 67	0
All	All	601/609 (98%)	-0.25	12 (1%) 65 65	12, 19, 39, 96	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	340	SER	7.3
1	A	211	GLU	4.6
1	A	209	VAL	3.4
2	B	22	THR	3.2
2	B	21	VAL	2.6
2	B	151	ASP	2.5
1	A	339	GLU	2.5
1	A	3	ASN	2.2
2	B	20	PRO	2.2
1	A	336	ALA	2.1
2	B	263	LYS	2.1
1	A	213	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAD	B	302	27/44	0.98	0.06	12,15,41,80	1
3	FAD	A	401	53/53	0.98	0.06	11,14,18,20	0
3	FAD	B	301	53/53	0.99	0.07	10,13,16,24	0
4	CL	A	402	1/1	0.99	0.04	23,23,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.