



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 05:40 am GMT

PDB ID : 3LMW  
Title : Crystal structure of iota-carrageenase family GH82 from *A. fortis* in absence of chloride ions  
Authors : Rebuffet, E.; Barbeyron, T.; Jeudy, A.; Czjzek, M.; Michel, G.  
Deposited on : 2010-02-01  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	473	 75% 21% •
1	B	473	 2% 69% 23% • 5%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7652 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iota-carrageenase, CgiA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	472	Total	C	N	O	S	0	4	0
			3768	2354	685	713	16			
1	B	449	Total	C	N	O	S	15	0	0
			3534	2212	629	678	15			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	492	LEU	-	EXPRESSION TAG	UNP Q9F5I8
A	493	GLU	-	EXPRESSION TAG	UNP Q9F5I8
A	494	HIS	-	EXPRESSION TAG	UNP Q9F5I8
A	495	HIS	-	EXPRESSION TAG	UNP Q9F5I8
A	496	HIS	-	EXPRESSION TAG	UNP Q9F5I8
A	497	HIS	-	EXPRESSION TAG	UNP Q9F5I8
A	498	HIS	-	EXPRESSION TAG	UNP Q9F5I8
A	499	HIS	-	EXPRESSION TAG	UNP Q9F5I8
B	492	LEU	-	EXPRESSION TAG	UNP Q9F5I8
B	493	GLU	-	EXPRESSION TAG	UNP Q9F5I8
B	494	HIS	-	EXPRESSION TAG	UNP Q9F5I8
B	495	HIS	-	EXPRESSION TAG	UNP Q9F5I8
B	496	HIS	-	EXPRESSION TAG	UNP Q9F5I8
B	497	HIS	-	EXPRESSION TAG	UNP Q9F5I8
B	498	HIS	-	EXPRESSION TAG	UNP Q9F5I8
B	499	HIS	-	EXPRESSION TAG	UNP Q9F5I8

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ni	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	196	Total	O	0	0
			196	196		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	136	Total 136	O 136	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Iota-carrageenase, CgiA



#### • Molecule 1: Iota-carrageenase, CgiA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.96Å 101.39Å 95.91Å 90.00° 100.40° 90.00°	Depositor
Resolution (Å)	69.06 – 2.60 69.06 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.1 (69.06-2.60) 95.1 (69.06-2.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.79 (at 2.62Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.180 , 0.260 0.177 , 0.257	Depositor DCC
$R_{free}$ test set	1512 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.5	Xtriage
Anisotropy	0.574	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7652	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, CA, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	0/3851	0.87	6/5202 (0.1%)
1	B	0.75	0/3601	0.78	0/4862
All	All	0.75	0/7452	0.83	6/10064 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	2	0

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	SER	CB-CA-C	-15.72	80.24	110.10
1	A	174	ASN	N-CA-C	13.12	146.44	111.00
1	A	184	PHE	N-CA-CB	13.00	133.99	110.60
1	A	174	ASN	N-CA-CB	-9.57	93.38	110.60
1	A	183	ILE	N-CA-C	-7.54	90.65	111.00

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	174	ASN	CA
1	A	184	PHE	CA

There are no planarity outliers.



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3768	0	3655	100	0
1	B	3534	0	3436	87	0
2	A	4	0	0	0	0
2	B	12	0	0	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	196	0	0	8	0
5	B	136	0	0	4	0
All	All	7652	0	7091	182	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:439:PRO:HG2	1:A:468:THR:OG1	1.50	1.09
1:A:440:GLU:O	1:A:441:THR:HG23	1.58	1.00
1:A:437:THR:C	1:A:439:PRO:HD2	1.83	0.98
1:B:329:GLU:HA	1:B:333:GLY:O	1.65	0.94
1:B:137:ASN:ND2	1:B:169:ASN:HD22	1.66	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/473 (100%)	442 (93%)	25 (5%)	7 (2%)	11	23
1	B	441/473 (93%)	402 (91%)	30 (7%)	9 (2%)	8	16
All	All	915/946 (97%)	844 (92%)	55 (6%)	16 (2%)	9	20

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	184	PHE
1	A	248	ASN
1	B	61	ASP
1	B	338	GLN
1	B	440	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/407 (101%)	381 (93%)	30 (7%)	15	31
1	B	386/407 (95%)	365 (95%)	21 (5%)	24	47
All	All	797/814 (98%)	746 (94%)	51 (6%)	20	39

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	437	THR
1	A	454	LYS
1	B	438	THR
1	A	440	GLU
1	A	492	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	499	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	67	ASN
1	B	212	ASN
1	A	395	GLN
1	A	418	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NO3	A	3	-	1,3,3	3.29	1 (100%)	0,3,3	0.00	-
2	NO3	B	1	-	1,3,3	3.20	1 (100%)	0,3,3	0.00	-
2	NO3	B	2	-	1,3,3	3.64	1 (100%)	0,3,3	0.00	-
2	NO3	B	6	-	1,3,3	3.56	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NO3	A	3	-	-	0/0/0/0	0/0/0/0
2	NO3	B	1	-	-	0/0/0/0	0/0/0/0
2	NO3	B	2	-	-	0/0/0/0	0/0/0/0
2	NO3	B	6	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NO3	O1-N	3.20	1.39	1.24
2	A	3	NO3	O1-N	3.29	1.39	1.24
2	B	6	NO3	O1-N	3.56	1.41	1.24
2	B	2	NO3	O1-N	3.64	1.41	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	472/473 (99%)	-0.43	2 (0%) 92 91	7, 21, 47, 62	0
1	B	449/473 (94%)	-0.28	11 (2%) 59 52	7, 24, 73, 87	3 (0%)
All	All	921/946 (97%)	-0.36	13 (1%) 75 71	7, 23, 58, 87	3 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	328	VAL	3.3
1	B	337	ALA	2.9
1	A	439	PRO	2.8
1	B	416	LYS	2.6
1	B	494	HIS	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NO3	B	6	4/4	0.83	0.18	59,60,60,60	0
3	CA	A	9	1/1	0.91	0.19	60,60,60,60	0
2	NO3	B	1	4/4	0.93	0.21	29,29,30,31	0
2	NO3	A	3	4/4	0.97	0.15	26,28,28,29	0
2	NO3	B	2	4/4	0.99	0.10	15,16,16,17	0
4	NI	A	1	1/1	0.99	0.12	23,23,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.