



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 05:40 pm GMT

PDB ID : 3LXJ  
Title : Crystal Structure of the Bromodomain of Human AAA domain containing 2B (ATAD2B)  
Authors : Filippakopoulos, P.; Keates, T.; Picaud, S.; Fedorov, O.; Krojer, T.; Vollmar, M.; Muniz, J.; Pike, A.C.W.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.; Weigelt, J.; Bountra, C.; Knapp, S.; Structural Genomics Consortium (SGC)  
Deposited on : 2010-02-25  
Resolution : 2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1763 (2.36-2.32)
Clashscore	122126	1858 (2.36-2.32)
Ramachandran outliers	120053	1834 (2.36-2.32)
Sidechain outliers	120020	1835 (2.36-2.32)
RSRZ outliers	108989	1737 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	<div> <div style="width: 93%;"></div> <div style="width: 6%;"></div> <div style="width: 1%;"></div> </div> <div>93% 6% .</div>
1	B	136	<div> <div style="width: 94%;"></div> <div style="width: 4%;"></div> <div style="width: 2%;"></div> </div> <div>94% . .</div>
1	C	136	<div> <div style="width: 88%;"></div> <div style="width: 12%;"></div> <div style="width: 1%;"></div> </div> <div>88% 12% .</div>
1	D	136	<div> <div style="width: 89%;"></div> <div style="width: 10%;"></div> <div style="width: 1%;"></div> </div> <div>89% 10% .</div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase family AAA domain-containing protein 2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	135	Total	C	N	O	S	0	0	0
			1063	671	183	204	5			
1	B	135	Total	C	N	O	S	0	0	0
			1054	664	183	202	5			
1	C	135	Total	C	N	O	S	0	0	0
			1073	677	184	207	5			
1	D	134	Total	C	N	O	S	0	0	0
			1078	679	183	211	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	951	SER	-	EXPRESSION TAG	UNP Q9ULI0
B	951	SER	-	EXPRESSION TAG	UNP Q9ULI0
C	951	SER	-	EXPRESSION TAG	UNP Q9ULI0
D	951	SER	-	EXPRESSION TAG	UNP Q9ULI0

- Molecule 2 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C<sub>3</sub>H<sub>8</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	3	1		
2	B	1	Total	C	O	0	0
			4	3	1		
2	C	1	Total	C	O	0	0
			4	3	1		
2	D	1	Total	C	O	0	0
			4	3	1		
2	D	1	Total	C	O	0	0
			4	3	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total	O	0	0
			20	20		
3	B	33	Total	O	0	0
			33	33		
3	C	37	Total	O	0	0
			37	37		
3	D	46	Total	O	0	0
			46	46		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATPase family AAA domain-containing protein 2B

Chain A:  93% 6%





- Molecule 1: ATPase family AAA domain-containing protein 2B

Chain B:  94%




- Molecule 1: ATPase family AAA domain-containing protein 2B

Chain C:  88% 12%



- Molecule 1: ATPase family AAA domain-containing protein 2B

Chain D:  89% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.64Å 79.64Å 204.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.81 – 2.33 37.09 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.0 (40.81-2.33) 99.5 (37.09-2.34)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.66 (at 2.34Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.212 , 0.253 0.206 , 0.239	Depositor DCC
$R_{free}$ test set	1593 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.0	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 12.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.348 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.512 for H, K, L 0.488 for -H-K, K, -L	Depositor
Outliers	1 of 30851 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4424	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.47 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7543e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/1078	0.71	0/1458
1	B	0.53	0/1069	0.67	0/1446
1	C	0.60	0/1088	0.73	2/1470 (0.1%)
1	D	0.57	0/1093	0.67	0/1474
All	All	0.57	0/4328	0.70	2/5848 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	977	ASP	CB-CG-OD2	-5.60	113.26	118.30
1	C	977	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	951	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1063	0	1030	4	0
1	B	1054	0	1010	3	0
1	C	1073	0	1053	7	0
1	D	1078	0	1065	5	0
2	A	4	0	8	0	0
2	B	4	0	8	0	0
2	C	4	0	8	0	0
2	D	8	0	16	0	0
3	A	20	0	0	0	0
3	B	33	0	0	0	0
3	C	37	0	0	0	0
3	D	46	0	0	0	0
All	All	4424	0	4198	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1022:ASP:OD2	1:B:1079:LYS:NZ	2.23	0.69
1:D:1044:GLY:HA2	1:D:1047:ILE:HD12	1.89	0.55
1:D:1042:ASP:HB2	1:D:1043:PRO:CD	2.39	0.52
1:C:1011:LYS:HG2	1:C:1016:ASN:HD22	1.77	0.49
1:B:966:PHE:CZ	1:B:1063:ILE:HD12	2.48	0.49

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/136 (98%)	126 (95%)	7 (5%)	0	100	100
1	B	133/136 (98%)	127 (96%)	5 (4%)	1 (1%)	21	22
1	C	133/136 (98%)	130 (98%)	3 (2%)	0	100	100
1	D	132/136 (97%)	128 (97%)	3 (2%)	1 (1%)	21	22
All	All	531/544 (98%)	511 (96%)	18 (3%)	2 (0%)	36	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	993	SER
1	B	1040	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/125 (88%)	109 (99%)	1 (1%)	81	88
1	B	107/125 (86%)	104 (97%)	3 (3%)	47	57
1	C	114/125 (91%)	112 (98%)	2 (2%)	62	73
1	D	117/125 (94%)	113 (97%)	4 (3%)	40	50
All	All	448/500 (90%)	438 (98%)	10 (2%)	55	67

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	952	MET
1	C	988	ASP
1	D	997	GLU
1	B	1063	ILE
1	D	952	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1016	ASN
1	A	1072	ASN
1	B	1016	ASN
1	C	1016	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	IPA	A	4	-	3,3,3	0.67	0	3,3,3	0.65	0
2	IPA	B	3	-	3,3,3	0.60	0	3,3,3	0.62	0
2	IPA	C	2	-	3,3,3	0.65	0	3,3,3	0.58	0
2	IPA	D	1	-	3,3,3	0.68	0	3,3,3	0.33	0
2	IPA	D	5	-	3,3,3	0.62	0	3,3,3	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IPA	A	4	-	-	0/0/0/0	0/0/0/0
2	IPA	B	3	-	-	0/0/0/0	0/0/0/0
2	IPA	C	2	-	-	0/0/0/0	0/0/0/0
2	IPA	D	1	-	-	0/0/0/0	0/0/0/0
2	IPA	D	5	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	135/136 (99%)	-0.71	0 100 100	28, 44, 72, 82	0
1	B	135/136 (99%)	-0.59	0 100 100	32, 46, 86, 103	0
1	C	135/136 (99%)	-0.59	0 100 100	24, 38, 72, 91	0
1	D	134/136 (98%)	-0.61	0 100 100	31, 43, 72, 88	0
All	All	539/544 (99%)	-0.62	0 100 100	24, 43, 79, 103	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	IPA	C	2	4/4	0.93	0.11	33,37,40,41	0
2	IPA	D	1	4/4	0.96	0.11	34,37,38,39	0
2	IPA	A	4	4/4	0.96	0.08	31,31,34,52	0
2	IPA	B	3	4/4	0.96	0.12	35,36,41,46	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	IPA	D	5	4/4	0.98	0.14	33,34,45,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.