



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 18, 2019 – 04:45 PM EDT

PDB ID : 4LX4  
Title : Crystal Structure Determination of Pseudomonas stutzeri endoglucanase Cel5A using a Twinned Data Set  
Authors : Dutoit, R.; Delsaute, M.; Berlemont, R.; Van Elder, D.; Galleni, M.; Bauvois, C.  
Deposited on : 2013-07-29  
Resolution : 1.56 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.3.2  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.3.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1224 (1.56-1.56)
Clashscore	122126	1265 (1.56-1.56)
Ramachandran outliers	120053	1240 (1.56-1.56)
Sidechain outliers	120020	1238 (1.56-1.56)
RSRZ outliers	108989	1207 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div> <div>3%</div> <div>80%</div> <div>17%</div> <div>..</div> </div>
1	B	330	<div> <div>6%</div> <div>80%</div> <div>18%</div> <div>..</div> </div>
1	C	330	<div> <div>7%</div> <div>77%</div> <div>20%</div> <div>.</div> </div>
1	D	330	<div> <div>7%</div> <div>76%</div> <div>20%</div> <div>..</div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 21329 atoms, of which 9851 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoglucanase(Endo-1,4-beta-glucanase)protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	322	Total	C	H	N	O	S	0	0	0
			5019	1651	2444	448	467	9			
1	B	328	Total	C	H	N	O	S	0	0	0
			5113	1678	2494	456	476	9			
1	C	321	Total	C	H	N	O	S	0	0	0
			5022	1651	2451	447	464	9			
1	D	319	Total	C	H	N	O	S	0	0	0
			4964	1637	2414	440	464	9			

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
2	B	1	Total	C	H	N	O	0	0
			20	4	12	1	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
2	D	1	Total	C	H	N	O	0	0
			20	4	12	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	258	Total	O	0	0
			258	258		
3	B	312	Total	O	0	0
			312	312		
3	C	276	Total	O	0	0
			276	276		
3	D	285	Total	O	0	0
			285	285		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Endoglucanase(Endo-1,4-beta-glucanase)protein




- Molecule 1: Endoglucanase(Endo-1,4-beta-glucanase)protein



- Molecule 1: Endoglucanase(Endo-1,4-beta-glucanase)protein



● Molecule 1: Endoglucanase(Endo-1,4-beta-glucanase)protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.39Å 82.90Å 104.74Å 90.00° 92.08° 90.00°	Depositor
Resolution (Å)	48.28 – 1.56 48.28 – 1.56	Depositor EDS
% Data completeness (in resolution range)	91.6 (48.28-1.56) 82.3 (48.28-1.56)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.41 (at 1.55Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.220 , 0.237 0.224 , 0.236	Depositor DCC
$R_{free}$ test set	7883 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.4	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 23.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.71$ , $\langle L^2 \rangle = 0.60$	Xtriage
Estimated twinning fraction	0.126 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	21329	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	2/2655 (0.1%)	0.60	2/3607 (0.1%)
1	B	0.41	2/2700 (0.1%)	0.56	2/3668 (0.1%)
1	C	0.40	1/2651 (0.0%)	0.58	2/3601 (0.1%)
1	D	0.38	1/2629 (0.0%)	0.56	1/3573 (0.0%)
All	All	0.40	6/10635 (0.1%)	0.57	7/14449 (0.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	279	PRO	N-CD	5.51	1.55	1.47
1	D	134	PRO	N-CD	5.36	1.55	1.47
1	C	153	PRO	N-CD	5.25	1.55	1.47
1	B	236	PRO	N-CD	5.16	1.55	1.47
1	A	256	PRO	N-CD	5.09	1.54	1.47

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	255	ILE	C-N-CD	5.81	140.60	128.40
1	A	255	ILE	C-N-CD	5.76	140.50	128.40
1	A	133	HIS	C-N-CD	5.70	140.38	128.40
1	B	235	ARG	C-N-CD	5.64	140.24	128.40
1	C	152	TRP	C-N-CD	5.50	139.96	128.40

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2575	2444	2457	38	1
1	B	2619	2494	2507	53	0
1	C	2571	2451	2464	52	0
1	D	2550	2414	2428	47	1
2	A	8	12	12	0	0
2	B	8	12	12	0	0
2	C	8	12	12	0	0
2	D	8	12	12	0	0
3	A	258	0	0	6	0
3	B	312	0	0	13	1
3	C	276	0	0	11	2
3	D	285	0	0	10	2
All	All	11478	9851	9904	186	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 186 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:246:GLN:HA	3:D:742:HOH:O	1.78	0.82
1:B:256:PRO:HG2	1:B:259:LEU:HG	1.62	0.82
1:A:172:ILE:HB	1:A:201:LEU:CD2	2.11	0.81
1:B:113:LYS:HB3	3:B:789:HOH:O	1.79	0.80
1:A:220:GLN:HB3	3:A:690:HOH:O	1.82	0.79

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:GLU:HG2	1:D:319:THR:HG1[2_556]	1.32	0.28
3:C:776:HOH:O	3:D:748:HOH:O[1_565]	2.03	0.17
3:C:765:HOH:O	3:D:780:HOH:O[1_565]	2.07	0.13
3:B:586:HOH:O	3:B:608:HOH:O[2_557]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/330 (97%)	317 (99%)	3 (1%)	0	100	100
1	B	326/330 (99%)	321 (98%)	5 (2%)	0	100	100
1	C	317/330 (96%)	311 (98%)	6 (2%)	0	100	100
1	D	315/330 (96%)	312 (99%)	3 (1%)	0	100	100
All	All	1278/1320 (97%)	1261 (99%)	17 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/267 (98%)	258 (99%)	3 (1%)	76	55
1	B	266/267 (100%)	263 (99%)	3 (1%)	76	55
1	C	262/267 (98%)	259 (99%)	3 (1%)	76	55
1	D	259/267 (97%)	257 (99%)	2 (1%)	83	69
All	All	1048/1068 (98%)	1037 (99%)	11 (1%)	78	60

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	178	ARG
1	B	221	THR
1	C	180	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	97	ASP
1	C	178	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	TRS	A	401	-	7,7,7	0.28	0	9,9,9	0.65	0
2	TRS	B	401	-	7,7,7	0.53	0	9,9,9	1.04	0
2	TRS	C	401	-	7,7,7	0.23	0	9,9,9	0.39	0
2	TRS	D	401	-	7,7,7	0.25	0	9,9,9	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	A	401	-	-	3/9/9/9	-
2	TRS	B	401	-	-	3/9/9/9	-
2	TRS	C	401	-	-	7/9/9/9	-
2	TRS	D	401	-	-	2/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	TRS	C1-C-C3-O3
2	A	401	TRS	C2-C-C3-O3
2	A	401	TRS	N-C-C3-O3
2	C	401	TRS	C2-C-C1-O1
2	C	401	TRS	C3-C-C1-O1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/330 (97%)	0.68	10 (3%) 49 57	6, 11, 18, 25	0
1	B	328/330 (99%)	0.82	21 (6%) 19 22	6, 11, 18, 24	0
1	C	321/330 (97%)	0.89	23 (7%) 15 18	6, 12, 20, 26	0
1	D	319/330 (96%)	0.88	24 (7%) 14 16	5, 12, 19, 28	0
All	All	1290/1320 (97%)	0.82	78 (6%) 22 25	5, 11, 19, 28	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	LEU	5.8
1	B	329	ALA	5.4
1	D	74	PHE	4.5
1	B	328	ILE	4.0
1	C	225	ILE	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	TRS	A	401	8/8	0.83	0.20	8,16,24,29	0
2	TRS	C	401	8/8	0.85	0.14	8,12,15,19	0
2	TRS	B	401	8/8	0.87	0.14	8,12,16,18	0
2	TRS	D	401	8/8	0.91	0.12	9,12,15,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.