



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 06:24 am GMT

PDB ID : 5LXC
Title : Crystal structure of DYRK2 in complex with EHT 5372 (Compound 1)
Authors : Chaikuad, A.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.; Bountra, C.;
Besson, T.; Knapp, S.; Structural Genomics Consortium (SGC)
Deposited on : 2016-09-20
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1287 (2.16-2.16)
Clashscore	122126	1390 (2.16-2.16)
Ramachandran outliers	120053	1368 (2.16-2.16)
Sidechain outliers	120020	1367 (2.16-2.16)
RSRZ outliers	108989	1262 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	408	 % 85% 9% 6%
1	B	408	 6% 86% 8% 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dual specificity tyrosine-phosphorylation-regulated kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	P	S	0	5	0
			3128	1991	557	556	1	23			
1	B	386	Total	C	N	O	P	S	0	5	0
			3141	1997	564	556	1	23			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	72	SER	-	expression tag	UNP Q92630
B	72	SER	-	expression tag	UNP Q92630

- Molecule 2 is methyl 9-[(2,4-dichlorophenyl)amino]-[1,3]thiazolo[5,4-f]quinazoline-2-carboximidate (three-letter code: 7AA) (formula: C₁₇H₁₁Cl₂N₅OS).



Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	Cl	N	O	S	0	0
			26	17	2	5	1	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	234	Total	O	0	0
			234	234		
4	B	162	Total	O	0	0
			162	162		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 2



- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	130.22Å 60.98Å 148.79Å 90.00° 105.04° 90.00°	Depositor
Resolution (Å)	49.43 – 2.15 49.43 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.3 (49.43-2.15) 97.3 (49.43-2.15)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.206 , 0.254 0.214 , 0.259	Depositor DCC
R_{free} test set	3047 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6761	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, 7AA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	2/3200 (0.1%)	0.79	4/4308 (0.1%)
1	B	0.72	2/3213 (0.1%)	0.74	3/4324 (0.1%)
All	All	0.76	4/6413 (0.1%)	0.77	7/8632 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	450	ARG	CZ-NH2	-11.91	1.17	1.33
1	B	450	ARG	CZ-NH2	-9.84	1.20	1.33
1	A	450	ARG	CZ-NH1	-9.52	1.20	1.33
1	B	450	ARG	CZ-NH1	-8.78	1.21	1.33

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	450	ARG	NE-CZ-NH1	18.57	129.59	120.30
1	A	450	ARG	NH1-CZ-NH2	-11.23	107.05	119.40
1	B	450	ARG	NE-CZ-NH1	9.57	125.08	120.30
1	B	450	ARG	NE-CZ-NH2	8.98	124.79	120.30
1	B	450	ARG	NH1-CZ-NH2	-8.49	110.06	119.40
1	A	450	ARG	NE-CZ-NH2	6.07	123.33	120.30
1	A	313	ARG	NE-CZ-NH2	-5.56	117.52	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3128	0	3125	21	0
1	B	3141	0	3143	24	0
2	A	26	0	0	2	0
2	B	26	0	0	1	0
3	A	32	0	48	3	0
3	B	12	0	18	1	0
4	A	234	0	0	3	0
4	B	162	0	0	5	0
All	All	6761	0	6334	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:501:7AA:SAP	2:B:501:7AA:NAN	2.64	0.70
2:A:501:7AA:SAP	2:A:501:7AA:NAN	2.72	0.62
3:A:508:EDO:H21	1:B:411:ARG:HD3	1.82	0.61
1:B:110:PHE:O	1:B:147:TYR:HA	2.00	0.61
1:A:87:MET:O	1:A:91:MET:HB2	2.00	0.61
1:B:165:LYS:HE3	4:B:637:HOH:O	2.00	0.61
1:B:84:GLU:N	1:B:84:GLU:OE1	2.35	0.59
1:B:270:ARG:NH2	1:B:302:GLU:OE1	2.34	0.57
1:A:407:GLY:O	1:A:414:LEU:HD12	2.07	0.54
1:A:436:LEU:HG	1:A:440:LYS:HE3	1.89	0.54
1:B:291:ILE:C	1:B:291:ILE:HD12	2.27	0.54
1:B:436:LEU:HG	1:B:440:LYS:HE3	1.89	0.54
1:A:270:ARG:NH2	1:A:302:GLU:OE1	2.40	0.52
1:B:350:PRO:HG2	4:B:707:HOH:O	2.08	0.52
1:A:352:GLU:CG	1:B:411:ARG:HH12	2.24	0.50
1:A:142:HIS:ND1	1:A:150:GLU:OE2	2.35	0.50
1:B:185:ARG:HB2	4:B:602:HOH:O	2.11	0.50
1:B:456:ALA:O	1:B:462:LEU:HD12	2.12	0.49
1:B:396:THR:HA	1:B:401:SER:O	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:413:LYS:HG2	3:A:508:EDO:H11	1.95	0.48
1:A:456:ALA:O	1:A:462:LEU:HD12	2.13	0.48
1:A:80:PRO:HA	4:A:696:HOH:O	2.12	0.48
1:A:232:SER:OG	1:A:233:MET:N	2.43	0.47
1:A:180:VAL:HG12	1:A:181:ARG:O	2.16	0.46
1:A:282:LEU:HD22	1:A:294:ILE:HG21	1.97	0.46
1:A:240:LYS:HE2	4:A:821:HOH:O	2.16	0.46
1:B:328:MET:N	1:B:329:PRO:CD	2.80	0.45
1:B:273:HIS:O	1:B:274:CYS:HB2	2.17	0.45
1:A:132:ASP:OD2	1:A:134:GLN:N	2.48	0.44
1:A:428:LYS:HD2	4:A:825:HOH:O	2.18	0.43
1:B:110:PHE:CZ	1:B:150:GLU:HG3	2.52	0.43
1:B:137:TYR:HB2	1:B:223:HIS:NE2	2.34	0.43
1:A:291:ILE:HD12	1:A:291:ILE:C	2.39	0.43
1:A:369:SER:OG	1:A:372[B]:LEU:HG	2.19	0.42
1:A:352:GLU:HG3	1:B:411:ARG:HH12	1.84	0.42
3:B:502:EDO:H11	4:B:719:HOH:O	2.19	0.42
1:B:452:THR:OG1	1:B:455:GLN:HG3	2.20	0.42
1:B:291:ILE:HD12	1:B:291:ILE:O	2.19	0.42
1:B:458[A]:ARG:NE	4:B:610:HOH:O	2.49	0.42
1:A:83:PRO:HB3	1:A:102:ILE:HG12	2.02	0.42
1:B:282:LEU:HD22	1:B:294:ILE:HG21	2.01	0.41
1:B:394:VAL:CG2	1:B:402:VAL:HG13	2.49	0.41
1:A:444[A]:GLU:HG2	1:A:449:VAL:HB	2.02	0.41
1:B:459:HIS:CG	1:B:460:PRO:HD2	2.55	0.41
1:B:108:ILE:HD12	1:B:147:TYR:CG	2.56	0.41
2:A:501:7AA:CAG	3:A:504:EDO:H22	2.52	0.40
1:A:270:ARG:HA	1:A:328:MET:SD	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/408 (95%)	370 (96%)	14 (4%)	2 (0%)	31	24
1	B	388/408 (95%)	361 (93%)	27 (7%)	0	100	100
All	All	774/816 (95%)	731 (94%)	41 (5%)	2 (0%)	43	39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	398	SER
1	A	406	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/355 (95%)	337 (100%)	0	100	100
1	B	338/355 (95%)	338 (100%)	0	100	100
All	All	675/710 (95%)	675 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	309	1	16,16,17	0.76	0	21,22,24	1.05	2 (9%)
1	PTR	B	309	1	16,16,17	1.14	0	21,22,24	0.93	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	309	1	-	0/9/11/13	0/1/1/1
1	PTR	B	309	1	-	0/9/11/13	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	309	PTR	O3P-P-OH	-2.05	98.58	105.50
1	A	309	PTR	O3P-P-OH	-2.04	98.60	105.50
1	A	309	PTR	O3P-P-O2P	2.67	118.15	107.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	7AA	A	501	-	23,29,29	4.02	9 (39%)	23,42,42	2.30	9 (39%)
3	EDO	A	502	-	3,3,3	0.67	0	2,2,2	0.30	0
3	EDO	A	503	-	3,3,3	0.46	0	2,2,2	0.48	0
3	EDO	A	504	-	3,3,3	0.27	0	2,2,2	0.52	0
3	EDO	A	505	-	3,3,3	0.46	0	2,2,2	0.39	0
3	EDO	A	506	-	3,3,3	0.89	0	2,2,2	0.28	0
3	EDO	A	507	-	3,3,3	0.50	0	2,2,2	0.27	0
3	EDO	A	508	-	3,3,3	0.29	0	2,2,2	0.47	0
3	EDO	A	509	-	3,3,3	0.52	0	2,2,2	0.36	0
2	7AA	B	501	-	23,29,29	3.84	9 (39%)	23,42,42	2.26	9 (39%)
3	EDO	B	502	-	3,3,3	0.62	0	2,2,2	0.25	0
3	EDO	B	503	-	3,3,3	0.57	0	2,2,2	0.29	0
3	EDO	B	504	-	3,3,3	0.56	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7AA	A	501	-	-	0/5/20/20	0/4/4/4
3	EDO	A	502	-	-	0/1/1/1	0/0/0/0
3	EDO	A	503	-	-	0/1/1/1	0/0/0/0
3	EDO	A	504	-	-	0/1/1/1	0/0/0/0
3	EDO	A	505	-	-	0/1/1/1	0/0/0/0
3	EDO	A	506	-	-	0/1/1/1	0/0/0/0
3	EDO	A	507	-	-	0/1/1/1	0/0/0/0
3	EDO	A	508	-	-	0/1/1/1	0/0/0/0
3	EDO	A	509	-	-	0/1/1/1	0/0/0/0
2	7AA	B	501	-	-	0/5/20/20	0/4/4/4
3	EDO	B	502	-	-	0/1/1/1	0/0/0/0
3	EDO	B	503	-	-	0/1/1/1	0/0/0/0
3	EDO	B	504	-	-	0/1/1/1	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	7AA	CAV-CAQ	-10.16	1.31	1.48
2	A	501	7AA	CAV-CAQ	-10.01	1.31	1.48
2	B	501	7AA	CAV-SAP	-9.54	1.60	1.73
2	A	501	7AA	CAV-SAP	-8.52	1.61	1.73
2	B	501	7AA	C6-NAN	-7.34	1.33	1.45
2	A	501	7AA	CAI-CAX	-6.51	1.30	1.41
2	A	501	7AA	C6-NAN	-6.12	1.35	1.45
2	B	501	7AA	CAI-CAX	-4.08	1.34	1.41
2	A	501	7AA	C2-N3	-3.12	1.41	1.45
2	B	501	7AA	C4-N3	-2.52	1.32	1.37
2	B	501	7AA	C2-N3	-2.44	1.42	1.45
2	A	501	7AA	CAX-CAY	-2.23	1.34	1.42
2	B	501	7AA	CAR-CL2	3.50	1.82	1.74
2	A	501	7AA	OAQ-CAA	3.58	1.53	1.45
2	B	501	7AA	CAQ-NAB	4.06	1.35	1.25
2	B	501	7AA	C4-C5	4.09	1.40	1.38
2	A	501	7AA	CAQ-NAB	4.42	1.35	1.25
2	A	501	7AA	C4-C5	7.04	1.41	1.38

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	7AA	CAX-CAY-SAP	-3.98	107.10	112.00
2	A	501	7AA	CAX-CAY-SAP	-3.84	107.27	112.00
2	B	501	7AA	C2-N3-C4	-3.29	117.96	122.28
2	A	501	7AA	C2-N3-C4	-3.15	118.14	122.28
2	B	501	7AA	CAH-C4-C5	-2.63	119.18	121.16
2	A	501	7AA	CAH-C4-C5	-2.40	119.36	121.16
2	A	501	7AA	CAI-CAH-C4	-2.12	116.12	120.05
2	B	501	7AA	CAS-CAJ-CAR	2.42	121.34	118.72
2	A	501	7AA	CAH-CAI-CAX	2.56	124.25	120.84
2	B	501	7AA	CAT-CAS-CL1	2.66	122.69	119.54
2	B	501	7AA	C5-C6-NAN	2.67	116.87	111.04
2	B	501	7AA	CAF-CAR-CL2	2.85	124.08	119.35
2	A	501	7AA	C5-C6-NAN	3.26	118.14	111.04
2	A	501	7AA	CAT-CAS-CL1	3.66	123.89	119.54
2	A	501	7AA	CAV-NAM-CAX	3.77	111.22	103.83
2	B	501	7AA	CAV-NAM-CAX	4.56	112.77	103.83
2	B	501	7AA	N1-C2-N3	4.57	121.76	112.00
2	A	501	7AA	N1-C2-N3	4.94	122.54	112.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	7AA	2	0
3	A	504	EDO	1	0
3	A	508	EDO	2	0
2	B	501	7AA	1	0
3	B	502	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/408 (93%)	0.00	5 (1%) 77 81	21, 41, 72, 109	0
1	B	385/408 (94%)	0.36	23 (5%) 22 28	21, 50, 92, 109	0
All	All	768/816 (94%)	0.18	28 (3%) 42 50	21, 45, 87, 109	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	93	LYS	3.9
1	B	171	VAL	3.9
1	B	125	PRO	3.6
1	A	402	VAL	3.5
1	B	288	ARG	3.4
1	B	124	GLY	3.1
1	B	172	HIS	3.1
1	B	223	HIS	3.0
1	A	400	GLY	2.9
1	B	151	VAL	2.9
1	B	106	PRO	2.9
1	B	186	PHE	2.8
1	A	124	GLY	2.6
1	B	152	LEU	2.5
1	B	82	THR	2.5
1	B	220	PHE	2.5
1	B	159	SER	2.4
1	B	184	LYS	2.4
1	B	113	LEU	2.4
1	B	147	TYR	2.3
1	B	140	VAL	2.3
1	A	396	THR	2.1
1	B	400	GLY	2.1
1	B	80	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	206	ASP	2.0
1	A	403	VAL	2.0
1	B	119	GLN	2.0
1	B	141	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	B	309	16/17	0.95	0.13	36,42,44,47	0
1	PTR	A	309	16/17	0.96	0.12	36,47,50,54	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	507	4/4	0.82	0.22	67,70,71,72	0
3	EDO	A	502	4/4	0.86	0.14	52,56,57,59	0
3	EDO	A	505	4/4	0.87	0.14	69,71,72,73	0
3	EDO	B	504	4/4	0.90	0.28	57,65,65,70	0
3	EDO	A	509	4/4	0.90	0.19	60,61,61,65	0
3	EDO	B	502	4/4	0.93	0.26	47,58,59,62	0
3	EDO	B	503	4/4	0.94	0.27	62,62,62,64	0
3	EDO	A	508	4/4	0.94	0.13	38,40,41,45	0
3	EDO	A	503	4/4	0.94	0.11	36,38,38,46	0
3	EDO	A	506	4/4	0.96	0.11	39,40,44,48	0
3	EDO	A	504	4/4	0.97	0.18	41,43,44,45	0
2	7AA	B	501	26/26	0.97	0.11	39,45,68,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	7AA	A	501	26/26	0.97	0.11	22,25,41,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.