



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 30, 2020 – 08:42 AM EDT

PDB ID : 6LXT  
Title : Structure of post fusion core of 2019-nCoV S2 subunit  
Authors : Zhu, Y.; Sun, F.  
Deposited on : 2020-02-11  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.10.1  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.10.1

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1716 (2.90-2.90)
Clashscore	122126	1924 (2.90-2.90)
Ramachandran outliers	120053	1884 (2.90-2.90)
Sidechain outliers	120020	1886 (2.90-2.90)
RSRZ outliers	108989	1669 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	 2% 72% 14% • 12%
1	B	132	 2% 73% 13% • 12%
1	C	132	 2% 77% 12% 11%
1	D	132	 4% 69% 14% • 17%
1	E	132	 7% 76% 11% 14%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	132	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PG4	A	1303	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S2, Spike protein S2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	0	0	0
			881	543	153	185			
1	B	116	Total	C	N	O	0	0	0
			880	543	152	185			
1	C	117	Total	C	N	O	0	0	0
			889	547	154	188			
1	D	110	Total	C	N	O	0	0	0
			831	511	146	174			
1	E	114	Total	C	N	O	0	0	0
			865	534	149	182			
1	F	113	Total	C	N	O	0	0	0
			859	530	150	179			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1156	SER	-	linker	UNP P0DTC2
A	1157	GLY	-	linker	UNP P0DTC2
A	1158	GLY	-	linker	UNP P0DTC2
A	1159	ARG	-	linker	UNP P0DTC2
A	1160	GLY	-	linker	UNP P0DTC2
A	1161	GLY	-	linker	UNP P0DTC2
A	1207	GLY	-	expression tag	UNP P0DTC2
A	1208	GLY	-	expression tag	UNP P0DTC2
B	1156	SER	-	linker	UNP P0DTC2
B	1157	GLY	-	linker	UNP P0DTC2
B	1158	GLY	-	linker	UNP P0DTC2
B	1159	ARG	-	linker	UNP P0DTC2
B	1160	GLY	-	linker	UNP P0DTC2
B	1161	GLY	-	linker	UNP P0DTC2
B	1207	GLY	-	expression tag	UNP P0DTC2
B	1208	GLY	-	expression tag	UNP P0DTC2
C	1156	SER	-	linker	UNP P0DTC2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	1157	GLY	-	linker	UNP P0DTC2
C	1158	GLY	-	linker	UNP P0DTC2
C	1159	ARG	-	linker	UNP P0DTC2
C	1160	GLY	-	linker	UNP P0DTC2
C	1161	GLY	-	linker	UNP P0DTC2
C	1207	GLY	-	expression tag	UNP P0DTC2
C	1208	GLY	-	expression tag	UNP P0DTC2
D	1156	SER	-	linker	UNP P0DTC2
D	1157	GLY	-	linker	UNP P0DTC2
D	1158	GLY	-	linker	UNP P0DTC2
D	1159	ARG	-	linker	UNP P0DTC2
D	1160	GLY	-	linker	UNP P0DTC2
D	1161	GLY	-	linker	UNP P0DTC2
D	1207	GLY	-	expression tag	UNP P0DTC2
D	1208	GLY	-	expression tag	UNP P0DTC2
E	1156	SER	-	linker	UNP P0DTC2
E	1157	GLY	-	linker	UNP P0DTC2
E	1158	GLY	-	linker	UNP P0DTC2
E	1159	ARG	-	linker	UNP P0DTC2
E	1160	GLY	-	linker	UNP P0DTC2
E	1161	GLY	-	linker	UNP P0DTC2
E	1207	GLY	-	expression tag	UNP P0DTC2
E	1208	GLY	-	expression tag	UNP P0DTC2
F	1156	SER	-	linker	UNP P0DTC2
F	1157	GLY	-	linker	UNP P0DTC2
F	1158	GLY	-	linker	UNP P0DTC2
F	1159	ARG	-	linker	UNP P0DTC2
F	1160	GLY	-	linker	UNP P0DTC2
F	1161	GLY	-	linker	UNP P0DTC2
F	1207	GLY	-	expression tag	UNP P0DTC2
F	1208	GLY	-	expression tag	UNP P0DTC2

- Molecule 2 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	8	5		
2	A	1	Total	C	O	0	0
			13	8	5		


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		
3	F	2	Total	Zn	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S2, Spike protein S2




- Molecule 1: Spike protein S2, Spike protein S2




- Molecule 1: Spike protein S2, Spike protein S2




- Molecule 1: Spike protein S2, Spike protein S2



- Molecule 1: Spike protein S2, Spike protein S2



● Molecule 1: Spike protein S2, Spike protein S2





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.24Å 57.58Å 115.72Å 90.00° 91.58° 90.00°	Depositor
Resolution (Å)	47.32 – 2.90 47.32 – 2.90	Depositor EDS
% Data completeness (in resolution range)	94.0 (47.32-2.90) 94.0 (47.32-2.90)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.259 , 0.290 0.258 , 0.277	Depositor DCC
$R_{free}$ test set	737 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.6	Xtriage
Anisotropy	0.188	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.54$ , $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.024 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5237	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	0/882	0.78	0/1191
1	B	0.67	0/882	0.75	0/1191
1	C	0.69	0/890	0.75	0/1202
1	D	0.70	0/832	0.78	0/1123
1	E	0.70	0/866	0.77	0/1169
1	F	0.69	0/860	0.78	0/1160
All	All	0.69	0/5212	0.77	0/7036

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	881	0	896	26	0
1	B	880	0	893	22	0
1	C	889	0	900	21	0
1	D	831	0	845	23	0
1	E	865	0	879	16	0
1	F	859	0	876	17	0
2	A	26	0	36	14	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	F	2	0	0	0	0
All	All	5237	0	5325	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1184:ASP:OD2	2:A:1303:PG4:H52	1.51	1.09
1:A:920:GLN:HG2	1:C:920:GLN:HE22	1.25	1.01
1:B:920:GLN:NE2	1:C:920:GLN:HG2	1.85	0.91
1:A:1188:GLU:HG2	2:A:1303:PG4:O1	1.71	0.91
1:B:920:GLN:HE21	1:C:920:GLN:HE21	1.20	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/132 (85%)	112 (100%)	0	0	100	100
1	B	112/132 (85%)	112 (100%)	0	0	100	100
1	C	113/132 (86%)	110 (97%)	3 (3%)	0	100	100
1	D	106/132 (80%)	104 (98%)	2 (2%)	0	100	100
1	E	110/132 (83%)	108 (98%)	2 (2%)	0	100	100
1	F	109/132 (83%)	102 (94%)	6 (6%)	1 (1%)	19	52
All	All	662/792 (84%)	648 (98%)	13 (2%)	1 (0%)	49	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	1200	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/110 (93%)	95 (93%)	7 (7%)	17	44
1	B	102/110 (93%)	94 (92%)	8 (8%)	14	37
1	C	103/110 (94%)	98 (95%)	5 (5%)	27	61
1	D	96/110 (87%)	91 (95%)	5 (5%)	25	59
1	E	100/110 (91%)	95 (95%)	5 (5%)	27	60
1	F	99/110 (90%)	95 (96%)	4 (4%)	34	69
All	All	602/660 (91%)	568 (94%)	34 (6%)	23	55

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	914	ASN
1	C	1199	ASP
1	F	1175	SER
1	C	985	ASP
1	A	1201	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	935	GLN
1	C	978	ASN
1	D	949	GLN
1	C	920	GLN
1	F	926	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PG4	A	1303	-	12,12,12	0.46	0	11,11,11	1.61	3 (27%)
2	PG4	A	1301	-	12,12,12	0.73	0	11,11,11	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	A	1303	-	-	6/10/10/10	-
2	PG4	A	1301	-	-	5/10/10/10	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1303	PG4	O2-C3-C4	-3.52	94.53	110.39
2	A	1303	PG4	O5-C8-C7	-2.01	100.14	111.81
2	A	1303	PG4	O4-C7-C8	2.00	118.87	110.07

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1301	PG4	O3-C5-C6-O4
2	A	1303	PG4	O4-C7-C8-O5
2	A	1303	PG4	O1-C1-C2-O2
2	A	1301	PG4	O2-C3-C4-O3
2	A	1301	PG4	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1303	PG4	13	0
2	A	1301	PG4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	116/132 (87%)	0.02	2 (1%) 70 69	49, 75, 124, 142	0
1	B	116/132 (87%)	0.01	3 (2%) 56 52	42, 72, 114, 124	0
1	C	117/132 (88%)	-0.01	2 (1%) 70 69	49, 71, 119, 148	0
1	D	110/132 (83%)	0.18	5 (4%) 33 29	68, 97, 137, 142	0
1	E	114/132 (86%)	0.43	9 (7%) 12 9	67, 98, 143, 155	0
1	F	113/132 (85%)	-0.04	3 (2%) 54 50	48, 82, 124, 135	0
All	All	686/792 (86%)	0.10	24 (3%) 44 38	42, 84, 129, 155	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	912	THR	6.2
1	E	939	SER	3.8
1	D	1196	SER	3.5
1	E	1202	GLU	3.5
1	D	1165	ASP	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	A	1302	1/1	0.63	0.21	150,150,150,150	0
3	ZN	F	1301	1/1	0.79	0.07	148,148,148,148	0
2	PG4	A	1301	13/13	0.82	0.20	54,67,84,84	0
3	ZN	A	1304	1/1	0.88	0.15	117,117,117,117	0
2	PG4	A	1303	13/13	0.93	0.32	41,53,83,83	0
3	ZN	F	1302	1/1	0.93	0.12	145,145,145,145	0
3	ZN	B	1302	1/1	0.94	0.07	145,145,145,145	0
3	ZN	B	1301	1/1	0.97	0.13	103,103,103,103	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.