



Full wwPDB NMR Structure Validation Report ⓘ

Feb 18, 2018 – 07:32 am GMT

PDB ID : 2M3V
Title : Solution structure of tyrosine phosphatase related to biofilm formation A (TpbA) from *Pseudomonas aeruginosa*
Authors : Koveal, D.; Peti, W.; Page, R.
Deposited on : 2013-01-26

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk30686
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 73%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136279	12091
Ramachandran outliers	132675	10835
Sidechain outliers	132484	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	193	 67% 10% • 22%

2 Ensemble composition and analysis

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:37-A:167, A:175-A:184 (141)	0.53	4
2	A:185-A:193 (9)	0.13	15

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 9, 10, 11, 13, 18
2	6, 12, 15, 20
3	7, 8, 14
4	16, 19
Single-model clusters	17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2962 atoms, of which 1467 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms						Trace
1	A	193	Total	C	H	N	O	S	0
			2962	926	1467	281	277	11	

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	26	GLY	-	EXPRESSION TAG	UNP Q02S50
A	27	HIS	-	EXPRESSION TAG	UNP Q02S50
A	28	MET	-	EXPRESSION TAG	UNP Q02S50

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Putative uncharacterized protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Putative uncharacterized protein



4.2.2 Score per residue for model 2


- Molecule 1: Putative uncharacterized protein






4.2.3 Score per residue for model 3

- Molecule 1: Putative uncharacterized protein




4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: Putative uncharacterized protein



4.2.5 Score per residue for model 5

- Molecule 1: Putative uncharacterized protein




4.2.6 Score per residue for model 6

- Molecule 1: Putative uncharacterized protein




4.2.7 Score per residue for model 7

- Molecule 1: Putative uncharacterized protein




4.2.8 Score per residue for model 8

- Molecule 1: Putative uncharacterized protein




4.2.9 Score per residue for model 9

- Molecule 1: Putative uncharacterized protein




4.2.10 Score per residue for model 10

- Molecule 1: Putative uncharacterized protein




4.2.11 Score per residue for model 11

- Molecule 1: Putative uncharacterized protein




4.2.12 Score per residue for model 12

- Molecule 1: Putative uncharacterized protein




4.2.13 Score per residue for model 13

- Molecule 1: Putative uncharacterized protein




4.2.14 Score per residue for model 14

- Molecule 1: Putative uncharacterized protein




4.2.15 Score per residue for model 15

- Molecule 1: Putative uncharacterized protein




4.2.16 Score per residue for model 16

- Molecule 1: Putative uncharacterized protein




4.2.17 Score per residue for model 17

- Molecule 1: Putative uncharacterized protein




4.2.18 Score per residue for model 18

- Molecule 1: Putative uncharacterized protein




4.2.19 Score per residue for model 19

- Molecule 1: Putative uncharacterized protein



4.2.20 Score per residue for model 20

- Molecule 1: Putative uncharacterized protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA 2.1	structure solution	
CNS 1.3	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2m3v_cs.str
Number of chemical shift lists	1
Total number of shifts	1919
Number of shifts mapped to atoms	1919
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	73%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.6±0.8
All	All	0	12

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	58	ARG	Sidechain	4
1	A	147	ARG	Sidechain	3
1	A	87	ARG	Sidechain	2
1	A	96	ARG	Sidechain	1
1	A	106	ARG	Sidechain	1
1	A	164	ARG	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1163	1165	1165	14±3
All	All	23260	23300	23300	282

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:52:MET:SD	1:A:56:LEU:HD11	0.71	2.26	14	5
1:A:39:TRP:CD1	1:A:160:GLU:HA	0.69	2.23	6	14
1:A:132:CYS:HA	1:A:139:THR:OG1	0.63	1.93	13	2
1:A:40:ALA:HA	1:A:51:ARG:O	0.60	1.96	9	15
1:A:78:VAL:O	1:A:97:VAL:HA	0.60	1.97	7	6
1:A:164:ARG:NE	1:A:164:ARG:HA	0.59	2.12	4	2
1:A:52:MET:SD	1:A:58:ARG:HG2	0.59	2.38	3	1
1:A:146:TYR:O	1:A:150:VAL:HG22	0.58	1.98	19	20
1:A:160:GLU:O	1:A:164:ARG:HB2	0.58	1.99	16	1
1:A:119:LEU:HB2	1:A:146:TYR:CE1	0.58	2.33	20	15
1:A:147:ARG:NH2	1:A:160:GLU:HB2	0.56	2.15	19	4
1:A:156:GLN:O	1:A:159:LEU:HB3	0.55	2.00	7	1
1:A:58:ARG:HA	1:A:130:MET:O	0.55	2.02	13	2
1:A:112:VAL:O	1:A:116:LEU:HG	0.54	2.03	2	10
1:A:66:SER:O	1:A:70:LEU:HG	0.53	2.03	10	17
1:A:109:ASP:O	1:A:112:VAL:HG12	0.53	2.04	15	3
1:A:138:ARG:NE	1:A:138:ARG:HA	0.53	2.19	14	1
1:A:123:GLU:HA	1:A:126:GLY:O	0.53	2.04	18	6
1:A:156:GLN:HE21	1:A:156:GLN:C	0.52	2.07	11	1
1:A:177:SER:O	1:A:181:ARG:HG2	0.52	2.05	11	10
1:A:86:ASP:HB2	1:A:90:LEU:HD12	0.52	1.81	7	3
1:A:142:PHE:O	1:A:146:TYR:HB3	0.52	2.04	5	2
1:A:109:ASP:O	1:A:113:LEU:HG	0.51	2.05	5	2
1:A:79:VAL:HG22	1:A:130:MET:HA	0.51	1.81	12	2
1:A:60:ALA:O	1:A:133:LYS:HA	0.51	2.06	12	2
1:A:48:ASN:O	1:A:59:SER:HA	0.50	2.06	20	10
1:A:176:ALA:O	1:A:180:VAL:HG23	0.50	2.07	10	7
1:A:68:ALA:O	1:A:72:ARG:HG2	0.50	2.07	5	8
1:A:146:TYR:CE1	1:A:150:VAL:HG21	0.50	2.42	20	5
1:A:185:VAL:O	1:A:189:ARG:HG2	0.49	2.07	1	3
1:A:79:VAL:HA	1:A:98:VAL:O	0.49	2.06	3	2
1:A:69:LEU:O	1:A:73:LEU:HG	0.49	2.07	20	5
1:A:141:LEU:HD23	1:A:176:ALA:HA	0.48	1.84	17	2
1:A:79:VAL:CG1	1:A:100:LEU:HB2	0.48	2.37	7	2
1:A:147:ARG:H	1:A:147:ARG:HD3	0.48	1.68	14	1
1:A:185:VAL:O	1:A:189:ARG:HD2	0.48	2.09	6	1
1:A:147:ARG:O	1:A:151:GLN:HB2	0.48	2.09	4	7
1:A:147:ARG:HG3	1:A:148:ILE:HD12	0.47	1.85	18	2
1:A:132:CYS:SG	1:A:139:THR:HB	0.47	2.49	14	1
1:A:39:TRP:NE1	1:A:160:GLU:HA	0.47	2.24	3	2
1:A:164:ARG:HA	1:A:164:ARG:NE	0.47	2.25	10	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:53:SER:OG	1:A:56:LEU:HG	0.47	2.08	15	1
1:A:67:VAL:HG12	1:A:90:LEU:HD23	0.47	1.87	6	9
1:A:86:ASP:HB3	1:A:97:VAL:HG22	0.47	1.86	15	2
1:A:106:ARG:O	1:A:106:ARG:HG2	0.47	2.09	8	1
1:A:79:VAL:CG2	1:A:130:MET:HA	0.47	2.40	12	3
1:A:44:ASP:OD2	1:A:47:ILE:HG12	0.47	2.10	7	2
1:A:67:VAL:O	1:A:71:GLN:HG3	0.46	2.11	3	2
1:A:186:ASP:HA	1:A:189:ARG:CD	0.46	2.40	8	2
1:A:159:LEU:HA	1:A:162:MET:HB3	0.45	1.89	20	1
1:A:115:VAL:O	1:A:119:LEU:HD22	0.45	2.12	10	7
1:A:79:VAL:CG2	1:A:131:HIS:H	0.45	2.24	17	1
1:A:113:LEU:O	1:A:117:ARG:HG3	0.45	2.12	13	3
1:A:138:ARG:HA	1:A:138:ARG:NE	0.45	2.26	13	1
1:A:51:ARG:HG3	1:A:51:ARG:O	0.44	2.12	17	1
1:A:147:ARG:HA	1:A:151:GLN:HG3	0.44	1.90	6	1
1:A:52:MET:HG3	1:A:53:SER:H	0.44	1.72	10	1
1:A:86:ASP:HA	1:A:89:TRP:NE1	0.44	2.28	9	1
1:A:113:LEU:O	1:A:117:ARG:HG2	0.44	2.12	17	1
1:A:111:GLU:O	1:A:115:VAL:HG12	0.44	2.12	9	1
1:A:47:ILE:HB	1:A:61:LEU:O	0.44	2.13	12	1
1:A:150:VAL:HG23	1:A:151:GLN:HG2	0.43	1.90	8	1
1:A:79:VAL:HG12	1:A:100:LEU:HG	0.43	1.90	11	1
1:A:114:SER:O	1:A:118:GLN:HB2	0.43	2.13	17	2
1:A:69:LEU:O	1:A:73:LEU:HD13	0.43	2.14	7	2
1:A:130:MET:HE1	1:A:143:ALA:HA	0.43	1.90	12	1
1:A:52:MET:HG2	1:A:160:GLU:OE1	0.43	2.14	16	1
1:A:100:LEU:HD11	1:A:115:VAL:HG13	0.42	1.91	8	1
1:A:136:ASN:ND2	1:A:165:GLY:HA3	0.42	2.29	13	1
1:A:111:GLU:O	1:A:114:SER:HB2	0.42	2.15	13	2
1:A:86:ASP:HB3	1:A:97:VAL:HB	0.42	1.92	1	1
1:A:59:SER:O	1:A:131:HIS:HB2	0.42	2.14	17	2
1:A:117:ARG:HA	1:A:120:GLN:CG	0.42	2.45	13	1
1:A:56:LEU:HD21	1:A:147:ARG:HD3	0.42	1.92	3	1
1:A:46:SER:O	1:A:61:LEU:HD12	0.42	2.15	14	1
1:A:161:GLU:HA	1:A:161:GLU:OE1	0.41	2.14	5	1
1:A:86:ASP:HB3	1:A:97:VAL:HG13	0.41	1.92	7	2
1:A:67:VAL:O	1:A:71:GLN:HG2	0.41	2.16	14	1
1:A:144:ALA:O	1:A:148:ILE:HD13	0.41	2.15	15	2
1:A:118:GLN:HE21	1:A:118:GLN:C	0.41	2.19	9	1
1:A:164:ARG:HA	1:A:164:ARG:HE	0.41	1.76	5	1
1:A:141:LEU:HA	1:A:167:PHE:CE1	0.41	2.50	8	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:108:ASP:CG	1:A:109:ASP:N	0.41	2.73	10	1
1:A:111:GLU:O	1:A:115:VAL:HG23	0.41	2.15	12	3
1:A:37:PRO:HB3	1:A:39:TRP:CE2	0.41	2.51	8	1
1:A:141:LEU:CD2	1:A:176:ALA:HA	0.40	2.46	1	1
1:A:161:GLU:OE1	1:A:164:ARG:HB2	0.40	2.16	5	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/193 (78%)	138±2 (92±2%)	10±2 (7±1%)	2±1 (1±1%)	17	62
All	All	3000/3860 (78%)	2757 (92%)	202 (7%)	41 (1%)	17	62

All 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	54	PRO	18
1	A	107	VAL	8
1	A	62	PRO	5
1	A	133	LYS	3
1	A	134	HIS	2
1	A	37	PRO	2
1	A	91	GLY	1
1	A	166	GLY	1
1	A	135	GLY	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/154 (78%)	113±1 (94±1%)	7±1 (6±1%)	25	73
All	All	2400/3080 (78%)	2256 (94%)	144 (6%)	25	73

All 34 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	76	LYS	20
1	A	119	LEU	15
1	A	156	GLN	15
1	A	118	GLN	15
1	A	179	TYR	10
1	A	146	TYR	7
1	A	53	SER	6
1	A	123	GLU	5
1	A	147	ARG	5
1	A	120	GLN	4
1	A	58	ARG	4
1	A	96	ARG	4
1	A	87	ARG	4
1	A	192	MET	3
1	A	86	ASP	3
1	A	161	GLU	3
1	A	145	MET	2
1	A	83	LYS	2
1	A	175	ASP	2
1	A	137	ASN	1
1	A	167	PHE	1
1	A	131	HIS	1
1	A	117	ARG	1
1	A	48	ASN	1
1	A	56	LEU	1
1	A	186	ASP	1
1	A	63	ASN	1
1	A	51	ARG	1
1	A	136	ASN	1
1	A	133	LYS	1
1	A	55	THR	1
1	A	155	LYS	1
1	A	189	ARG	1
1	A	159	LEU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 73% for the well-defined parts and 71% for the entire structure.

7.1 Chemical shift list 1

File name: 2m3v_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1919
Number of shifts mapped to atoms	1919
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	182	-0.19 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	173	0.08 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	159	0.27 ± 0.40	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 73%, i.e. 1343 atoms were assigned a chemical shift out of a possible 1839. 32 out of 32 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	542/736 (74%)	271/293 (92%)	144/300 (48%)	127/143 (89%)
Sidechain	756/987 (77%)	460/576 (80%)	288/354 (81%)	8/57 (14%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	45/116 (39%)	42/61 (69%)	0/49 (0%)	3/6 (50%)
Overall	1343/1839 (73%)	773/930 (83%)	432/703 (61%)	138/206 (67%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 71%, i.e. 1679 atoms were assigned a chemical shift out of a possible 2352. 35 out of 35 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	682/945 (72%)	341/376 (91%)	182/386 (47%)	159/183 (87%)
Sidechain	944/1256 (75%)	580/737 (79%)	355/445 (80%)	9/74 (12%)
Aromatic	53/151 (35%)	49/80 (61%)	0/62 (0%)	4/9 (44%)
Overall	1679/2352 (71%)	970/1193 (81%)	537/893 (60%)	172/266 (65%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	161	GLU	HA	1.70	6.30 – 2.20	-6.2

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

