



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 26, 2018 – 12:58 AM EDT

PDB ID : 1MC5  
Title : Ternary complex of Human glutathione-dependent formaldehyde dehydrogenase with S-(hydroxymethyl)glutathione and NADH  
Authors : Sanghani, P.C.; Bosron, W.F.; Hurley, T.D.  
Deposited on : 2002-08-05  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031021  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031021


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	<div> <div>2%</div> <div>79%</div> <div>21%</div> <div>.</div> </div>
1	B	374	<div> <div>79%</div> <div>20%</div> <div>.</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol dehydrogenase class III chi chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	373	Total	C	N	O	S	0	0	0
			2772	1765	465	520	22			
1	B	373	Total	C	N	O	S	0	0	0
			2772	1765	465	520	22			


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
5	B	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 6 is 2-AMINO-4-[1-CARBOXYMETHYL-CARBAMOYL)-2-HYDROXYMETHYLSULFANYL-ETHYLCARBAMOYL]-BUTYRIC ACID (three-letter code: AHE) (formula:  $C_{11}H_{19}N_3O_7S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			22	11	3	7	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	176	Total O 176 176	0	0
7	B	219	Total O 219 219	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alcohol dehydrogenase class III chi chain



- Molecule 1: Alcohol dehydrogenase class III chi chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.66 Å 78.66 Å 311.43 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.83 – 2.60 48.83 – 2.60	Depositor EDS
% Data completeness (in resolution range)	88.9 (48.83-2.60) 89.0 (48.83-2.60)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.09 (at 2.61 Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.197 , 0.249 0.197 , 0.250	Depositor DCC
$R_{free}$ test set	1402 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.9	Xtriage
Anisotropy	0.464	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6069	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, ZN, PO4, NAD, AHE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/2823	0.61	0/3817
1	B	0.35	0/2823	0.62	0/3817
All	All	0.35	0/5646	0.62	0/7634

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2772	0	2827	53	0
1	B	2772	0	2828	58	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
4	A	10	0	0	0	0
4	B	5	0	0	0	0
5	A	44	0	26	4	0
5	B	44	0	26	2	0
6	A	22	0	16	1	0
7	A	176	0	0	3	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	219	0	0	5	0
All	All	6069	0	5723	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:GLN:HE22	1:A:325:GLU:HG3	1.39	0.85
1:B:89:ILE:HD11	1:B:168:VAL:HG13	1.62	0.82
1:A:192:SER:HB3	7:A:614:HOH:O	1.82	0.80
5:A:500:NAD:C4N	6:A:404:AHE:HD21	2.12	0.80
1:A:39:ILE:HD12	1:A:165:LEU:HD12	1.67	0.75
1:A:195:ALA:HB2	1:A:261:VAL:HG11	1.69	0.74
1:B:95:GLN:HE22	1:B:325:GLU:HG3	1.55	0.69
1:B:111:GLN:HE22	1:B:114:ARG:HD2	1.58	0.68
1:B:29:PRO:HG3	1:B:35:ARG:HB2	1.77	0.67
1:A:95:GLN:HE22	1:A:325:GLU:CG	2.09	0.66
1:A:222:ASP:OD2	5:A:500:NAD:H1B	1.96	0.65
1:B:91:LEU:HD13	1:B:323:SER:HB2	1.80	0.64
1:B:284:GLY:HA2	1:B:310:GLY:HA3	1.81	0.62
1:A:275:ARG:O	1:A:279:GLU:HG3	2.00	0.61
1:B:89:ILE:HD11	1:B:168:VAL:CG1	2.30	0.61
1:B:253:LEU:HD23	1:B:256:MET:HE3	1.82	0.60
1:B:45:HIS:CE1	1:B:361:MET:HG2	2.36	0.60
1:B:208:MET:O	1:B:212:VAL:HG13	2.02	0.59
1:B:27:ALA:HB3	1:B:128:ARG:HG3	1.83	0.59
1:B:281:CYS:HB3	1:B:286:GLY:HA3	1.84	0.59
1:A:95:GLN:NE2	1:A:325:GLU:HG3	2.14	0.59
1:B:7:CYS:HB2	1:B:147:GLU:OE1	2.03	0.57
1:A:44:CYS:HB3	1:A:66:HIS:CE1	2.40	0.57
1:B:116:THR:HG23	7:B:672:HOH:O	2.04	0.57
1:A:94:PRO:HG3	7:A:548:HOH:O	2.05	0.56
1:B:178:GLY:HA3	1:B:205:ALA:HB3	1.88	0.56
1:B:44:CYS:HB2	1:B:67:GLU:OE2	2.06	0.55
1:A:270:ASN:O	1:A:274:MET:HG3	2.06	0.55
1:A:93:ILE:HD12	1:A:318:PHE:HB2	1.88	0.55
1:B:62:VAL:HG22	1:B:63:ILE:N	2.21	0.55
1:A:56:PRO:HG3	1:A:296:SER:HB3	1.87	0.55
1:A:250:GLN:HG2	1:A:279:GLU:HB2	1.89	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:PRO:HG3	1:A:35:ARG:HB2	1.89	0.54
1:A:115:VAL:HB	7:A:666:HOH:O	2.08	0.53
1:A:15:ALA:HB2	1:A:60:PHE:CZ	2.44	0.53
1:A:336:SER:O	1:A:337:LYS:HB2	2.09	0.53
1:B:291:VAL:O	5:B:501:NAD:H2N	2.09	0.52
1:B:271:VAL:HG11	1:B:299:GLU:O	2.09	0.52
1:B:336:SER:O	1:B:337:LYS:HB2	2.09	0.52
1:B:111:GLN:HE22	1:B:114:ARG:CD	2.23	0.52
1:A:92:TYR:CD2	1:A:93:ILE:HG13	2.45	0.51
1:A:30:LYS:HD2	1:A:128:ARG:NH2	2.26	0.51
1:B:347:HIS:HE1	1:B:365:LYS:O	1.94	0.50
1:A:278:LEU:HD22	1:A:313:TRP:CE3	2.46	0.50
1:A:12:ALA:HB2	1:A:19:LEU:HD22	1.93	0.49
1:A:350:SER:OG	1:A:353:GLU:HG2	2.13	0.49
1:B:222:ASP:HA	5:B:501:NAD:N3A	2.27	0.49
1:A:12:ALA:HB2	1:A:19:LEU:CD2	2.43	0.48
1:B:284:GLY:CA	1:B:310:GLY:HA3	2.44	0.48
1:B:188:LEU:HD11	1:B:213:ALA:HB3	1.95	0.48
1:A:327:VAL:O	1:A:331:VAL:HG13	2.14	0.47
1:A:266:GLU:HG3	1:A:274:MET:HG2	1.95	0.47
1:A:229:ALA:O	1:A:233:GLU:HG3	2.14	0.47
1:A:89:ILE:HG13	1:A:159:ILE:HD13	1.96	0.47
1:B:64:LEU:HA	1:B:139:TYR:CD2	2.50	0.47
1:B:250:GLN:HG2	1:B:279:GLU:HB2	1.97	0.47
1:B:30:LYS:O	1:B:76:GLY:HA3	2.15	0.47
1:A:281:CYS:HB3	1:A:286:GLY:HA3	1.97	0.46
1:B:128:ARG:HD3	7:B:605:HOH:O	2.16	0.46
1:B:278:LEU:HD22	1:B:313:TRP:CE3	2.50	0.46
1:A:62:VAL:HG22	1:A:63:ILE:N	2.30	0.46
1:B:168:VAL:HG22	1:B:171:LEU:HD12	1.96	0.46
1:A:129:PHE:HB2	1:A:136:ILE:O	2.15	0.46
1:A:283:LYS:HG2	1:A:284:GLY:N	2.31	0.46
1:A:93:ILE:HD12	1:A:318:PHE:CB	2.45	0.46
1:A:209:GLY:O	1:A:212:VAL:HG22	2.16	0.46
1:A:291:VAL:O	5:A:500:NAD:H2N	2.16	0.46
1:B:207:ILE:HD13	1:B:236:ALA:HB2	1.97	0.45
1:B:195:ALA:HB2	1:B:261:VAL:HG11	1.97	0.45
1:B:43:VAL:HG12	1:B:361:MET:HE2	1.97	0.45
1:A:37:LYS:O	1:A:70:GLY:HA3	2.17	0.45
1:A:222:ASP:HA	5:A:500:NAD:C2A	2.47	0.44
1:B:372:LYS:HE2	1:B:373:ILE:OXT	2.18	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:253:LEU:HA	1:B:256:MET:HE2	1.99	0.44
1:A:202:VAL:O	1:A:206:VAL:HG23	2.18	0.44
1:B:27:ALA:HB3	1:B:128:ARG:CG	2.48	0.44
1:B:361:MET:HE3	1:B:361:MET:HB2	1.74	0.43
1:B:95:GLN:HB3	1:B:324:VAL:HG23	2.00	0.43
1:A:211:LYS:HA	1:A:211:LYS:HD2	1.82	0.43
1:A:278:LEU:HD22	1:A:313:TRP:CD2	2.52	0.43
1:A:39:ILE:HD12	1:A:165:LEU:CD1	2.43	0.43
1:B:202:VAL:O	1:B:206:VAL:HG23	2.19	0.43
1:B:252:VAL:HG12	1:B:256:MET:HE1	2.00	0.43
1:B:27:ALA:HB3	1:B:128:ARG:CD	2.48	0.43
1:B:252:VAL:HG12	1:B:256:MET:CE	2.48	0.43
1:A:302:THR:O	1:B:299:GLU:HB2	2.19	0.43
1:A:56:PRO:HG3	1:A:296:SER:CB	2.49	0.43
1:B:62:VAL:CG2	1:B:63:ILE:N	2.81	0.43
1:B:45:HIS:HB3	7:B:697:HOH:O	2.19	0.43
1:A:241:ASN:HB3	1:A:244:ASP:OD2	2.18	0.42
1:B:283:LYS:HG2	1:B:284:GLY:N	2.34	0.42
1:B:306:GLN:O	1:B:311:ARG:HB2	2.19	0.42
1:B:8:LYS:HE2	1:B:21:ILE:CG2	2.49	0.42
1:B:91:LEU:HA	7:B:705:HOH:O	2.19	0.42
1:B:268:ILE:HD12	1:B:273:VAL:HG11	2.00	0.42
1:A:270:ASN:HB3	1:A:273:VAL:HB	2.01	0.42
1:B:209:GLY:O	1:B:212:VAL:HG22	2.20	0.42
1:A:333:GLU:CD	1:A:338:LYS:HD3	2.40	0.42
1:A:142:THR:O	1:A:143:SER:C	2.58	0.42
1:B:266:GLU:HG3	1:B:274:MET:HA	2.01	0.41
1:A:33:GLU:OE2	1:A:122:MET:HG2	2.21	0.41
1:B:55:ASP:HA	1:B:56:PRO:HD3	1.93	0.41
1:A:122:MET:HG3	1:A:123:PRO:HD2	2.03	0.41
1:A:188:LEU:HD11	1:A:213:ALA:HB3	2.02	0.40
1:A:250:GLN:CG	1:A:276:ALA:HA	2.52	0.40
1:B:327:VAL:O	1:B:331:VAL:HG23	2.21	0.40
1:B:44:CYS:SG	1:B:45:HIS:N	2.93	0.40
1:A:266:GLU:OE2	1:A:268:ILE:HB	2.21	0.40
1:A:283:LYS:HD2	1:B:101:PHE:CE2	2.56	0.40
1:B:55:ASP:HB3	7:B:567:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/374 (99%)	346 (93%)	23 (6%)	2 (0%)	31	56
1	B	371/374 (99%)	354 (95%)	17 (5%)	0	100	100
All	All	742/748 (99%)	700 (94%)	40 (5%)	2 (0%)	43	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173	CYS
1	A	174	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/300 (100%)	294 (98%)	5 (2%)	63	83
1	B	299/300 (100%)	295 (99%)	4 (1%)	71	88
All	All	598/600 (100%)	589 (98%)	9 (2%)	67	86

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	THR
1	A	116	THR
1	A	122	MET
1	A	307	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	313	TRP
1	B	34	VAL
1	B	128	ARG
1	B	188	LEU
1	B	313	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	GLN
1	A	184	ASN
1	A	243	GLN
1	B	111	GLN
1	B	184	ASN
1	B	243	GLN
1	B	347	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PO4	A	401	-	4,4,4	1.39	0	6,6,6	0.39	0
4	PO4	A	403	-	4,4,4	1.42	0	6,6,6	0.39	0
6	AHE	A	404	2	11,21,21	0.77	0	16,26,26	1.18	1 (6%)
5	NAD	A	500	-	40,48,48	2.14	10 (25%)	44,73,73	1.78	7 (15%)
4	PO4	B	402	-	4,4,4	1.41	0	6,6,6	0.38	0
5	NAD	B	501	-	40,48,48	2.15	10 (25%)	44,73,73	1.96	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PO4	A	401	-	-	0/0/0/0	0/0/0/0
4	PO4	A	403	-	-	0/0/0/0	0/0/0/0
6	AHE	A	404	2	-	0/19/26/26	0/0/0/0
5	NAD	A	500	-	-	0/22/62/62	0/5/5/5
4	PO4	B	402	-	-	0/0/0/0	0/0/0/0
5	NAD	B	501	-	-	0/22/62/62	0/5/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	500	NAD	C3N-C7N	-9.34	1.35	1.50
5	B	501	NAD	C3N-C7N	-8.95	1.36	1.50
5	B	501	NAD	C5N-C4N	-3.14	1.32	1.38
5	B	501	NAD	C4N-C3N	-2.76	1.34	1.39
5	A	500	NAD	C5N-C4N	-2.71	1.33	1.38
5	B	501	NAD	C2N-C3N	-2.26	1.35	1.39
5	A	500	NAD	C2N-C3N	-2.25	1.35	1.39
5	A	500	NAD	C4N-C3N	-2.14	1.35	1.39
5	A	500	NAD	C8A-N9A	2.11	1.39	1.36
5	B	501	NAD	C4A-N3A	2.44	1.39	1.35
5	A	500	NAD	C8A-N7A	2.46	1.39	1.34
5	A	500	NAD	C4A-N3A	2.57	1.39	1.35
5	B	501	NAD	C8A-N9A	2.60	1.40	1.36
5	B	501	NAD	O4D-C1D	2.79	1.45	1.41
5	A	500	NAD	O4D-C1D	2.93	1.45	1.41
5	B	501	NAD	C2A-N1A	3.24	1.40	1.33
5	B	501	NAD	C8A-N7A	3.24	1.40	1.34
5	A	500	NAD	C2A-N1A	3.70	1.40	1.33
5	B	501	NAD	C2A-N3A	4.74	1.39	1.32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	500	NAD	C2A-N3A	4.98	1.40	1.32

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	501	NAD	N3A-C2A-N1A	-9.50	120.73	128.86
5	A	500	NAD	N3A-C2A-N1A	-7.66	122.31	128.86
5	B	501	NAD	C4B-O4B-C1B	-3.64	106.03	109.83
5	A	500	NAD	C2B-C3B-C4B	-3.22	96.43	102.62
5	B	501	NAD	O7N-C7N-N7N	-2.93	118.33	122.60
5	A	500	NAD	O7N-C7N-N7N	-2.82	118.49	122.60
5	A	500	NAD	C4B-O4B-C1B	-2.79	106.92	109.83
5	A	500	NAD	C6N-C5N-C4N	-2.40	115.85	119.43
5	B	501	NAD	C2D-C3D-C4D	-2.32	98.17	102.62
5	B	501	NAD	O4B-C4B-C3B	2.28	109.66	105.15
5	B	501	NAD	O5B-C5B-C4B	2.31	117.02	109.00
6	A	404	AHE	CG1-CD1-N2	2.45	120.07	115.83
5	A	500	NAD	O5D-C5D-C4D	2.50	117.69	109.00
5	A	500	NAD	C3N-C7N-N7N	2.55	120.72	117.76
5	B	501	NAD	O5D-C5D-C4D	2.69	118.35	109.00
5	B	501	NAD	C3N-C7N-N7N	2.94	121.17	117.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	404	AHE	1	0
5	A	500	NAD	4	0
5	B	501	NAD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	373/374 (99%)	-0.16	9 (2%) 59 52	19, 36, 59, 69	0
1	B	373/374 (99%)	-0.40	1 (0%) 93 93	18, 31, 46, 63	0
All	All	746/748 (99%)	-0.28	10 (1%) 77 73	18, 33, 55, 69	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	12	ALA	3.2
1	A	15	ALA	3.2
1	B	1	ALA	3.2
1	A	13	TRP	2.7
1	A	133	GLY	2.6
1	A	131	CYS	2.4
1	A	246	SER	2.4
1	A	14	GLU	2.4
1	A	1	ALA	2.2
1	A	16	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PO4	A	403	5/5	0.85	0.33	80,81,81,81	5
4	PO4	B	402	5/5	0.90	0.29	81,81,82,82	5
5	NAD	A	500	44/44	0.92	0.19	35,44,52,54	0
6	AHE	A	404	22/22	0.93	0.21	54,57,63,64	0
4	PO4	A	401	5/5	0.95	0.18	74,74,75,75	0
5	NAD	B	501	44/44	0.96	0.14	34,39,41,42	0
2	ZN	B	376	1/1	0.98	0.07	59,59,59,59	0
2	ZN	A	375	1/1	0.98	0.04	38,38,38,38	0
3	K	A	400	1/1	0.98	0.07	38,38,38,38	0
2	ZN	A	376	1/1	0.99	0.07	41,41,41,41	0
2	ZN	B	375	1/1	0.99	0.06	29,29,29,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.