



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 16, 2020 – 01:18 PM EDT

PDB ID : 5MC4  
Title : Crystal Structure of Gly448Arg mutant of Human Prolidase with Mn ions and GlyPro ligand  
Authors : Wilk, P.; Mueller, U.; Dobbek, H.; Weiss, M.S.  
Deposited on : 2016-11-09  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.10.1  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.10.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	5253 (1.80-1.80)
Clashscore	122126	6077 (1.80-1.80)
Ramachandran outliers	120053	6011 (1.80-1.80)
Sidechain outliers	120020	6010 (1.80-1.80)
RSRZ outliers	108989	5157 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div> <div>6%</div> <div>93%</div> <div>6%</div> </div>
1	B	482	<div> <div>6%</div> <div>92%</div> <div>6%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xaa-Pro dipeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	0	18	0
			3725	2355	642	695	33			
1	B	473	Total	C	N	O	S	0	11	0
			3755	2375	656	694	30			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	448	ARG	GLY	engineered mutation	UNP P12955
B	448	ARG	GLY	engineered mutation	UNP P12955

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mn	0	0
			2	2		
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is HYDROXIDE ION (three-letter code: OH) (formula: HO).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O		0	0
			1	1			
3	B	1	Total	O		0	0
			1	1			

- Molecule 4 is GLYCINE (three-letter code: GLY) (formula: C<sub>2</sub>H<sub>5</sub>NO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			4	2	1	1		
4	B	1	Total	C	N	O	0	0
			4	2	1	1		

- Molecule 5 is PROLINE (three-letter code: PRO) (formula:  $C_5H_9NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	5	1	2		
5	B	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	330	Total	O	0	0
			330	330		
7	B	314	Total	O	0	2
			316	316		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xaa-Pro dipeptidase



- Molecule 1: Xaa-Pro dipeptidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.74Å 107.74Å 210.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.97 – 1.80 47.97 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.97-1.80) 99.9 (47.97-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.21 (at 1.79Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.163 , 0.192 0.173 , 0.198	Depositor DCC
$R_{free}$ test set	2101 reflections (1.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.7	Xtriage
Anisotropy	0.406	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 52.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.048 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8168	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN, OH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/3863	0.56	0/5220
1	B	0.39	0/3869	0.55	0/5229
All	All	0.39	0/7732	0.55	0/10449

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3725	0	3675	12	0
1	B	3755	0	3709	23	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	2	0	0
4	B	4	0	2	0	0
5	A	8	0	7	0	0
5	B	8	0	7	0	0
6	A	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	6	0	8	0	0
7	A	330	0	0	1	0
7	B	316	0	0	2	0
All	All	8168	0	7418	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:ILE:CG2	1:B:85[B]:VAL:HG13	2.23	0.69
1:B:210:MET:HE3	1:B:475:ILE:HG23	1.78	0.64
1:B:39:ALA:O	7:B:601:HOH:O	2.16	0.62
1:A:331[A]:ARG:NH2	7:A:602:HOH:O	2.33	0.60
1:A:416:TYR:CD2	1:A:448:ARG:HB2	2.37	0.60
1:B:159[B]:ARG:NH1	1:B:160:GLU:O	2.35	0.59
1:A:237:ARG:NH2	1:B:223:GLU:OE2	2.39	0.55
1:A:244:ILE:HB	1:A:276:ASP:HB3	1.90	0.54
1:B:85[A]:VAL:HG22	1:B:94:THR:HB	1.89	0.54
1:B:240:SER:HB3	1:B:376:VAL:CG2	2.39	0.53
1:A:110:LYS:HB3	1:B:254:LEU:HD21	1.91	0.53
1:B:159[B]:ARG:NH1	7:B:610:HOH:O	2.42	0.52
1:A:240:SER:HB3	1:A:376[B]:VAL:CG2	2.42	0.50
1:A:241:TYR:OH	1:A:276:ASP:OD1	2.30	0.49
1:B:201:ILE:HB	1:B:236:MET:HE1	1.95	0.48
1:A:246:GLY:N	1:A:274:LEU:O	2.47	0.47
1:A:262:PRO:HG3	1:B:237:ARG:HB2	1.96	0.47
1:A:198:THR:HA	1:A:236[A]:MET:HE1	1.98	0.45
1:B:210:MET:HB3	1:B:210:MET:HE2	1.90	0.44
1:B:244:ILE:HB	1:B:276:ASP:HB3	1.99	0.44
1:A:79:GLU:HG3	1:B:257:GLY:HA3	2.00	0.43
1:B:328:ASP:OD1	1:B:331[A]:ARG:NH1	2.52	0.42
1:B:45:ILE:CD1	1:B:141:PRO:HB3	2.50	0.42
1:A:108:MET:HB2	1:B:253:VAL:HG11	2.02	0.42
1:B:73:TRP:CD1	1:B:184:ARG:HG2	2.55	0.42
1:B:45:ILE:HG23	1:B:85[B]:VAL:HG13	2.02	0.41
1:B:96:PHE:HA	1:B:127:GLN:O	2.20	0.41
1:B:243:CYS:HB3	1:B:277:MET:HG2	2.03	0.40
1:B:45:ILE:HD12	1:B:141:PRO:HB3	2.03	0.40
1:B:146:THR:HB	1:B:160:GLU:HG3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/482 (99%)	469 (98%)	10 (2%)	0	100	100
1	B	478/482 (99%)	466 (98%)	12 (2%)	0	100	100
All	All	957/964 (99%)	935 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/404 (102%)	409 (100%)	2 (0%)	90	89
1	B	409/404 (101%)	408 (100%)	1 (0%)	94	93
All	All	820/808 (102%)	817 (100%)	3 (0%)	92	91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	MET
1	A	448	ARG
1	B	255	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 2 are modelled with single atom and 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	GOL	A	506	-	5,5,5	0.37	0	5,5,5	0.48	0
6	GOL	B	506	-	5,5,5	0.37	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	506	-	-	2/4/4/4	-
6	GOL	B	506	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	506	GOL	O1-C1-C2-C3
6	B	506	GOL	C1-C2-C3-O3
6	B	506	GOL	O1-C1-C2-O2
6	B	506	GOL	O2-C2-C3-O3
6	A	506	GOL	O1-C1-C2-O2
6	A	506	GOL	O1-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	465/482 (96%)	0.13	29 (6%) 20 16	20, 30, 64, 157	0
1	B	473/482 (98%)	0.13	29 (6%) 21 17	19, 30, 69, 163	0
All	All	938/964 (97%)	0.13	58 (6%) 20 16	19, 30, 69, 163	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	242	THR	15.0
1	B	243	CYS	12.1
1	B	111	ILE	7.5
1	A	107	TRP	7.5
1	A	104	HIS	7.1
1	A	101	PRO	6.8
1	A	109	GLY	6.7
1	A	246	GLY	6.2
1	B	245	CYS	6.2
1	B	102	ALA	6.0
1	B	244	ILE	5.9
1	B	105	ALA	5.8
1	B	103	SER	5.7
1	A	241	TYR	5.6
1	B	104	HIS	5.5
1	B	253	VAL	5.3
1	B	101	PRO	5.0
1	B	487	THR	4.9
1	A	111	ILE	4.9
1	A	242	THR	4.8
1	A	245	CYS	4.6
1	A	247	SER	4.6
1	A	105	ALA	4.2
1	A	243	CYS	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	241	TYR	3.9
1	A	164[A]	ASP	3.8
1	A	108	MET	3.7
1	B	260	GLY	3.7
1	A	261	ALA	3.5
1	B	259	ALA	3.5
1	B	252	ALA	3.2
1	B	159[A]	ARG	3.1
1	A	119	GLU	3.0
1	A	168	LYS	3.0
1	B	485	ALA	3.0
1	A	167	SER	2.9
1	B	256	TYR	2.9
1	A	244	ILE	2.9
1	B	110	LYS	2.6
1	B	427	ALA	2.6
1	A	102	ALA	2.5
1	B	112	HIS	2.5
1	B	441	GLN	2.4
1	A	239	SER	2.4
1	A	159	ARG	2.4
1	B	239	SER	2.3
1	B	255	HIS	2.3
1	A	100	LEU	2.3
1	A	166	ILE	2.3
1	A	110	LYS	2.3
1	A	6	GLY	2.2
1	A	115	GLU	2.2
1	B	486	PHE	2.2
1	B	254	LEU	2.1
1	B	356	GLN	2.1
1	A	106	THR	2.1
1	A	356	GLN	2.0
1	B	437	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	GOL	A	506	6/6	0.80	0.20	54,58,59,64	0
6	GOL	B	506	6/6	0.88	0.22	52,61,65,69	0
3	OH	A	501	1/1	0.88	0.15	35,35,35,35	0
4	GLY	B	503	4/5	0.90	0.23	37,42,45,46	0
4	GLY	A	503	4/5	0.92	0.19	34,35,38,40	0
5	PRO	B	504	8/8	0.92	0.14	45,52,55,56	0
5	PRO	A	504	8/8	0.93	0.11	42,43,50,52	0
3	OH	B	501	1/1	0.97	0.09	28,28,28,28	0
2	MN	B	500	1/1	0.98	0.07	32,32,32,32	1
2	MN	A	500	1/1	0.99	0.06	34,34,34,34	1
2	MN	A	499	1/1	1.00	0.07	24,24,24,24	1
2	MN	B	499	1/1	1.00	0.08	23,23,23,23	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.