



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 17, 2018 – 02:48 am GMT

PDB ID : 2MQP  
Title : Structural Investigation of hnRNP L bound to RNA  
Authors : Blatter, M.; Allain, F.  
Deposited on : 2014-06-24

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk30686
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30686


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 83%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136279	12091
Ramachandran outliers	132675	10835
Sidechain outliers	132484	10811
RNA backbone	3744	647

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	118	
2	B	6	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:190-A:283 (94)	0.11	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 7, 8, 11, 15, 18, 19
2	4, 5, 6, 14, 20
3	9, 13, 16
Single-model clusters	10; 12; 17

### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2013 atoms, of which 974 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Protein Hnrnpl.

Mol	Chain	Residues	Atoms						Trace
1	A	118	Total	C	H	N	O	S	0
			1824	572	908	159	180	5	

- Molecule 2 is a RNA chain called RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3').


Mol	Chain	Residues	Atoms						Trace
2	B	6	Total	C	H	N	O	P	0
			189	57	66	24	37	5	

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Protein Hnrnp1



- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.2 Score per residue for model 2

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.3 Score per residue for model 3

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.4 Score per residue for model 4

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.5 Score per residue for model 5

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.6 Score per residue for model 6

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.7 Score per residue for model 7

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.8 Score per residue for model 8

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.9 Score per residue for model 9

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.10 Score per residue for model 10

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')






### 4.2.11 Score per residue for model 11

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.12 Score per residue for model 12

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.13 Score per residue for model 13

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



#### 4.2.14 Score per residue for model 14

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



#### 4.2.15 Score per residue for model 15

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



#### 4.2.16 Score per residue for model 16

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.17 Score per residue for model 17

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')




### 4.2.18 Score per residue for model 18

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



### 4.2.19 Score per residue for model 19 (medoid)

- Molecule 1: Protein Hnrnp1




- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



4.2.20 Score per residue for model 20

- Molecule 1: Protein Hnrnp1



- Molecule 2: RNA (5'-R(\*AP\*CP\*AP\*CP\*AP\*C)-3')



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 250 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
AMBER	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2mqp_cs.str
Number of chemical shift lists	2
Total number of shifts	1425
Number of shifts mapped to atoms	1425
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	83%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.51±0.00	0±0/755 (0.0±0.0%)	0.75±0.01	0±0/1029 (0.0±0.0%)
2	B	1.04±0.01	0±0/137 (0.0±0.0%)	1.54±0.05	1±1/211 (0.5±0.6%)
All	All	0.62	0/17840 (0.0%)	0.93	19/24800 (0.1%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	6	C	O4'-C1'-N1	9.93	116.15	108.20	7	5
2	B	3	A	O4'-C1'-N9	6.17	113.14	108.20	11	8
2	B	3	A	C5'-C4'-C3'	-5.67	106.93	116.00	19	2
2	B	2	C	O4'-C1'-N1	5.55	112.64	108.20	19	2
2	B	6	C	C1'-O4'-C4'	-5.21	105.73	109.90	19	1
2	B	6	C	N1-C2-O2	5.11	121.97	118.90	7	1

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	740	742	742	2±1
2	B	123	66	68	0±1
All	All	17260	16160	16200	39

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:271:ASN:ND2	2:B:4:C:C2	0.55	2.74	11	5
1:A:213:ASN:N	1:A:214:PRO:HD2	0.45	2.27	20	20
1:A:266:LYS:N	1:A:267:PRO:HD2	0.43	2.27	6	7
1:A:273:PHE:CZ	2:B:5:A:C2	0.43	3.06	11	3
1:A:266:LYS:N	1:A:267:PRO:CD	0.41	2.84	13	4

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/118 (80%)	87±1 (92±1%)	6±1 (6±1%)	1±0 (1±0%)	20	67
All	All	1880/2360 (80%)	1739 (92%)	121 (6%)	20 (1%)	20	67

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	256	GLY	20

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/105 (80%)	77±1 (92±1%)	7±1 (8±1%)	17	64
All	All	1680/2100 (80%)	1544 (92%)	136 (8%)	17	64

All 12 unique residues with a non-rotameric sidechain are listed below. They are sorted by the

frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	200	ILE	20
1	A	259	THR	20
1	A	204	THR	20
1	A	195	THR	18
1	A	268	THR	17
1	A	279	THR	15
1	A	269	ARG	9
1	A	245	LYS	8
1	A	225	ARG	4
1	A	274	LYS	3
1	A	261	LYS	1
1	A	205	THR	1

### 6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	B	5/6 (83%)	4±0 (77±10%)	1±0 (18±6%)	0.01±0.01
All	All	100/120 (83%)	77 (77%)	18 (18%)	0.01

The overall RNA backbone suiteness is 0.01.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	2	C	20
2	B	4	C	20
2	B	6	C	19
2	B	3	A	18

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	5	A	18

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 83% for the well-defined parts and 82% for the entire structure.

### 7.1 Chemical shift list 1

File name: 2mqp\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1348
Number of shifts mapped to atoms	1348
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	116	$-0.41 \pm 0.16$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	111	$-0.04 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
$^{15}\text{N}$	111	$0.20 \pm 0.20$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 77%, i.e. 968 atoms were assigned a chemical shift out of a possible 1262. 15 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	368/462 (80%)	184/184 (100%)	94/188 (50%)	90/90 (100%)
Sidechain	533/598 (89%)	326/348 (94%)	191/222 (86%)	16/28 (57%)

*Continued on next page...*

Continued from previous page...

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	67/88 (76%)	44/46 (96%)	22/41 (54%)	1/1 (100%)
Overall	968/1262 (77%)	554/644 (86%)	307/493 (62%)	107/125 (86%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 77%, i.e. 1187 atoms were assigned a chemical shift out of a possible 1537. 17 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	454/578 (79%)	227/230 (99%)	116/236 (49%)	111/112 (99%)
Sidechain	666/757 (88%)	411/443 (93%)	233/274 (85%)	22/40 (55%)
Aromatic	67/88 (76%)	44/46 (96%)	22/41 (54%)	1/1 (100%)
Overall	1187/1537 (77%)	682/785 (87%)	371/593 (63%)	134/159 (84%)

#### 7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	275	ASN	HB3	-0.27	4.41 – 1.11	-9.2
1	A	220	ARG	CB	40.06	39.81 – 21.51	5.1

#### 7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



## 7.2 Chemical shift list 2

File name: 2mqp\_cs.str

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	77
Number of shifts mapped to atoms	77
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

### 7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 6%, i.e. 77 atoms were assigned a chemical shift out of a possible 1262. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	0/462 (0%)	0/184 (0%)	0/188 (0%)	0/90 (0%)
Sidechain	0/598 (0%)	0/348 (0%)	0/222 (0%)	0/28 (0%)
Aromatic	0/88 (0%)	0/46 (0%)	0/41 (0%)	0/1 (0%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Overall	77/1262 (6%)	41/644 (6%)	36/493 (7%)	0/125 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 5%, i.e. 77 atoms were assigned a chemical shift out of a possible 1537. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	0/578 (0%)	0/230 (0%)	0/236 (0%)	0/112 (0%)
Sidechain	0/757 (0%)	0/443 (0%)	0/274 (0%)	0/40 (0%)
Aromatic	0/88 (0%)	0/46 (0%)	0/41 (0%)	0/1 (0%)
Overall	77/1537 (5%)	41/785 (5%)	36/593 (6%)	0/159 (0%)

#### 7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.2.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned\_chem\_shift\_list\_2). RCI is only applicable to proteins.