



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2018 – 01:55 am GMT

PDB ID : 5MQL
Title : Crystal structure of dCK mutant C3S in complex with masitinib and UDP
Authors : Rebuffet, E.; Hammam, K.; Saez-Ayala, M.; Gros, L.; Lopez, S.; Hajem, B.;
Humbert, M.; Baudalet, E.; Audebert, S.; Betzi, S.; Lugari, A.; Combes, S.;
Pez, D.; Letard, S.; Mansfield, C.; Moussy, A.; de Sepulveda, P.; Morelli, X.;
Dubreuil, P.
Deposited on : 2016-12-20
Resolution : 3.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1008 (3.30-3.22)
Clashscore	122126	1066 (3.30-3.22)
Ramachandran outliers	120053	1046 (3.30-3.22)
Sidechain outliers	120020	1045 (3.30-3.22)
RSRZ outliers	108989	1993 (3.32-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 62%; width: 38%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 99%; width: 1%; height: 10px; background-color: grey;"></div> </div> <div>62% 17% 20%</div> </div>
1	B	289	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 58%; width: 42%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 99%; width: 1%; height: 10px; background-color: grey;"></div> </div> <div>3% 58% 20% 21%</div> </div>
1	C	289	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 56%; width: 44%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 99%; width: 1%; height: 10px; background-color: grey;"></div> </div> <div>% 56% 21% 21%</div> </div>
1	D	289	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 58%; width: 42%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 99%; width: 1%; height: 10px; background-color: grey;"></div> </div> <div>2% 58% 18% 22%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	G65	A	302	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxycytidine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1858	1192	305	353	8			
1	B	227	Total	C	N	O	S	0	0	0
			1806	1167	299	333	7			
1	C	228	Total	C	N	O	S	0	0	0
			1853	1190	308	348	7			
1	D	225	Total	C	N	O	S	0	0	0
			1815	1169	294	345	7			

There are 128 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-28	MET	-	initiating methionine	UNP P27707
A	-27	SER	-	expression tag	UNP P27707
A	-26	TYR	-	expression tag	UNP P27707
A	-25	TYR	-	expression tag	UNP P27707
A	-24	HIS	-	expression tag	UNP P27707
A	-23	HIS	-	expression tag	UNP P27707
A	-22	HIS	-	expression tag	UNP P27707
A	-21	HIS	-	expression tag	UNP P27707
A	-20	HIS	-	expression tag	UNP P27707
A	-19	HIS	-	expression tag	UNP P27707
A	-18	LEU	-	expression tag	UNP P27707
A	-17	GLU	-	expression tag	UNP P27707
A	-16	SER	-	expression tag	UNP P27707
A	-15	THR	-	expression tag	UNP P27707
A	-14	SER	-	expression tag	UNP P27707
A	-13	LEU	-	expression tag	UNP P27707
A	-12	TYR	-	expression tag	UNP P27707
A	-11	LYS	-	expression tag	UNP P27707
A	-10	LYS	-	expression tag	UNP P27707
A	-9	ALA	-	expression tag	UNP P27707
A	-8	GLY	-	expression tag	UNP P27707

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	LEU	-	expression tag	UNP P27707
A	-6	GLU	-	expression tag	UNP P27707
A	-5	ASN	-	expression tag	UNP P27707
A	-4	LEU	-	expression tag	UNP P27707
A	-3	TYR	-	expression tag	UNP P27707
A	-2	PHE	-	expression tag	UNP P27707
A	-1	GLN	-	expression tag	UNP P27707
A	0	GLY	-	expression tag	UNP P27707
A	9	SER	CYS	engineered mutation	UNP P27707
A	45	SER	CYS	engineered mutation	UNP P27707
A	59	SER	CYS	engineered mutation	UNP P27707
B	-28	MET	-	initiating methionine	UNP P27707
B	-27	SER	-	expression tag	UNP P27707
B	-26	TYR	-	expression tag	UNP P27707
B	-25	TYR	-	expression tag	UNP P27707
B	-24	HIS	-	expression tag	UNP P27707
B	-23	HIS	-	expression tag	UNP P27707
B	-22	HIS	-	expression tag	UNP P27707
B	-21	HIS	-	expression tag	UNP P27707
B	-20	HIS	-	expression tag	UNP P27707
B	-19	HIS	-	expression tag	UNP P27707
B	-18	LEU	-	expression tag	UNP P27707
B	-17	GLU	-	expression tag	UNP P27707
B	-16	SER	-	expression tag	UNP P27707
B	-15	THR	-	expression tag	UNP P27707
B	-14	SER	-	expression tag	UNP P27707
B	-13	LEU	-	expression tag	UNP P27707
B	-12	TYR	-	expression tag	UNP P27707
B	-11	LYS	-	expression tag	UNP P27707
B	-10	LYS	-	expression tag	UNP P27707
B	-9	ALA	-	expression tag	UNP P27707
B	-8	GLY	-	expression tag	UNP P27707
B	-7	LEU	-	expression tag	UNP P27707
B	-6	GLU	-	expression tag	UNP P27707
B	-5	ASN	-	expression tag	UNP P27707
B	-4	LEU	-	expression tag	UNP P27707
B	-3	TYR	-	expression tag	UNP P27707
B	-2	PHE	-	expression tag	UNP P27707
B	-1	GLN	-	expression tag	UNP P27707
B	0	GLY	-	expression tag	UNP P27707
B	9	SER	CYS	engineered mutation	UNP P27707
B	45	SER	CYS	engineered mutation	UNP P27707

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	59	SER	CYS	engineered mutation	UNP P27707
C	-28	MET	-	initiating methionine	UNP P27707
C	-27	SER	-	expression tag	UNP P27707
C	-26	TYR	-	expression tag	UNP P27707
C	-25	TYR	-	expression tag	UNP P27707
C	-24	HIS	-	expression tag	UNP P27707
C	-23	HIS	-	expression tag	UNP P27707
C	-22	HIS	-	expression tag	UNP P27707
C	-21	HIS	-	expression tag	UNP P27707
C	-20	HIS	-	expression tag	UNP P27707
C	-19	HIS	-	expression tag	UNP P27707
C	-18	LEU	-	expression tag	UNP P27707
C	-17	GLU	-	expression tag	UNP P27707
C	-16	SER	-	expression tag	UNP P27707
C	-15	THR	-	expression tag	UNP P27707
C	-14	SER	-	expression tag	UNP P27707
C	-13	LEU	-	expression tag	UNP P27707
C	-12	TYR	-	expression tag	UNP P27707
C	-11	LYS	-	expression tag	UNP P27707
C	-10	LYS	-	expression tag	UNP P27707
C	-9	ALA	-	expression tag	UNP P27707
C	-8	GLY	-	expression tag	UNP P27707
C	-7	LEU	-	expression tag	UNP P27707
C	-6	GLU	-	expression tag	UNP P27707
C	-5	ASN	-	expression tag	UNP P27707
C	-4	LEU	-	expression tag	UNP P27707
C	-3	TYR	-	expression tag	UNP P27707
C	-2	PHE	-	expression tag	UNP P27707
C	-1	GLN	-	expression tag	UNP P27707
C	0	GLY	-	expression tag	UNP P27707
C	9	SER	CYS	engineered mutation	UNP P27707
C	45	SER	CYS	engineered mutation	UNP P27707
C	59	SER	CYS	engineered mutation	UNP P27707
D	-28	MET	-	initiating methionine	UNP P27707
D	-27	SER	-	expression tag	UNP P27707
D	-26	TYR	-	expression tag	UNP P27707
D	-25	TYR	-	expression tag	UNP P27707
D	-24	HIS	-	expression tag	UNP P27707
D	-23	HIS	-	expression tag	UNP P27707
D	-22	HIS	-	expression tag	UNP P27707
D	-21	HIS	-	expression tag	UNP P27707
D	-20	HIS	-	expression tag	UNP P27707

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	HIS	-	expression tag	UNP P27707
D	-18	LEU	-	expression tag	UNP P27707
D	-17	GLU	-	expression tag	UNP P27707
D	-16	SER	-	expression tag	UNP P27707
D	-15	THR	-	expression tag	UNP P27707
D	-14	SER	-	expression tag	UNP P27707
D	-13	LEU	-	expression tag	UNP P27707
D	-12	TYR	-	expression tag	UNP P27707
D	-11	LYS	-	expression tag	UNP P27707
D	-10	LYS	-	expression tag	UNP P27707
D	-9	ALA	-	expression tag	UNP P27707
D	-8	GLY	-	expression tag	UNP P27707
D	-7	LEU	-	expression tag	UNP P27707
D	-6	GLU	-	expression tag	UNP P27707
D	-5	ASN	-	expression tag	UNP P27707
D	-4	LEU	-	expression tag	UNP P27707
D	-3	TYR	-	expression tag	UNP P27707
D	-2	PHE	-	expression tag	UNP P27707
D	-1	GLN	-	expression tag	UNP P27707
D	0	GLY	-	expression tag	UNP P27707
D	9	SER	CYS	engineered mutation	UNP P27707
D	45	SER	CYS	engineered mutation	UNP P27707
D	59	SER	CYS	engineered mutation	UNP P27707

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	C	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	D	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 3 is Masitinib (three-letter code: G65) (formula: $C_{28}H_{30}N_6OS$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			36	28	6	1	1		

- Molecule 4 is 2'-DEOXYCYTIDINE-5'-MONOPHOSPHATE (three-letter code: DCM) (formula: $C_9H_{14}N_3O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total 20	C 9	N 3	O 7	P 1	0	0
4	C	1	Total 20	C 9	N 3	O 7	P 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	5	Total O 5 5	0	0
6	B	3	Total O 3 3	0	0
6	C	5	Total O 5 5	0	0
6	D	5	Total O 5 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Deoxycytidine kinase




• Molecule 1: Deoxycytidine kinase




• Molecule 1: Deoxycytidine kinase





● Molecule 1: Deoxycytidine kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.33Å 88.33Å 342.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.80 – 3.25 47.95 – 3.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.80-3.25) 99.9 (47.95-3.25)	Depositor EDS
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 3.25Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.215 , 0.277 0.225 , 0.284	Depositor DCC
R_{free} test set	1120 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	100.5	Xtriage
Anisotropy	0.443	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 97.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7527	wwPDB-VP
Average B, all atoms (Å ²)	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DCM, UDP, G65

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/1904	0.74	0/2590
1	B	0.52	0/1852	0.72	0/2521
1	C	0.57	0/1899	0.73	0/2581
1	D	0.56	0/1861	0.73	0/2532
All	All	0.55	0/7516	0.73	0/10224

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1858	0	1734	39	0
1	B	1806	0	1690	61	0
1	C	1853	0	1750	58	0
1	D	1815	0	1688	44	0
2	A	25	0	11	0	0
2	B	25	0	11	0	0
2	C	25	0	11	1	0
2	D	25	0	11	1	0
3	A	36	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	20	0	12	1	0
4	C	20	0	12	1	0
5	C	1	0	0	0	0
6	A	5	0	0	0	0
6	B	3	0	0	0	0
6	C	5	0	0	0	0
6	D	5	0	0	0	0
All	All	7527	0	6930	192	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 192 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:ARG:NH1	1:B:233:ILE:HD12	1.32	1.38
1:B:219:ARG:HH11	1:B:233:ILE:CD1	1.40	1.34
1:B:219:ARG:NH1	1:B:233:ILE:CD1	1.91	1.30
1:B:47:ASP:OD1	1:B:119:ALA:HB1	1.36	1.25
1:C:41:LEU:HD11	1:C:48:TRP:CZ3	1.80	1.15

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	227/289 (78%)	202 (89%)	24 (11%)	1 (0%)	36 70
1	B	223/289 (77%)	200 (90%)	21 (9%)	2 (1%)	19 54
1	C	224/289 (78%)	209 (93%)	13 (6%)	2 (1%)	19 54
1	D	221/289 (76%)	188 (85%)	31 (14%)	2 (1%)	19 54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	895/1156 (77%)	799 (89%)	89 (10%)	7 (1%)	21 56

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	128	ARG
1	C	115	LYS
1	A	120	GLU
1	B	228	LEU
1	D	241	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	195/265 (74%)	186 (95%)	9 (5%)	29 62
1	B	185/265 (70%)	180 (97%)	5 (3%)	48 74
1	C	196/265 (74%)	186 (95%)	10 (5%)	26 59
1	D	189/265 (71%)	174 (92%)	15 (8%)	13 42
All	All	765/1060 (72%)	726 (95%)	39 (5%)	26 59

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	102	LEU
1	C	224	ASN
1	D	211	LYS
1	C	109	LEU
1	C	112	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	97	GLN
1	C	108	GLN
1	D	77	ASN
1	B	212	HIS
1	C	156	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UDP	A	301	-	20,26,26	1.04	2 (10%)	23,40,40	3.28	2 (8%)
3	G65	A	302	-	37,40,40	2.42	7 (18%)	49,55,55	1.65	9 (18%)
2	UDP	B	301	-	20,26,26	1.05	2 (10%)	23,40,40	3.26	2 (8%)
4	DCM	B	302	-	18,21,21	0.83	1 (5%)	24,31,31	0.60	0
2	UDP	C	302	-	20,26,26	0.97	2 (10%)	23,40,40	3.26	2 (8%)
4	DCM	C	303	5	18,21,21	0.91	1 (5%)	24,31,31	0.52	0
2	UDP	D	500	-	20,26,26	1.00	2 (10%)	23,40,40	3.27	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	301	-	-	0/12/32/32	0/2/2/2
3	G65	A	302	-	-	0/18/30/30	0/5/5/5
2	UDP	B	301	-	-	0/12/32/32	0/2/2/2
4	DCM	B	302	-	-	0/6/22/22	0/2/2/2
2	UDP	C	302	-	-	0/12/32/32	0/2/2/2
4	DCM	C	303	5	-	0/6/22/22	0/2/2/2
2	UDP	D	500	-	-	0/12/32/32	0/2/2/2

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	G65	CAF-CAG	-8.27	1.36	1.48
3	A	302	G65	CBC-CAZ	-6.31	1.40	1.51
3	A	302	G65	CAS-CAR	-6.04	1.39	1.51
3	A	302	G65	CAV-CAU	-5.15	1.39	1.50
3	A	302	G65	CAO-NAT	-3.68	1.34	1.41

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	G65	CAH-CAG-CAF	-4.25	123.53	129.44
2	C	302	UDP	C5-C4-N3	-3.66	114.66	123.17
2	B	301	UDP	C5-C4-N3	-3.62	114.77	123.17
2	D	500	UDP	C5-C4-N3	-3.60	114.81	123.17
2	A	301	UDP	C5-C4-N3	-3.59	114.82	123.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	G65	3	0
4	B	302	DCM	1	0
2	C	302	UDP	1	0
4	C	303	DCM	1	0
2	D	500	UDP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/289 (79%)	0.13	4 (1%) 70 62	75, 99, 128, 153	0
1	B	227/289 (78%)	0.41	8 (3%) 44 35	70, 96, 134, 145	0
1	C	228/289 (78%)	-0.03	2 (0%) 84 79	67, 88, 125, 170	0
1	D	225/289 (77%)	0.11	7 (3%) 49 40	72, 95, 127, 141	0
All	All	911/1156 (78%)	0.15	21 (2%) 60 53	67, 95, 129, 170	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	37	PHE	3.7
1	B	260	LEU	3.4
1	D	55	VAL	3.3
1	B	256	PHE	3.1
1	B	234	LEU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	G65	A	302	36/36	0.75	0.43	122,133,142,143	0
4	DCM	B	302	20/20	0.90	0.27	105,114,127,129	0
2	UDP	D	500	25/25	0.93	0.21	98,100,104,106	0
2	UDP	B	301	25/25	0.93	0.24	106,109,111,111	0
5	MG	C	301	1/1	0.94	0.24	67,67,67,67	0
2	UDP	A	301	25/25	0.94	0.26	113,119,124,125	0
2	UDP	C	302	25/25	0.94	0.23	91,95,105,107	0
4	DCM	C	303	20/20	0.95	0.27	80,89,108,109	0

6.5 Other polymers [i](#)

There are no such residues in this entry.