



Full wwPDB NMR Structure Validation Report ⓘ

Apr 28, 2019 – 07:17 AM EDT

PDB ID : 5MQX
Title : NMR solution structure of macro domain from Venezuelan equine encephalitis virus(VEEV) in complex with ADP-ribose
Authors : Makrynitsa, G.I.; Ntonti, D.; Marousis, K.D.; Matsoukas, M.T.; Papageorgiou, N.; Coutard, B.; Bentrop, D.; Spyroulias, G.A.
Deposited on : 2016-12-21

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.8.0 (224370), CSD as540be (2019)
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20031633
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20031633


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136327	12091
Ramachandran outliers	132723	10835
Sidechain outliers	132532	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	166	<div> <div style="width: 84%; background-color: green;"></div> <div style="width: 11%; background-color: yellow;"></div> <div style="width: 5%; background-color: cyan;"></div> <div style="width: 0%; background-color: grey;"></div> </div> <div>84% 11% . .</div>

2 Ensemble composition and analysis

This entry contains 21 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:160 (158)	0.52	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 16, 19, 21
2	11, 12, 13, 14, 15, 17, 18, 20

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2500 atoms, of which 1250 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called Non-structural protein3.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	160	2443	764	1229	214	233	3	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	161	HIS	-	expression tag	UNP P36328
A	162	HIS	-	expression tag	UNP P36328
A	163	HIS	-	expression tag	UNP P36328
A	164	HIS	-	expression tag	UNP P36328
A	165	HIS	-	expression tag	UNP P36328
A	166	HIS	-	expression tag	UNP P36328

- Molecule 2 is ADENOSINE-5-DIPHOSPHORIBOSE (three-letter code: APR) (formula: C₁₅H₂₃N₅O₁₄P₂).




4.2.3 Score per residue for model 3

- Molecule 1: Non-structural protein3




4.2.4 Score per residue for model 4

- Molecule 1: Non-structural protein3




4.2.5 Score per residue for model 5

- Molecule 1: Non-structural protein3




4.2.6 Score per residue for model 6

- Molecule 1: Non-structural protein3



4.2.7 Score per residue for model 7

- Molecule 1: Non-structural protein3





4.2.8 Score per residue for model 8

- Molecule 1: Non-structural protein3


Chain A: 77% 17% ..



4.2.9 Score per residue for model 9

- Molecule 1: Non-structural protein3

Chain A: 78% 16% ..



4.2.10 Score per residue for model 10

- Molecule 1: Non-structural protein3

Chain A: 80% 14% ..



4.2.11 Score per residue for model 11


- Molecule 1: Non-structural protein3

Chain A: 80% 14% ..




4.2.12 Score per residue for model 12

- Molecule 1: Non-structural protein3




4.2.13 Score per residue for model 13

- Molecule 1: Non-structural protein3




4.2.14 Score per residue for model 14

- Molecule 1: Non-structural protein3




4.2.15 Score per residue for model 15

- Molecule 1: Non-structural protein3



4.2.16 Score per residue for model 16


- Molecule 1: Non-structural protein3



HIS
HIS


4.2.17 Score per residue for model 17

- Molecule 1: Non-structural protein3

Chain A:  82% 12% ..


4.2.18 Score per residue for model 18

- Molecule 1: Non-structural protein3

Chain A:  84% 11% ..

4.2.19 Score per residue for model 19


- Molecule 1: Non-structural protein3

Chain A:  79% 14% ..

HIS


4.2.20 Score per residue for model 20

- Molecule 1: Non-structural protein3

Chain A:  80% 13% ..

4.2.21 Score per residue for model 21

- Molecule 1: Non-structural protein3



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 112 calculated structures, 21 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure calculation	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	5mqx_cs.cif
Number of chemical shift lists	1
Total number of shifts	1860
Number of shifts mapped to atoms	1860
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: APR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.66±0.01	0±0/1221 (0.0±0.0%)	1.17±0.10	5±2/1650 (0.3±0.1%)
All	All	0.66	0/25641 (0.0%)	1.17	108/34650 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.8±1.6
All	All	0	37

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	74	ASN	OD1-CG-ND2	-20.75	74.17	121.90	1	1
1	A	117	ASN	OD1-CG-ND2	-20.00	75.89	121.90	1	1
1	A	117	ASN	CB-CG-ND2	18.87	161.99	116.70	1	1
1	A	74	ASN	CB-CG-ND2	18.81	161.85	116.70	1	1
1	A	117	ASN	CB-CG-OD1	-17.75	86.10	121.60	1	1
1	A	74	ASN	CB-CG-OD1	-16.82	87.95	121.60	1	1
1	A	56	ARG	NE-CZ-NH2	14.04	127.32	120.30	2	16
1	A	158	ARG	NE-CZ-NH2	9.52	125.06	120.30	18	14
1	A	120	ARG	NE-CZ-NH2	9.46	125.03	120.30	8	13
1	A	159	ARG	NE-CZ-NH2	8.53	124.56	120.30	12	17
1	A	144	ARG	NE-CZ-NH2	8.46	124.53	120.30	17	12
1	A	56	ARG	CD-NE-CZ	7.39	133.95	123.60	2	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	142	TYR	CB-CG-CD1	-6.87	116.88	121.00	7	5
1	A	8	ARG	NE-CZ-NH1	-6.83	116.88	120.30	19	1
1	A	89	TYR	CB-CG-CD1	-6.49	117.10	121.00	5	2
1	A	4	TYR	CB-CG-CD1	-5.98	117.41	121.00	16	2
1	A	8	ARG	NE-CZ-NH2	5.89	123.25	120.30	19	9
1	A	56	ARG	NE-CZ-NH1	-5.82	117.39	120.30	2	2
1	A	8	ARG	CD-NE-CZ	5.77	131.67	123.60	7	2
1	A	144	ARG	NE-CZ-NH1	5.39	123.00	120.30	13	2
1	A	101	TYR	CB-CG-CD2	-5.37	117.78	121.00	3	1
1	A	38	TYR	CB-CG-CD2	-5.26	117.84	121.00	19	1
1	A	160	GLU	OE1-CD-OE2	-5.18	117.08	123.30	11	1
1	A	38	TYR	CB-CG-CD1	-5.01	117.99	121.00	5	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	89	TYR	Sidechain	6
1	A	101	TYR	Sidechain	4
1	A	159	ARG	Sidechain,Peptide	4
1	A	144	ARG	Sidechain	3
1	A	8	ARG	Sidechain	2
1	A	158	ARG	Sidechain	2
1	A	69	VAL	Peptide	2
1	A	17	GLY	Peptide	2
1	A	107	PRO	Peptide	1
1	A	60	GLY	Peptide	1
1	A	102	LYS	Peptide	1
1	A	73	PHE	Sidechain	1
1	A	4	TYR	Sidechain	1
1	A	110	SER	Peptide	1
1	A	56	ARG	Sidechain	1
1	A	114	PHE	Sidechain	1
1	A	119	ASP	Peptide	1
1	A	142	TYR	Sidechain	1
1	A	70	GLY	Peptide	1
1	A	38	TYR	Sidechain	1

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1202	1215	1215	1±1
2	A	36	21	21	0±0
All	All	25998	25956	25956	20

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:106:ILE:HD12	1:A:139:VAL:HG13	0.51	1.82	5	1
1:A:33:VAL:HG11	2:A:201:APR:O2A	0.48	2.08	6	1
1:A:23:ALA:HB2	1:A:67:HIS:CE1	0.48	2.43	3	1
1:A:158:ARG:HA	1:A:158:ARG:CZ	0.46	2.40	14	1
1:A:57:LEU:CD2	1:A:59:LYS:HE2	0.46	2.41	10	1
1:A:51:GLU:H	1:A:51:GLU:CD	0.46	2.14	1	1
1:A:51:GLU:CD	1:A:51:GLU:H	0.45	2.15	12	1
1:A:21:ASN:HD21	1:A:65:ILE:HG22	0.44	1.70	9	1
1:A:56:ARG:HG2	1:A:56:ARG:HH21	0.44	1.72	2	1
1:A:38:TYR:CE1	1:A:45:PHE:CZ	0.44	3.06	8	1
1:A:89:TYR:CD1	1:A:128:LEU:HD13	0.42	2.49	19	1
1:A:6:VAL:HG13	1:A:152:LEU:HD12	0.41	1.90	21	1
1:A:49:PRO:C	1:A:50:ILE:HD12	0.41	2.36	2	1
1:A:75:LYS:HE2	1:A:75:LYS:HA	0.41	1.90	6	1
1:A:19:ILE:HD13	1:A:41:PHE:CE2	0.41	2.50	7	1
1:A:11:ILE:CD1	1:A:19:ILE:HG23	0.41	2.46	21	2
1:A:121:LEU:HD11	1:A:148:TRP:CD1	0.40	2.50	1	1
1:A:11:ILE:HG12	1:A:142:TYR:CD1	0.40	2.52	12	1
1:A:99:ASN:ND2	1:A:101:TYR:CZ	0.40	2.90	16	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/166 (95%)	137±2 (88±1%)	17±2 (11±2%)	2±1 (2±1%)	15	59
All	All	3297/3486 (95%)	2885 (88%)	360 (11%)	52 (2%)	15	59

All 16 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	47	LEU	15
1	A	110	SER	11
1	A	111	THR	5
1	A	72	ASN	3
1	A	30	GLY	3
1	A	73	PHE	3
1	A	112	GLY	2
1	A	115	SER	2
1	A	114	PHE	1
1	A	45	PHE	1
1	A	107	PRO	1
1	A	117	ASN	1
1	A	62	ALA	1
1	A	46	ASP	1
1	A	146	LYS	1
1	A	29	PRO	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/134 (95%)	109±3 (86±2%)	18±3 (14±2%)	7	48
All	All	2667/2814 (95%)	2297 (86%)	370 (14%)	7	48

All 76 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	125	LEU	20
1	A	5	HIS	17
1	A	118	LYS	16
1	A	135	THR	15
1	A	87	GLU	14
1	A	40	LYS	13
1	A	119	ASP	12
1	A	99	ASN	12
1	A	83	LYS	12
1	A	56	ARG	10
1	A	75	LYS	10
1	A	109	LEU	9
1	A	59	LYS	8
1	A	82	ASP	8
1	A	74	ASN	8
1	A	54	LYS	7
1	A	134	THR	7
1	A	90	GLU	7
1	A	50	ILE	7
1	A	8	ARG	7
1	A	153	LYS	6
1	A	117	ASN	6
1	A	147	LYS	6
1	A	154	GLU	6
1	A	26	LYS	5
1	A	150	MET	5
1	A	146	LYS	5
1	A	19	ILE	5
1	A	102	LYS	5
1	A	144	ARG	4
1	A	136	ASP	4
1	A	113	ILE	4
1	A	77	SER	4
1	A	18	VAL	4
1	A	98	ASP	3
1	A	45	PHE	3
1	A	72	ASN	3
1	A	73	PHE	3
1	A	158	ARG	3
1	A	67	HIS	3
1	A	38	TYR	3
1	A	89	TYR	3
1	A	143	CYS	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	114	PHE	3
1	A	133	ASP	3
1	A	84	GLN	3
1	A	15	THR	3
1	A	46	ASP	3
1	A	104	VAL	3
1	A	11	ILE	3
1	A	28	GLN	2
1	A	94	LYS	2
1	A	51	GLU	2
1	A	3	SER	2
1	A	63	LYS	2
1	A	123	GLN	2
1	A	76	VAL	2
1	A	97	ASN	2
1	A	149	GLU	1
1	A	58	VAL	1
1	A	142	TYR	1
1	A	138	ASP	1
1	A	21	ASN	1
1	A	4	TYR	1
1	A	65	ILE	1
1	A	126	ASN	1
1	A	16	GLU	1
1	A	25	SER	1
1	A	39	LYS	1
1	A	145	ASP	1
1	A	156	VAL	1
1	A	111	THR	1
1	A	100	ASN	1
1	A	47	LEU	1
1	A	66	ILE	1
1	A	128	LEU	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	APR	A	201	-	33,39,39	1.07±0.05	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	APR	A	201	-	38,60,60	1.51±0.14	0±1 (1±1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	APR	A	201	-	-	0±0,18,54,54	0±0,4,4,4

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)	Models	
								Worst	Total
2	A	201	APR	O1D-C1D-O4D	7.22	101.86	111.15	9	6
2	A	201	APR	C4-C5-N7	5.29	114.92	109.40	12	2
2	A	201	APR	C4'-O4'-C1'	5.20	104.41	109.83	10	1

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 86% for the entire structure.

7.1 Chemical shift list 1

File name: 5mqx_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1860
Number of shifts mapped to atoms	1860
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	5

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	158	-0.06 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	145	0.13 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}'$	147	0.16 ± 0.18	None needed (< 0.5 ppm)
^{15}N	149	-0.16 ± 0.42	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 1614 atoms were assigned a chemical shift out of a possible 1876. 25 out of 25 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	753/780 (97%)	304/311 (98%)	301/316 (95%)	148/153 (97%)
Sidechain	819/980 (84%)	501/569 (88%)	315/366 (86%)	3/45 (7%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	42/116 (36%)	22/62 (35%)	19/49 (39%)	1/5 (20%)
Overall	1614/1876 (86%)	827/942 (88%)	635/731 (87%)	152/203 (75%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 86%, i.e. 1632 atoms were assigned a chemical shift out of a possible 1895. 25 out of 25 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	760/788 (96%)	306/314 (97%)	305/320 (95%)	149/154 (97%)
Sidechain	830/991 (84%)	508/576 (88%)	319/370 (86%)	3/45 (7%)
Aromatic	42/116 (36%)	22/62 (35%)	19/49 (39%)	1/5 (20%)
Overall	1632/1895 (86%)	836/952 (88%)	643/739 (87%)	153/204 (75%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	41	PHE	CE1	120.81	137.92 – 123.42	-6.8
1	A	72	ASN	HB2	0.89	4.36 – 1.26	-6.2
1	A	64	HIS	HE1	4.98	10.53 – 5.43	-5.9
1	A	72	ASN	HB3	0.89	4.41 – 1.11	-5.7
1	A	7	VAL	HB	0.37	3.59 – 0.39	-5.1

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

