



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 7, 2019 – 04:33 PM EST

PDB ID : 6NSF
Title : Crystal structure of the A/Brisbane/10/2007 (H3N2) influenza virus hemagglutinin G186V/L194P mutant in complex with 3'-SLNLN
Authors : Wu, N.C.; Wilson, I.A.
Deposited on : 2019-01-24
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.4
buster-report : 1.1.7 (2018)
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	<div> <div>13%</div> <div>90%</div> <div>9%</div> <div>.</div> </div>
2	B	176	<div> <div>%</div> <div>94%</div> <div>...</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BMA	A	412	-	-	-	X
7	SIA	A	416	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	7	0
			2518	1578	446	482	12			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	PRO	-	expression tag	UNP C3PR70
A	10	GLY	-	expression tag	UNP C3PR70
A	190	ASP	ASN	conflict	UNP C3PR70
A	194	PRO	LEU	engineered mutation	UNP C3PR70

- Molecule 2 is a protein called Hemagglutinin HA2 chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	173	Total	C	N	O	S	0	6	0
			1426	894	247	279	6			

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is BETA-L-FUCOSE (three-letter code: FUL) (formula: $C_6H_{12}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is O-SIALIC ACID (three-letter code: SIA) (formula: $C_{11}H_{19}NO_9$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			21	11	1	9		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	187	Total 187	O 187	0	0
8	B	181	Total 181	O 181	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hemagglutinin HA1 chain



• Molecule 2: Hemagglutinin HA2 chain



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	100.17Å 100.17Å 382.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10 43.10 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.10) 99.9 (43.10-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.182 , 0.214 0.189 , 0.220	Depositor DCC
R_{free} test set	2185 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4541	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, BMA, NAG, FUL, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/2588	0.72	1/3522 (0.0%)
2	B	0.54	0/1465	0.80	3/1969 (0.2%)
All	All	0.52	0/4053	0.75	4/5491 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	54	ARG	NE-CZ-NH1	9.55	125.07	120.30
2	B	54	ARG	NE-CZ-NH2	-9.16	115.72	120.30
2	B	145	ASP	CB-CG-OD1	5.80	123.52	118.30
1	A	299	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2518	0	2468	12	0
2	B	1426	0	1375	6	0
3	A	140	0	123	1	0
3	B	14	0	13	0	0
4	A	22	0	18	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	22	0	20	0	0
6	A	10	0	10	0	0
7	A	21	0	18	0	0
8	A	187	0	0	0	0
8	B	181	0	0	1	0
All	All	4541	0	4045	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:60[B]:ASN:H	2:B:60[B]:ASN:HD22	1.08	0.98
2:B:60[B]:ASN:H	2:B:60[B]:ASN:ND2	1.70	0.87
2:B:15:GLU:OE2	8:B:301:HOH:O	2.07	0.73
2:B:60[B]:ASN:HD22	2:B:60[B]:ASN:N	1.88	0.67
1:A:141:ARG:NH1	1:A:147:PHE:O	2.38	0.56
1:A:193:PHE:HB3	1:A:194:PRO:HD3	1.90	0.54
1:A:50[A]:GLU:HG2	1:A:273:PRO:HG2	1.91	0.51
1:A:188:ASP:HA	1:A:191:GLN:HB3	1.95	0.48
1:A:264:LYS:HD2	2:B:63:PHE:CE2	2.49	0.48
1:A:141:ARG:HB3	1:A:146:SER:CB	2.45	0.46
1:A:24:THR:OG1	3:A:401:NAG:H81	2.17	0.45
1:A:169:PRO:HA	1:A:242:ILE:HA	2.00	0.44
1:A:141:ARG:HB3	1:A:146:SER:HB2	2.01	0.43
1:A:166:VAL:O	1:A:244:LEU:HD12	2.20	0.41
1:A:264:LYS:HB3	2:B:63:PHE:CG	2.56	0.41
1:A:102:VAL:HG22	1:A:232:ILE:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/321 (100%)	308 (96%)	14 (4%)	0	100	100
2	B	177/176 (101%)	168 (95%)	9 (5%)	0	100	100
All	All	499/497 (100%)	476 (95%)	23 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/285 (101%)	276 (96%)	12 (4%)	32	32
2	B	152/148 (103%)	148 (97%)	4 (3%)	49	53
All	All	440/433 (102%)	424 (96%)	16 (4%)	41	39

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	128	THR
1	A	142	ARG
1	A	143	SER
1	A	167	THR
1	A	187	THR
1	A	236	ILE
1	A	246	ASN
1	A	276	LYS
1	A	289	PRO
1	A	296[A]	ASN
1	A	296[B]	ASN
2	B	38	LEU
2	B	60[A]	ASN
2	B	60[B]	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	168	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
1	A	246	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	401	1	14,14,15	0.58	0	17,19,21	1.18	2 (11%)
3	NAG	A	402	1,3	14,14,15	0.48	0	17,19,21	1.30	1 (5%)
3	NAG	A	403	3,4	14,14,15	0.32	0	17,19,21	0.82	1 (5%)
4	BMA	A	404	3,5	11,11,12	0.34	0	15,15,17	0.66	0
5	MAN	A	405	4	11,11,12	0.70	0	15,15,17	2.11	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MAN	A	406	4	11,11,12	0.40	0	15,15,17	0.89	1 (6%)
3	NAG	A	407	1,3,6	14,14,15	0.39	0	17,19,21	1.02	1 (5%)
6	FUL	A	408	3	9,10,11	0.49	0	13,14,16	1.08	2 (15%)
3	NAG	A	409	3	14,14,15	0.29	0	17,19,21	1.04	1 (5%)
3	NAG	A	410	1,3	14,14,15	0.42	0	17,19,21	1.06	1 (5%)
3	NAG	A	411	3,4	14,14,15	0.60	0	17,19,21	1.31	2 (11%)
4	BMA	A	412	3	11,11,12	0.75	0	15,15,17	2.09	2 (13%)
3	NAG	A	413	1,3	14,14,15	0.51	0	17,19,21	0.99	2 (11%)
3	NAG	A	414	3	14,14,15	0.27	0	17,19,21	0.55	0
3	NAG	A	415	1	14,14,15	0.43	0	17,19,21	0.86	0
7	SIA	A	416	-	18,21,21	1.09	1 (5%)	20,31,31	0.86	1 (5%)
3	NAG	B	201	2	14,14,15	0.32	0	17,19,21	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1	-	2/6/23/26	0/1/1/1
3	NAG	A	402	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	403	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	404	3,5	-	2/2/19/22	0/1/1/1
5	MAN	A	405	4	-	0/2/19/22	0/1/1/1
5	MAN	A	406	4	-	2/2/19/22	0/1/1/1
3	NAG	A	407	1,3,6	-	2/6/23/26	0/1/1/1
6	FUL	A	408	3	-	-	0/1/1/1
3	NAG	A	409	3	-	2/6/23/26	0/1/1/1
3	NAG	A	410	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	411	3,4	-	2/6/23/26	0/1/1/1
4	BMA	A	412	3	-	2/2/19/22	0/1/1/1
3	NAG	A	413	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	414	3	-	2/6/23/26	0/1/1/1
3	NAG	A	415	1	-	0/6/23/26	0/1/1/1
7	SIA	A	416	-	-	2/14/38/38	0/1/1/1
3	NAG	B	201	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	416	SIA	O2-C2	4.00	1.44	1.39

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	405	MAN	C1-C2-C3	6.06	117.11	109.66
4	A	412	BMA	C1-C2-C3	5.48	116.40	109.66
4	A	412	BMA	C1-O5-C5	4.77	118.69	112.20
5	A	405	MAN	C1-O5-C5	4.48	118.30	112.20
3	A	402	NAG	C1-O5-C5	4.42	118.21	112.20
3	A	410	NAG	C1-O5-C5	3.54	117.01	112.20
3	A	401	NAG	C1-O5-C5	2.89	116.13	112.20
6	A	408	FUL	C1-C2-C3	2.77	113.06	109.66
3	A	407	NAG	O5-C5-C6	2.70	111.42	107.15
3	A	411	NAG	C1-O5-C5	2.69	115.86	112.20
7	A	416	SIA	O6-C6-C5	2.55	112.25	109.74
3	A	409	NAG	C4-C3-C2	2.54	114.74	111.02
5	A	406	MAN	C1-O5-C5	2.42	115.50	112.20
3	A	403	NAG	C1-O5-C5	2.27	115.29	112.20
5	A	405	MAN	O5-C1-C2	2.23	114.23	110.79
3	A	401	NAG	O5-C5-C6	2.23	110.67	107.15
3	A	413	NAG	O5-C5-C4	-2.22	105.42	110.83
3	A	413	NAG	C1-O5-C5	-2.14	109.30	112.20
6	A	408	FUL	O5-C5-C4	-2.06	106.30	109.61
3	A	411	NAG	C3-C4-C5	-2.02	106.59	110.23

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	407	NAG	O5-C5-C6-O6
3	A	401	NAG	O5-C5-C6-O6
4	A	404	BMA	O5-C5-C6-O6
4	A	404	BMA	C4-C5-C6-O6
5	A	406	MAN	C4-C5-C6-O6
3	A	407	NAG	C4-C5-C6-O6
3	A	401	NAG	C4-C5-C6-O6
4	A	412	BMA	C4-C5-C6-O6
3	A	411	NAG	O5-C5-C6-O6
5	A	406	MAN	O5-C5-C6-O6
3	A	411	NAG	C4-C5-C6-O6
3	B	201	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	A	416	SIA	C7-C8-C9-O9
4	A	412	BMA	O5-C5-C6-O6
3	A	414	NAG	C4-C5-C6-O6
7	A	416	SIA	O8-C8-C9-O9
3	A	409	NAG	C4-C5-C6-O6
3	A	414	NAG	O5-C5-C6-O6
3	A	409	NAG	O5-C5-C6-O6
3	B	201	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/321 (98%)	0.81	42 (13%) 3 4	24, 46, 144, 182	0
2	B	173/176 (98%)	-0.31	2 (1%) 79 82	22, 32, 48, 91	0
All	All	490/497 (98%)	0.41	44 (8%) 9 12	22, 37, 119, 182	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	ALA	21.9
1	A	193	PHE	21.0
1	A	156	HIS	13.7
1	A	159	PHE	13.4
1	A	192	ILE	13.1
1	A	195	TYR	12.8
1	A	197	GLN	12.4
1	A	157	LEU	12.1
1	A	189	ASN	10.3
1	A	188	ASP	8.8
1	A	190	ASP	8.7
1	A	158	LYS	8.5
1	A	129	GLY	8.0
1	A	199	SER	8.0
1	A	194	PRO	7.9
1	A	161	TYR	7.8
1	A	198	ALA	7.4
1	A	191	GLN	7.4
1	A	162	PRO	7.4
1	A	186	VAL	6.9
1	A	187	THR	5.8
1	A	225	ASN	5.0
1	A	220	ARG	4.8
1	A	219	SER	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	128	THR	4.3
1	A	221	PRO	4.1
1	A	223	VAL	4.1
1	A	160	LYS	3.9
1	A	130	VAL	3.7
1	A	142	ARG	3.6
1	A	214	ILE	3.6
1	A	222	ARG	3.6
1	A	200	GLY	3.4
1	A	250	ASN	3.4
1	A	217	ILE	3.3
1	A	226	ILE	3.1
1	A	201	ARG	2.9
2	B	170	ARG	2.9
1	A	131	THR	2.8
1	A	218	GLY	2.7
1	A	224	ARG	2.6
2	B	2	ILE	2.3
1	A	246	ASN	2.1
1	A	164	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	A	412	11/12	0.23	0.63	136,140,142,142	0
3	NAG	A	401	14/15	0.62	0.40	65,72,76,77	0
3	NAG	A	410	14/15	0.64	0.37	93,101,104,110	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	B	201	14/15	0.64	0.33	72,81,82,85	0
7	SIA	A	416	21/21	0.65	0.43	96,104,106,106	0
4	BMA	A	404	11/12	0.71	0.25	92,98,101,105	0
5	MAN	A	406	11/12	0.74	0.35	103,107,114,114	0
3	NAG	A	409	14/15	0.77	0.39	88,99,103,105	0
5	MAN	A	405	11/12	0.78	0.27	96,101,102,102	0
3	NAG	A	411	14/15	0.80	0.46	116,119,122,128	0
6	FUL	A	408	10/11	0.80	0.38	92,95,97,98	0
3	NAG	A	407	14/15	0.80	0.28	58,66,85,86	0
3	NAG	A	413	14/15	0.81	0.18	42,50,54,64	0
3	NAG	A	415	14/15	0.82	0.20	77,87,90,92	0
3	NAG	A	402	14/15	0.82	0.16	47,51,53,60	0
3	NAG	A	414	14/15	0.85	0.40	72,78,81,84	0
3	NAG	A	403	14/15	0.85	0.23	65,71,74,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SIA A 416:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.