



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 10:45 pm GMT

PDB ID : 3NT3
Title : CRYSTAL STRUCTURE OF LSSmKate2 red fluorescent proteins with large Stokes shift
Authors : Malashkevich, V.N.; Piatkevich, K.; Almo, S.C.; Verkhusha, V.
Deposited on : 2010-07-02
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2534 (1.50-1.50)
Clashscore	122126	2727 (1.50-1.50)
Ramachandran outliers	120053	2661 (1.50-1.50)
Sidechain outliers	120020	2659 (1.50-1.50)
RSRZ outliers	108989	2481 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	
1	B	243	
1	C	243	
1	D	243	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	501	-	-	X	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8259 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LSSmKate2 red fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	10	0
			1836	1173	303	346	14			
1	B	224	Total	C	N	O	S	0	7	0
			1820	1160	302	345	13			
1	C	225	Total	C	N	O	S	0	6	0
			1819	1158	304	344	13			
1	D	224	Total	C	N	O	S	0	5	0
			1806	1150	301	342	13			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	251	Total 251	O 251	0	1
3	B	261	Total 261	O 261	0	1
3	C	223	Total 223	O 223	0	0
3	D	231	Total 231	O 231	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LSSmKate2 red fluorescent protein




- Molecule 1: LSSmKate2 red fluorescent protein



- Molecule 1: LSSmKate2 red fluorescent protein



- Molecule 1: LSSmKate2 red fluorescent protein



GLY
HIS
LYS
LEU
ASN

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.70Å 49.45Å 127.17Å 90.00° 93.13° 90.00°	Depositor
Resolution (Å)	19.99 – 1.50 19.99 – 1.50	Depositor EDS
% Data completeness (in resolution range)	96.0 (19.99-1.50) 96.0 (19.99-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 1.50Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.174 , 0.202 0.181 , 0.210	Depositor DCC
R_{free} test set	6886 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8259	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NRQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/1872	0.75	1/2525 (0.0%)
1	B	0.59	0/1846	0.75	1/2489 (0.0%)
1	C	0.59	0/1854	0.73	0/2500
1	D	0.60	0/1837	0.74	0/2477
All	All	0.60	0/7409	0.74	2/9991 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	222	CYS	C-N-CA	5.34	135.06	121.70
1	B	92	ARG	NE-CZ-NH2	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1836	0	1804	29	0
1	B	1820	0	1778	23	0
1	C	1819	0	1791	13	0
1	D	1806	0	1779	11	0
2	A	6	0	8	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	6	0	8	0	0
3	A	251	0	0	4	0
3	B	261	0	0	1	0
3	C	223	0	0	2	0
3	D	231	0	0	3	0
All	All	8259	0	7168	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159[B]:ASP:H	2:A:501:GOL:C1	1.85	0.90
1:A:159[A]:ASP:H	2:A:501:GOL:C1	1.85	0.90
1:A:159[B]:ASP:H	2:A:501:GOL:H11	1.44	0.81
1:A:159[A]:ASP:H	2:A:501:GOL:H11	1.44	0.81
1:A:105[A]:THR:HG23	3:A:850:HOH:O	1.81	0.80
1:A:11[B]:MET:CG	1:A:41[B]:MET:CE	2.62	0.77
1:C:18[A]:THR:CG2	1:C:21:ASN:HA	2.16	0.75
1:A:11[B]:MET:HG3	1:A:41[B]:MET:HE2	1.69	0.74
1:A:182:LYS:HD3	1:A:186:ASN:HB2	1.70	0.72
1:A:11[B]:MET:HG2	1:A:41[B]:MET:HE1	1.73	0.70
1:C:151:ASP:CG	1:C:151:ASP:O	2.30	0.68
1:A:127[A]:THR:HG23	3:A:858:HOH:O	1.93	0.67
1:B:11[B]:MET:SD	1:B:41[B]:MET:SD	2.92	0.67
1:D:18[A]:THR:OG1	1:D:122:ARG:NH2	2.28	0.67
1:A:11[B]:MET:CG	1:A:41[B]:MET:HE2	2.26	0.66
1:A:11[B]:MET:SD	1:A:41[B]:MET:SD	2.94	0.65
2:A:501:GOL:H2	1:B:158:SER:HA	1.79	0.65
1:B:11[B]:MET:CG	1:B:41[B]:MET:CE	2.76	0.64
1:A:11[B]:MET:HG3	1:A:41[B]:MET:CE	2.28	0.63
1:A:11[B]:MET:CG	1:A:41[B]:MET:HE1	2.27	0.62
1:D:18[A]:THR:HG21	3:D:293:HOH:O	1.99	0.62
1:B:187:LEU:HD22	3:B:538:HOH:O	2.00	0.62
1:B:25:LYS:HE3	1:B:47:GLU:OE2	2.01	0.60
1:A:63[B]:NRQ:HE2	1:A:199:LEU:HB2	1.85	0.59
1:B:185:LYS:HE2	1:B:185:LYS:H	1.68	0.59
1:B:11[B]:MET:HG2	1:B:41[B]:MET:HE1	1.85	0.58
1:B:11[B]:MET:HG3	1:B:41[B]:MET:HE2	1.86	0.57
1:C:41[A]:MET:SD	1:C:62:PHE:HB3	2.45	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:11[B]:MET:SD	1:C:41[B]:MET:HE2	2.45	0.57
1:C:18[A]:THR:HG21	1:C:21:ASN:HA	1.88	0.56
1:D:135:LYS:HE3	3:D:694:HOH:O	2.04	0.56
1:D:79:PHE:CE1	1:D:187:LEU:HD21	2.42	0.55
1:B:150:ALA:HB3	1:B:155:GLU:HG3	1.89	0.55
1:A:182:LYS:HD2	1:A:187:LEU:HD12	1.89	0.54
1:B:41[A]:MET:SD	1:B:62:PHE:HB3	2.47	0.54
1:C:2:SER:HB2	3:C:866:HOH:O	2.09	0.52
1:B:11[B]:MET:CG	1:B:41[B]:MET:HE2	2.39	0.52
1:C:63:NRQ:HD2	1:C:63:NRQ:O2	2.09	0.52
1:A:105[A]:THR:HG22	3:A:833:HOH:O	2.10	0.51
1:C:72:HIS:HA	1:C:219:ALA:HB3	1.93	0.51
1:B:151:ASP:OD2	1:B:151:ASP:O	2.29	0.50
1:D:78:ASP:O	1:D:82:GLN:HG3	2.11	0.50
1:B:63[B]:NRQ:HE2	1:B:199:LEU:HB2	1.93	0.50
1:A:151:ASP:OD1	1:A:151:ASP:O	2.30	0.50
1:A:18[A]:THR:CG2	1:A:21:ASN:HA	2.42	0.50
1:D:63:NRQ:O2	1:D:63:NRQ:HD2	2.13	0.48
1:B:11[B]:MET:HG2	1:B:41[B]:MET:CE	2.42	0.48
1:A:11[B]:MET:HG2	1:A:41[B]:MET:CE	2.37	0.46
1:A:67:TYR:OH	1:A:92:ARG:NH2	2.48	0.46
1:C:18[A]:THR:HG22	1:C:21:ASN:HA	1.97	0.46
1:C:11[A]:MET:HB3	1:C:115:LEU:HB2	1.98	0.46
1:A:63[B]:NRQ:HD2	1:A:215:GLU:OE2	2.16	0.45
1:A:11[B]:MET:SD	1:A:41[B]:MET:CE	3.05	0.45
1:D:41[A]:MET:SD	1:D:62:PHE:HB3	2.58	0.44
1:B:11[B]:MET:HG3	1:B:41[B]:MET:CE	2.47	0.43
1:B:182:LYS:HB3	1:B:187:LEU:HD21	2.00	0.43
1:D:72:HIS:HA	1:D:219:ALA:HB3	2.01	0.42
1:A:72:HIS:HA	1:A:219:ALA:HB3	2.01	0.42
1:B:188:LYS:NZ	1:B:189:VAL:O	2.53	0.42
1:D:41[A]:MET:CE	1:D:62:PHE:O	2.67	0.42
1:B:41[A]:MET:CE	1:B:62:PHE:O	2.68	0.42
1:B:11[B]:MET:CG	1:B:41[B]:MET:HE1	2.46	0.41
1:A:105[A]:THR:CG2	3:A:850:HOH:O	2.53	0.41
1:B:37:GLY:HA2	1:B:69:PHE:O	2.20	0.41
1:A:41[A]:MET:CE	1:A:62:PHE:O	2.69	0.41
1:D:85:PRO:O	3:D:283:HOH:O	2.22	0.41
1:B:185:LYS:CE	1:B:185:LYS:H	2.34	0.41
1:C:67:TYR:OH	1:C:92:ARG:NH2	2.53	0.41
1:A:159[A]:ASP:HB2	2:A:501:GOL:H12	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:LYS:HD2	1:C:122:ARG:NH2	2.36	0.41
1:A:41[A]:MET:SD	1:A:62:PHE:HB3	2.61	0.41
1:C:23:HIS:HB2	3:C:897:HOH:O	2.21	0.40
1:D:12:LYS:HB3	1:D:12:LYS:HE3	1.32	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/243 (94%)	224 (98%)	4 (2%)	0	100	100
1	B	225/243 (93%)	223 (99%)	2 (1%)	0	100	100
1	C	226/243 (93%)	222 (98%)	4 (2%)	0	100	100
1	D	224/243 (92%)	222 (99%)	2 (1%)	0	100	100
All	All	903/972 (93%)	891 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/207 (97%)	197 (98%)	3 (2%)	67	41
1	B	197/207 (95%)	194 (98%)	3 (2%)	67	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	198/207 (96%)	193 (98%)	5 (2%)	50	19
1	D	196/207 (95%)	191 (97%)	5 (3%)	49	17
All	All	791/828 (96%)	775 (98%)	16 (2%)	62	26

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	185	LYS
1	A	187	LEU
1	B	151	ASP
1	B	185	LYS
1	B	188	LYS
1	C	3	GLU
1	C	11[A]	MET
1	C	11[B]	MET
1	C	151	ASP
1	C	185	LYS
1	D	11[A]	MET
1	D	11[B]	MET
1	D	12	LYS
1	D	21	ASN
1	D	159	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	21	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	NRQ	A	63[A]	-	23,24,25	1.46	3 (13%)	25,32,34	4.58	9 (36%)
1	NRQ	A	63[B]	-	23,24,25	1.40	3 (13%)	25,32,34	3.57	6 (24%)
1	NRQ	B	63[A]	-	23,24,25	2.19	6 (26%)	25,32,34	3.80	7 (28%)
1	NRQ	B	63[B]	-	23,24,25	1.73	6 (26%)	25,32,34	3.83	10 (40%)
1	NRQ	C	63	1	23,24,25	1.54	3 (13%)	25,32,34	2.74	8 (32%)
1	NRQ	D	63	1	23,24,25	1.37	3 (13%)	25,32,34	2.28	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	NRQ	A	63[A]	-	-	0/9/31/32	0/2/2/2
1	NRQ	A	63[B]	-	-	0/9/31/32	0/2/2/2
1	NRQ	B	63[A]	-	-	0/9/31/32	0/2/2/2
1	NRQ	B	63[B]	-	-	0/9/31/32	0/2/2/2
1	NRQ	C	63	1	-	0/9/31/32	0/2/2/2
1	NRQ	D	63	1	-	0/9/31/32	0/2/2/2

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	63	NRQ	C2-N3	-4.18	1.29	1.39
1	C	63	NRQ	C1-N3	-3.96	1.31	1.38
1	B	63[B]	NRQ	C1-N3	-3.89	1.31	1.38
1	B	63[A]	NRQ	C1-N3	-3.89	1.31	1.38
1	A	63[B]	NRQ	C1-N3	-3.68	1.31	1.38
1	A	63[A]	NRQ	C1-N3	-3.68	1.31	1.38
1	A	63[B]	NRQ	C2-N3	-3.60	1.31	1.39
1	A	63[A]	NRQ	C2-N3	-3.60	1.31	1.39
1	D	63	NRQ	C1-N3	-3.57	1.32	1.38
1	B	63[B]	NRQ	C2-N3	-3.38	1.31	1.39
1	B	63[A]	NRQ	C2-N3	-3.38	1.31	1.39
1	D	63	NRQ	C2-N3	-2.98	1.32	1.39
1	B	63[B]	NRQ	CA2-C2	-2.75	1.45	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	63[A]	NRQ	CA2-C2	-2.75	1.45	1.48
1	B	63[B]	NRQ	CA3-N3	-2.26	1.42	1.47
1	B	63[A]	NRQ	CA3-N3	-2.26	1.42	1.47
1	A	63[B]	NRQ	CA2-C2	-2.13	1.46	1.48
1	A	63[A]	NRQ	CA2-C2	-2.13	1.46	1.48
1	D	63	NRQ	CA1-N1	2.35	1.33	1.27
1	B	63[B]	NRQ	CE1-CD1	2.47	1.43	1.38
1	B	63[B]	NRQ	CA1-N1	2.53	1.33	1.27
1	B	63[A]	NRQ	CA1-N1	2.53	1.33	1.27
1	C	63	NRQ	CA1-N1	2.65	1.34	1.27
1	B	63[A]	NRQ	CB2-CA2	7.31	1.41	1.35

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63[B]	NRQ	O2-C2-CA2	-10.06	125.63	130.97
1	A	63[A]	NRQ	O2-C2-CA2	-10.06	125.63	130.97
1	B	63[B]	NRQ	O2-C2-CA2	-7.98	126.74	130.97
1	B	63[A]	NRQ	O2-C2-CA2	-7.98	126.74	130.97
1	A	63[A]	NRQ	CB2-CA2-N2	-7.21	118.41	128.82
1	C	63	NRQ	O3-C3-CA3	-7.09	102.93	126.38
1	A	63[B]	NRQ	C2-CA2-N2	-6.38	104.42	108.93
1	A	63[A]	NRQ	C2-CA2-N2	-6.38	104.42	108.93
1	B	63[B]	NRQ	O3-C3-CA3	-6.25	105.71	126.38
1	B	63[A]	NRQ	O3-C3-CA3	-6.25	105.71	126.38
1	A	63[B]	NRQ	O3-C3-CA3	-6.07	106.30	126.38
1	A	63[A]	NRQ	O3-C3-CA3	-6.07	106.30	126.38
1	B	63[B]	NRQ	C2-CA2-N2	-5.70	104.91	108.93
1	B	63[A]	NRQ	C2-CA2-N2	-5.70	104.91	108.93
1	D	63	NRQ	CB2-CA2-N2	-5.67	120.64	128.82
1	B	63[A]	NRQ	CB2-CA2-N2	-5.23	121.27	128.82
1	B	63[B]	NRQ	CB2-CA2-C2	-4.64	116.61	122.28
1	C	63	NRQ	O2-C2-CA2	-4.62	128.52	130.97
1	B	63[B]	NRQ	CE1-CD1-CG2	-4.49	115.36	121.27
1	C	63	NRQ	C2-CA2-N2	-4.28	105.91	108.93
1	C	63	NRQ	CB2-CA2-N2	-3.53	123.72	128.82
1	D	63	NRQ	O3-C3-CA3	-2.58	117.84	126.38
1	D	63	NRQ	C2-CA2-N2	-2.55	107.13	108.93
1	B	63[B]	NRQ	CB1-CA1-N1	-2.47	117.03	125.38
1	B	63[A]	NRQ	CB1-CA1-N1	-2.47	117.03	125.38
1	A	63[B]	NRQ	CB1-CA1-N1	-2.12	118.20	125.38
1	A	63[A]	NRQ	CB1-CA1-N1	-2.12	118.20	125.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	63	NRQ	CB1-CA1-N1	-2.06	118.41	125.38
1	D	63	NRQ	CA2-C2-N3	2.13	104.22	103.30
1	A	63[A]	NRQ	CD1-CE1-CZ	2.17	122.30	119.88
1	A	63[B]	NRQ	CB2-CA2-C2	2.24	125.01	122.28
1	A	63[A]	NRQ	CE2-CD2-CG2	2.32	124.33	121.27
1	B	63[B]	NRQ	CE2-CD2-CG2	2.51	124.57	121.27
1	C	63	NRQ	CG2-CB2-CA2	2.54	133.00	130.10
1	C	63	NRQ	CA2-C2-N3	3.94	105.00	103.30
1	B	63[B]	NRQ	CD1-CE1-CZ	5.34	125.82	119.88
1	C	63	NRQ	CB2-CA2-C2	6.62	130.37	122.28
1	B	63[B]	NRQ	CB2-CA2-N2	6.68	138.47	128.82
1	D	63	NRQ	CB2-CA2-C2	8.15	132.24	122.28
1	B	63[A]	NRQ	CB2-CA2-C2	9.41	133.78	122.28
1	B	63[B]	NRQ	CA2-C2-N3	9.48	107.39	103.30
1	B	63[A]	NRQ	CA2-C2-N3	9.48	107.39	103.30
1	A	63[B]	NRQ	CA2-C2-N3	10.94	108.02	103.30
1	A	63[A]	NRQ	CA2-C2-N3	10.94	108.02	103.30
1	A	63[A]	NRQ	CB2-CA2-C2	12.18	137.16	122.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	63[A]	NRQ	1	0
1	A	63[B]	NRQ	2	0
1	B	63[A]	NRQ	1	0
1	B	63[B]	NRQ	1	0
1	C	63	NRQ	1	0
1	D	63	NRQ	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	501	-	5,5,5	0.72	0	5,5,5	1.31	1 (20%)
2	GOL	C	501	-	5,5,5	0.37	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	501	-	-	0/4/4/4	0/0/0/0
2	GOL	C	501	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	GOL	C3-C2-C1	-2.15	103.35	111.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	GOL	7	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/243 (91%)	0.12	10 (4%) 33 37	10, 17, 31, 55	0
1	B	223/243 (91%)	0.15	12 (5%) 26 28	11, 16, 32, 51	0
1	C	224/243 (92%)	0.13	13 (5%) 23 25	11, 17, 34, 46	0
1	D	223/243 (91%)	0.11	9 (4%) 38 42	12, 18, 32, 51	0
All	All	893/972 (91%)	0.13	44 (4%) 29 32	10, 17, 32, 55	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	186	ASN	10.6
1	D	184	ALA	10.3
1	A	184	ALA	8.4
1	B	186	ASN	8.3
1	B	187	LEU	7.4
1	D	186	ASN	6.1
1	C	184	ALA	6.1
1	B	189	VAL	5.3
1	A	187	LEU	5.0
1	A	183	PRO	4.9
1	C	186	ASN	4.8
1	A	152	GLY	4.5
1	D	187	LEU	4.4
1	B	185	LYS	4.3
1	A	185	LYS	4.1
1	C	152	GLY	4.0
1	B	151	ASP	3.8
1	A	151	ASP	3.8
1	C	3	GLU	3.8
1	A	189	VAL	3.7
1	B	188	LYS	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	166	GLY	3.6
1	C	185	LYS	3.4
1	B	73	THR	3.3
1	B	74	GLN	3.2
1	D	151	ASP	3.2
1	C	187	LEU	3.1
1	C	129	ASN	3.1
1	B	184	ALA	3.1
1	C	228	LEU	3.1
1	A	188	LYS	2.8
1	B	75	GLY	2.7
1	D	183	PRO	2.6
1	C	151	ASP	2.6
1	B	152	GLY	2.5
1	C	167	GLY	2.5
1	B	112	ASP	2.4
1	D	188	LYS	2.3
1	A	129	ASN	2.3
1	D	129	ASN	2.2
1	C	183	PRO	2.2
1	D	166	GLY	2.1
1	D	189	VAL	2.0
1	C	188	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	NRQ	B	63[B]	23/24	0.94	0.12	15,16,20,22	8
1	NRQ	A	63[B]	23/24	0.94	0.12	17,17,21,22	8
1	NRQ	B	63[A]	23/24	0.94	0.12	15,17,20,22	0
1	NRQ	A	63[A]	23/24	0.94	0.12	16,18,21,22	0
1	NRQ	D	63	23/24	0.95	0.08	16,17,21,23	0
1	NRQ	C	63	23/24	0.95	0.08	15,16,19,21	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	501	6/6	0.85	0.23	21,24,39,39	0
2	GOL	C	501	6/6	0.94	0.10	20,23,24,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.