



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 05:27 pm GMT

PDB ID : 2NXP  
Title : Structure of NTD2 domain of the human TAF5 subunit of TFIID  
Authors : Bhattacharya, S.; Takada, S.; Jacobson, R.H.  
Deposited on : 2006-11-17  
Resolution : 2.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	6027 (2.20-2.16)
Clashscore	122126	6837 (2.20-2.16)
Ramachandran outliers	120053	6731 (2.20-2.16)
Sidechain outliers	120020	6731 (2.20-2.16)
RSRZ outliers	108989	5899 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	156	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 88%, yellow 88%, yellow 95%, grey 95%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>88%</span> <span>7% • •</span> </div> </div>
1	B	156	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 12%, green 12%, green 90%, yellow 90%, yellow 97%, grey 97%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>12%</span> <span>90%</span> <span>• • 7%</span> </div> </div>
1	C	156	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 2%, green 2%, green 87%, yellow 87%, yellow 94%, grey 94%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>87%</span> <span>7% • 6%</span> </div> </div>
1	D	156	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 5%, green 5%, green 86%, yellow 86%, yellow 93%, grey 93%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>5%</span> <span>86%</span> <span>7% • 6%</span> </div> </div>
1	E	156	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 85%, yellow 85%, yellow 94%, grey 94%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>85%</span> <span>9% • 5%</span> </div> </div>
1	F	156	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 2%, green 2%, green 87%, yellow 87%, yellow 93%, grey 93%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>87%</span> <span>6% 6%</span> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	156	<div><div>4%</div><div><div></div></div><div>82%</div><div>12%</div><div>6%</div></div>
1	H	156	<div><div>4%</div><div><div></div></div><div>82%</div><div>11%</div><div>7%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10515 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription initiation factor TFIID subunit 5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	149	Total	C	N	O	S	Se	0	1	0
			1269	810	212	240	3	4			
1	B	145	Total	C	N	O	S	Se	0	0	0
			1235	790	207	231	3	4			
1	C	147	Total	C	N	O	S	Se	0	0	0
			1251	798	209	237	3	4			
1	D	147	Total	C	N	O	S	Se	0	0	0
			1251	798	209	237	3	4			
1	E	148	Total	C	N	O	S	Se	0	0	0
			1258	803	210	238	3	4			
1	F	146	Total	C	N	O	S	Se	0	0	0
			1243	794	208	234	3	4			
1	G	146	Total	C	N	O	S	Se	0	0	0
			1243	794	208	234	3	4			
1	H	145	Total	C	N	O	S	Se	0	0	0
			1236	789	207	233	3	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542
A	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
A	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
A	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542
B	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542
B	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
B	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
B	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542
C	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542
C	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
C	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
C	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542
D	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
D	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
D	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542
E	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542
E	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
E	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
E	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542
F	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542
F	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
F	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
F	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542
G	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542
G	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
G	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
G	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542
H	213	MSE	MET	MODIFIED RESIDUE	UNP Q15542
H	247	MSE	MET	MODIFIED RESIDUE	UNP Q15542
H	291	MSE	MET	MODIFIED RESIDUE	UNP Q15542
H	297	MSE	MET	MODIFIED RESIDUE	UNP Q15542

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0
2	E	1	Total Ca 1 1	0	0
2	H	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	F	1	Total Ca 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	67	Total 67	O 67	0	0
3	B	37	Total 37	O 37	0	0
3	C	102	Total 102	O 102	0	0
3	D	54	Total 54	O 54	0	0
3	E	95	Total 95	O 95	0	0
3	F	68	Total 68	O 68	0	0
3	G	45	Total 45	O 45	0	0
3	H	53	Total 53	O 53	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription initiation factor TFIID subunit 5




- Molecule 1: Transcription initiation factor TFIID subunit 5




- Molecule 1: Transcription initiation factor TFIID subunit 5




- Molecule 1: Transcription initiation factor TFIID subunit 5




- Molecule 1: Transcription initiation factor TFIID subunit 5




- Molecule 1: Transcription initiation factor TFIID subunit 5



• Molecule 1: Transcription initiation factor TFIID subunit 5



• Molecule 1: Transcription initiation factor TFIID subunit 5





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.62Å 61.83Å 133.36Å 90.00° 105.16° 90.00°	Depositor
Resolution (Å)	64.42 – 2.17 64.36 – 2.17	Depositor EDS
% Data completeness (in resolution range)	97.9 (64.42-2.17) 97.9 (64.36-2.17)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.213 , 0.263 0.221 , 0.267	Depositor DCC
$R_{free}$ test set	3837 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10515	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.9596e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.27	0/1300	0.42	0/1746
1	B	0.24	0/1262	0.37	0/1693
1	C	0.27	0/1278	0.41	0/1715
1	D	0.25	0/1278	0.39	0/1715
1	E	0.26	0/1286	0.41	0/1726
1	F	0.26	0/1270	0.40	0/1704
1	G	0.25	0/1270	0.38	0/1704
1	H	0.26	0/1263	0.40	0/1694
All	All	0.26	0/10207	0.40	0/13697

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1269	0	1201	8	0
1	B	1235	0	1176	5	0
1	C	1251	0	1184	7	0
1	D	1251	0	1184	9	0
1	E	1258	0	1192	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1243	0	1180	6	0
1	G	1243	0	1180	11	0
1	H	1236	0	1171	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	67	0	0	2	0
3	B	37	0	0	2	0
3	C	102	0	0	1	0
3	D	54	0	0	1	0
3	E	95	0	0	0	0
3	F	68	0	0	0	0
3	G	45	0	0	0	0
3	H	53	0	0	1	0
All	All	10515	0	9468	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:235:GLU:O	1:F:238:GLN:HG2	1.92	0.69
1:B:254:ASN:HB2	3:B:638:HOH:O	1.93	0.68
1:D:243:LEU:HD13	1:D:337:LEU:HD11	1.77	0.66
1:D:206:ASN:ND2	1:H:259:GLU:OE1	2.30	0.65
1:E:249:LEU:HD13	1:E:297:MSE:HE1	1.80	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	148/156 (95%)	147 (99%)	1 (1%)	0	100	100
1	B	143/156 (92%)	141 (99%)	2 (1%)	0	100	100
1	C	145/156 (93%)	143 (99%)	2 (1%)	0	100	100
1	D	145/156 (93%)	142 (98%)	3 (2%)	0	100	100
1	E	146/156 (94%)	145 (99%)	1 (1%)	0	100	100
1	F	144/156 (92%)	143 (99%)	1 (1%)	0	100	100
1	G	144/156 (92%)	141 (98%)	3 (2%)	0	100	100
1	H	143/156 (92%)	140 (98%)	3 (2%)	0	100	100
All	All	1158/1248 (93%)	1142 (99%)	16 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/142 (99%)	139 (99%)	2 (1%)	69	80
1	B	137/142 (96%)	136 (99%)	1 (1%)	85	92
1	C	139/142 (98%)	137 (99%)	2 (1%)	69	80
1	D	139/142 (98%)	136 (98%)	3 (2%)	55	66
1	E	140/142 (99%)	138 (99%)	2 (1%)	69	80
1	F	138/142 (97%)	138 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	138/142 (97%)	137 (99%)	1 (1%)	85	92
1	H	137/142 (96%)	136 (99%)	1 (1%)	85	92
All	All	1109/1136 (98%)	1097 (99%)	12 (1%)	76	85

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	207	GLN
1	D	243	LEU
1	E	339	ILE
1	C	340	ASP
1	E	201	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	258	ASN
1	E	315	GLN
1	H	223	HIS
1	C	322	GLN
1	D	331	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	145/156 (92%)	0.18	1 (0%) 87 87	21, 36, 59, 77	0
1	B	141/156 (90%)	0.81	19 (13%) 3 3	26, 45, 68, 78	0
1	C	143/156 (91%)	0.29	3 (2%) 63 63	18, 30, 58, 69	0
1	D	143/156 (91%)	0.49	8 (5%) 24 25	21, 40, 69, 94	0
1	E	144/156 (92%)	0.22	1 (0%) 87 87	19, 30, 53, 74	0
1	F	142/156 (91%)	0.22	3 (2%) 63 63	22, 33, 54, 72	0
1	G	142/156 (91%)	0.50	6 (4%) 36 37	23, 41, 64, 73	0
1	H	141/156 (90%)	0.30	7 (4%) 29 29	21, 37, 61, 71	0
All	All	1141/1248 (91%)	0.38	48 (4%) 36 37	18, 37, 63, 94	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	306	VAL	5.3
1	D	342	PHE	4.9
1	B	253	TYR	4.7
1	G	338	TYR	4.5
1	C	338	TYR	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	H	607	1/1	0.97	0.12	38,38,38,38	0
2	CA	G	606	1/1	0.98	0.06	33,33,33,33	0
2	CA	C	602	1/1	0.99	0.12	27,27,27,27	0
2	CA	F	605	1/1	0.99	0.11	30,30,30,30	0
2	CA	E	604	1/1	0.99	0.14	26,26,26,26	0
2	CA	A	600	1/1	0.99	0.10	38,38,38,38	0
2	CA	D	603	1/1	1.00	0.11	24,24,24,24	0
2	CA	B	601	1/1	1.00	0.08	32,32,32,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.