



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 09:33 pm GMT

PDB ID : 4NX2  
Title : Crystal structure of DCYRS complexed with DCY  
Authors : Wang, J.; Gong, W.; Li, J.; Gao, F.; Li, H.  
Deposited on : 2013-12-08  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	7193 (2.00-2.00)
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)
RSRZ outliers	108989	7011 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	<div> <div>3%</div> <div>87%</div> <div>11% ..</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	313	2482	1583	422	461	16	0	3	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	LEU	TYR	ENGINEERED MUTATION	UNP Q57834
A	65	ILE	LEU	ENGINEERED MUTATION	UNP Q57834
A	70	GLY	HIS	ENGINEERED MUTATION	UNP Q57834
A	108	ILE	PHE	ENGINEERED MUTATION	UNP Q57834
A	109	LEU	GLN	ENGINEERED MUTATION	UNP Q57834
A	114	GLY	TYR	ENGINEERED MUTATION	UNP Q57834
A	158	SER	ASP	ENGINEERED MUTATION	UNP Q57834
A	162	MET	LEU	ENGINEERED MUTATION	UNP Q57834
A	307	ALA	-	EXPRESSION TAG	UNP Q57834
A	308	HIS	-	EXPRESSION TAG	UNP Q57834
A	309	HIS	-	EXPRESSION TAG	UNP Q57834
A	310	HIS	-	EXPRESSION TAG	UNP Q57834
A	311	HIS	-	EXPRESSION TAG	UNP Q57834
A	312	HIS	-	EXPRESSION TAG	UNP Q57834
A	313	HIS	-	EXPRESSION TAG	UNP Q57834

- Molecule 2 is 3,5-dichloro-L-tyrosine (three-letter code: 2LT) (formula: C<sub>9</sub>H<sub>9</sub>Cl<sub>2</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			15	9	2	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	224	Total	O	0	0
			224	224		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-tRNA ligase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.10Å 160.96Å 39.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.01 – 2.00 39.01 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.5 (39.01-2.00) 92.3 (39.01-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.99 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, $R_{free}$	0.172 , 0.220 0.177 , 0.226	Depositor DCC
$R_{free}$ test set	1150 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.7	Xtriage
Anisotropy	0.397	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 52.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2721	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2LT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	2/2526 (0.1%)	0.61	3/3391 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	257	ARG	C-N	5.60	1.44	1.34
1	A	258	PRO	N-CD	5.42	1.55	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	258	PRO	CA-N-CD	-7.31	101.27	111.50
1	A	231	CYS	C-N-CD	5.84	140.66	128.40
1	A	256	LYS	C-N-CA	5.13	134.54	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	257	ARG	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2482	0	2553	57	0
2	A	15	0	7	2	0
3	A	224	0	0	1	0
All	All	2721	0	2560	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:PRO:HD2	1:A:259:GLU:CA	1.14	1.62
1:A:258:PRO:CG	1:A:259:GLU:HA	1.39	1.52
1:A:258:PRO:CD	1:A:259:GLU:CA	1.78	1.39
1:A:258:PRO:HD2	1:A:259:GLU:CB	1.57	1.33
1:A:258:PRO:CG	1:A:259:GLU:CA	2.03	1.30
1:A:258:PRO:CD	1:A:259:GLU:HA	0.83	1.30
1:A:258:PRO:HG2	1:A:259:GLU:C	1.49	1.28
1:A:258:PRO:HG2	1:A:259:GLU:CA	1.66	1.22
1:A:258:PRO:HG3	1:A:261:PHE:O	1.45	1.16
1:A:258:PRO:HD3	1:A:263:GLY:CA	1.79	1.11
1:A:258:PRO:CG	1:A:261:PHE:O	2.00	1.09
1:A:258:PRO:HD3	1:A:263:GLY:HA2	1.36	1.05
1:A:258:PRO:CD	1:A:263:GLY:HA2	1.87	1.05
1:A:261:PHE:O	1:A:263:GLY:N	1.96	0.98
1:A:112:LYS:HG3	1:A:113:ASP:N	1.79	0.98
1:A:258:PRO:HG2	1:A:260:LYS:N	1.80	0.96
1:A:113:ASP:HA	1:A:115:THR:H	1.32	0.95
1:A:258:PRO:HD3	1:A:263:GLY:N	1.81	0.94
1:A:112:LYS:O	1:A:115:THR:HG23	1.72	0.88
1:A:258:PRO:HG3	1:A:261:PHE:C	1.94	0.86
1:A:258:PRO:CG	1:A:259:GLU:C	2.35	0.85
1:A:112:LYS:HG3	1:A:113:ASP:H	1.43	0.84
1:A:258:PRO:HD3	1:A:263:GLY:H	1.41	0.84
1:A:113:ASP:HA	1:A:115:THR:N	1.94	0.82

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:PRO:HG3	1:A:261:PHE:CA	2.16	0.75
1:A:258:PRO:CD	1:A:261:PHE:O	2.35	0.74
1:A:258:PRO:HD2	1:A:259:GLU:HB3	1.69	0.72
1:A:109:LEU:O	1:A:115:THR:HG22	1.89	0.71
1:A:258:PRO:HD2	1:A:263:GLY:HA2	1.75	0.65
1:A:258:PRO:CG	1:A:261:PHE:N	2.61	0.64
1:A:259:GLU:HB3	1:A:263:GLY:HA2	1.82	0.61
1:A:258:PRO:HD3	1:A:261:PHE:O	2.04	0.58
2:A:401:2LT:OXT	3:A:711:HOH:O	2.17	0.58
1:A:258:PRO:HG3	1:A:261:PHE:N	2.19	0.57
1:A:258:PRO:HD2	1:A:259:GLU:CG	2.30	0.56
1:A:258:PRO:CG	1:A:259:GLU:N	2.64	0.54
1:A:258:PRO:HG3	1:A:261:PHE:CB	2.38	0.54
1:A:109:LEU:C	1:A:115:THR:HG22	2.29	0.53
1:A:259:GLU:HB3	1:A:263:GLY:CA	2.43	0.48
1:A:258:PRO:HG2	1:A:261:PHE:N	2.28	0.47
1:A:161:TYR:HD2	1:A:162[A]:MET:HE2	1.78	0.47
1:A:258:PRO:CD	1:A:259:GLU:CB	2.54	0.47
1:A:258:PRO:CG	1:A:261:PHE:H	2.28	0.46
1:A:69:LEU:HD22	1:A:106:SER:HA	1.99	0.45
1:A:258:PRO:CD	1:A:263:GLY:H	2.18	0.44
1:A:311:HIS:O	1:A:312:HIS:CB	2.65	0.43
1:A:150:ILE:HG22	1:A:154[B]:MET:SD	2.59	0.43
1:A:169:GLY:O	1:A:192:HIS:HA	2.19	0.42
1:A:259:GLU:CB	1:A:263:GLY:HA2	2.47	0.42
1:A:207:SER:HA	1:A:212:PHE:CD2	2.55	0.42
1:A:210:GLY:HA2	1:A:225:LYS:HD3	2.01	0.42
1:A:132:ARG:NH1	1:A:175:LYS:HE2	2.35	0.42
1:A:258:PRO:HG2	1:A:259:GLU:N	2.27	0.41
1:A:129:ARG:NE	1:A:132:ARG:HH21	2.17	0.41
1:A:34:GLY:HA3	2:A:401:2LT:HD2	2.02	0.41
1:A:261:PHE:C	1:A:263:GLY:H	2.09	0.41
1:A:231:CYS:HB3	1:A:288:LYS:HE3	2.02	0.40
1:A:251:TYR:HA	1:A:252:PRO:C	2.41	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/313 (100%)	304 (97%)	6 (2%)	4 (1%)	13	6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	258	PRO
1	A	262	GLY
1	A	312	HIS
1	A	257	ARG

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/272 (98%)	259 (97%)	8 (3%)	44	44

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LYS
1	A	32	LEU
1	A	69	LEU
1	A	139	ARG
1	A	254	THR
1	A	258	PRO
1	A	264	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	265	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	2LT	A	401	-	11,15,15	4.41	3 (27%)	16,21,21	2.10	9 (56%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2LT	A	401	-	-	0/4/8/8	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	2LT	CE2-CL2	-10.07	1.49	1.73
2	A	401	2LT	CE1-CL1	-9.94	1.49	1.73
2	A	401	2LT	CD1-CE1	2.30	1.42	1.38

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	2LT	CD1-CE1-CZ	-3.09	117.27	121.72
2	A	401	2LT	CE1-CD1-CG	-2.75	118.59	120.44
2	A	401	2LT	CD2-CE2-CZ	-2.49	118.13	121.72
2	A	401	2LT	OH-CZ-CE2	-2.20	114.58	119.78
2	A	401	2LT	CE2-CD2-CG	-2.14	119.00	120.44
2	A	401	2LT	CG-CB-CA	2.31	119.08	114.38
2	A	401	2LT	CD2-CG-CD1	2.57	122.63	118.97
2	A	401	2LT	CE2-CZ-CE1	2.77	124.35	117.66
2	A	401	2LT	CZ-CE1-CL1	3.32	123.28	118.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	2LT	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	313/313 (100%)	-0.24	10 (3%)	47 47	15, 23, 55, 81	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	PRO	5.7
1	A	262	GLY	3.8
1	A	261	PHE	3.7
1	A	312	HIS	3.5
1	A	263	GLY	3.4
1	A	139	ARG	3.0
1	A	260	LYS	2.9
1	A	313	HIS	2.6
1	A	109	LEU	2.4
1	A	140	GLU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	2LT	A	401	15/15	0.75	0.34	14,29,37,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.