



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 11, 2018 – 05:31 am GMT

PDB ID : 5NXU  
Title : Complex structure with maltose of Providencia stuartii Omp-Pst1 porin  
Authors : Colletier, J.P.; Nasrallah, C.  
Deposited on : 2017-05-10  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1851 (3.00-3.00)
Clashscore	122126	2167 (3.00-3.00)
Ramachandran outliers	120053	2101 (3.00-3.00)
Sidechain outliers	120020	2104 (3.00-3.00)
RSRZ outliers	108989	1751 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
1	B	352	<div> <div>3%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> </div>
1	C	352	<div> <div></div> <div> <div></div> <div>93%</div> <div>7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	LDA	A	401	-	-	-	X
2	LDA	B	406	-	-	-	X
2	LDA	B	407	-	-	-	X
2	LDA	B	409	-	-	-	X
2	LDA	C	405	-	-	-	X
2	LDA	C	406	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Porin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2758	1728	469	556	5			
1	B	352	Total	C	N	O	S	0	2	0
			2775	1738	472	559	6			
1	C	352	Total	C	N	O	S	0	1	0
			2767	1734	471	557	5			

- Molecule 2 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula:  $C_{14}H_{31}NO$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	B	1	Total	C	N	O	0	0
			16	14	1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			16	14	1	1		
2	B	1	Total	C	N	O	0	0
			11	9	1	1		
2	B	1	Total	C			0	0
			10	10				
2	B	1	Total	C	N	O	0	0
			15	13	1	1		
2	B	1	Total	C	N	O	0	0
			11	9	1	1		
2	B	1	Total	C	N	O	0	0
			11	9	1	1		
2	B	1	Total	C	N	O	0	0
			16	14	1	1		
2	B	1	Total	C	N	O	0	0
			16	14	1	1		
2	C	1	Total	C	N	O	0	0
			13	11	1	1		
2	C	1	Total	C	N	O	0	0
			16	14	1	1		
2	C	1	Total	C	N	O	0	0
			16	14	1	1		
2	C	1	Total	C	N	O	0	0
			16	14	1	1		
2	C	1	Total	C	N	O	0	0
			11	9	1	1		
2	C	1	Total	C	N	O	0	0
			16	14	1	1		
2	C	1	Total	C	N	O	0	0
			16	14	1	1		
2	C	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 3 is MALTOSE (three-letter code: MAL) (formula: C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			23	12	11		
3	B	1	Total	C	O	0	0
			23	12	11		
3	C	1	Total	C	O	0	0
			23	12	11		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	226	Total	O	0	0
			226	226		
5	B	166	Total	O	0	0
			166	166		
5	C	145	Total	O	0	0
			145	145		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Porin 1



#### • Molecule 1: Porin 1



#### • Molecule 1: Porin 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.20Å 127.12Å 150.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.17 – 3.00 39.41 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (97.17-3.00) 99.6 (39.41-3.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.211 , 0.255 0.210 , 0.256	Depositor DCC
$R_{free}$ test set	2030 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	82.3	Xtriage
Anisotropy	0.319	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 56.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9181	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	98.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MAL, LDA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/2815	0.79	4/3800 (0.1%)
1	B	0.40	0/2832	0.73	4/3822 (0.1%)
1	C	0.39	0/2824	0.68	4/3811 (0.1%)
All	All	0.40	0/8471	0.74	12/11433 (0.1%)

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	20	ARG	NE-CZ-NH1	18.43	129.51	120.30
1	A	20	ARG	NE-CZ-NH2	-17.84	111.38	120.30
1	B	39	ARG	NE-CZ-NH2	13.21	126.90	120.30
1	B	20	ARG	NE-CZ-NH1	-10.20	115.20	120.30
1	B	39	ARG	NE-CZ-NH1	-10.12	115.24	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2758	0	2591	7	0
1	B	2775	0	2606	11	0
1	C	2767	0	2603	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	32	0	62	0	0
2	B	122	0	223	4	0
2	C	120	0	226	3	0
3	A	23	0	22	0	0
3	B	23	0	22	2	0
3	C	23	0	22	0	0
4	B	1	0	0	0	0
5	A	226	0	0	1	0
5	B	166	0	0	1	0
5	C	145	0	0	1	0
All	All	9181	0	8377	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 23 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:344:LEU:HD22	2:B:402:LDA:H51	1.65	0.79
1:B:192:GLY:HA3	2:B:405:LDA:H101	1.66	0.77
1:B:199:ARG:NH1	5:B:501:HOH:O	2.31	0.63
1:B:4:TYR:HB3	1:B:11:LEU:HB2	1.87	0.56
1:B:154:TYR:CE2	2:B:401:LDA:H32	2.40	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/352 (99%)	321 (92%)	26 (7%)	3 (1%)	19	59
1	B	352/352 (100%)	321 (91%)	28 (8%)	3 (1%)	19	59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	351/352 (100%)	323 (92%)	24 (7%)	4 (1%)	16	53
All	All	1053/1056 (100%)	965 (92%)	78 (7%)	10 (1%)	17	59

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	7	ASP
1	B	7	ASP
1	C	7	ASP
1	A	186	TRP
1	C	88	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/284 (100%)	270 (95%)	14 (5%)	27	66
1	B	286/284 (101%)	276 (96%)	10 (4%)	39	75
1	C	285/284 (100%)	275 (96%)	10 (4%)	39	75
All	All	855/852 (100%)	821 (96%)	34 (4%)	34	72

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	30	LYS
1	B	131	ARG
1	C	242	MET
1	B	88	ASP
1	A	88	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	235	ASN
1	C	202	ASN
1	A	270	GLN
1	A	155	GLN
1	B	297	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 1 is monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LDA	A	401	-	12,15,15	0.75	1 (8%)	14,17,17	0.45	0
2	LDA	A	402	-	12,15,15	0.71	1 (8%)	14,17,17	0.64	0
3	MAL	A	403	-	24,24,24	0.67	0	35,35,35	1.32	6 (17%)
2	LDA	B	401	-	12,15,15	1.02	1 (8%)	14,17,17	3.31	4 (28%)
2	LDA	B	402	-	12,15,15	0.75	1 (8%)	14,17,17	0.62	0
2	LDA	B	403	-	7,10,15	0.91	1 (14%)	9,12,17	0.41	0
2	LDA	B	404	-	9,9,15	0.41	0	8,8,17	0.43	0
2	LDA	B	405	-	11,14,15	1.07	1 (9%)	13,16,17	3.27	4 (30%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LDA	B	406	-	7,10,15	1.29	1 (14%)	9,12,17	3.96	4 (44%)
2	LDA	B	407	-	7,10,15	1.31	1 (14%)	9,12,17	3.82	4 (44%)
2	LDA	B	408	-	12,15,15	1.01	1 (8%)	14,17,17	3.31	4 (28%)
2	LDA	B	409	-	12,15,15	0.71	0	14,17,17	0.61	0
3	MAL	B	410	-	24,24,24	0.56	0	35,35,35	1.09	2 (5%)
2	LDA	C	401	-	9,12,15	1.20	1 (11%)	11,14,17	3.59	4 (36%)
2	LDA	C	402	-	12,15,15	0.73	0	14,17,17	0.60	0
2	LDA	C	403	-	12,15,15	0.76	1 (8%)	14,17,17	0.87	1 (7%)
2	LDA	C	404	-	12,15,15	1.06	1 (8%)	14,17,17	3.18	4 (28%)
2	LDA	C	405	-	7,10,15	1.47	1 (14%)	9,12,17	3.89	4 (44%)
2	LDA	C	406	-	12,15,15	0.83	1 (8%)	14,17,17	0.70	0
2	LDA	C	407	-	12,15,15	0.70	0	14,17,17	0.60	0
2	LDA	C	408	-	12,15,15	0.71	1 (8%)	14,17,17	0.45	0
3	MAL	C	409	-	24,24,24	0.56	0	35,35,35	0.96	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LDA	A	401	-	-	0/13/13/13	0/0/0/0
2	LDA	A	402	-	-	0/13/13/13	0/0/0/0
3	MAL	A	403	-	-	0/8/48/48	0/2/2/2
2	LDA	B	401	-	-	0/13/13/13	0/0/0/0
2	LDA	B	402	-	-	0/13/13/13	0/0/0/0
2	LDA	B	403	-	-	0/8/8/13	0/0/0/0
2	LDA	B	404	-	-	0/7/7/13	0/0/0/0
2	LDA	B	405	-	-	0/12/12/13	0/0/0/0
2	LDA	B	406	-	-	0/8/8/13	0/0/0/0
2	LDA	B	407	-	-	0/8/8/13	0/0/0/0
2	LDA	B	408	-	-	0/13/13/13	0/0/0/0
2	LDA	B	409	-	-	0/13/13/13	0/0/0/0
3	MAL	B	410	-	-	0/8/48/48	0/2/2/2
2	LDA	C	401	-	-	0/10/10/13	0/0/0/0
2	LDA	C	402	-	-	0/13/13/13	0/0/0/0
2	LDA	C	403	-	-	0/13/13/13	0/0/0/0
2	LDA	C	404	-	-	0/13/13/13	0/0/0/0
2	LDA	C	405	-	-	0/8/8/13	0/0/0/0
2	LDA	C	406	-	-	0/13/13/13	0/0/0/0

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LDA	C	407	-	-	0/13/13/13	0/0/0/0
2	LDA	C	408	-	-	0/13/13/13	0/0/0/0
3	MAL	C	409	-	-	0/8/48/48	0/2/2/2

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	408	LDA	O1-N1	2.03	1.47	1.42
2	B	403	LDA	O1-N1	2.07	1.47	1.42
2	C	403	LDA	O1-N1	2.09	1.47	1.42
2	A	401	LDA	O1-N1	2.16	1.47	1.42
2	A	402	LDA	O1-N1	2.17	1.47	1.42

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	408	LDA	O1-N1-C1	-9.18	86.75	109.27
2	B	401	LDA	O1-N1-C1	-9.11	86.93	109.27
2	B	406	LDA	O1-N1-C1	-8.72	87.88	109.27
2	C	401	LDA	O1-N1-C1	-8.71	87.90	109.27
2	B	405	LDA	O1-N1-C1	-8.38	88.71	109.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	LDA	1	0
2	B	402	LDA	1	0
2	B	405	LDA	1	0
2	B	407	LDA	1	0
3	B	410	MAL	2	0
2	C	401	LDA	2	0
2	C	404	LDA	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	352/352 (100%)	-0.14	8 (2%) 60 31	65, 92, 146, 193	0
1	B	352/352 (100%)	-0.16	11 (3%) 49 21	66, 91, 142, 175	0
1	C	352/352 (100%)	-0.34	1 (0%) 93 83	69, 97, 139, 184	0
All	All	1056/1056 (100%)	-0.21	20 (1%) 66 37	65, 93, 145, 193	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	28	LYS	5.2
1	A	7	ASP	4.8
1	B	142	PHE	3.2
1	A	352	PHE	3.2
1	B	186	TRP	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	LDA	C	406	16/16	0.71	0.45	104,112,141,142	0
2	LDA	B	407	11/16	0.71	0.59	101,109,137,137	0
2	LDA	A	401	16/16	0.73	0.70	97,119,151,151	0
2	LDA	B	403	11/16	0.77	0.33	101,122,136,136	0
2	LDA	B	406	11/16	0.77	0.48	94,110,127,127	0
2	LDA	B	409	16/16	0.78	0.44	85,104,156,157	0
2	LDA	C	405	11/16	0.79	0.40	94,118,146,148	0
2	LDA	C	402	16/16	0.81	0.47	98,109,121,121	0
3	MAL	C	409	23/23	0.82	0.23	139,149,154,156	0
2	LDA	C	408	16/16	0.83	0.49	105,117,161,161	0
2	LDA	C	407	16/16	0.84	0.36	100,121,149,150	0
4	CA	B	411	1/1	0.84	0.48	101,101,101,101	0
2	LDA	A	402	16/16	0.87	0.42	88,111,145,145	0
3	MAL	A	403	23/23	0.87	0.18	115,124,128,129	0
2	LDA	B	405	15/16	0.89	0.26	87,93,127,131	0
2	LDA	C	401	13/16	0.89	0.40	93,102,120,124	0
3	MAL	B	410	23/23	0.90	0.18	103,116,118,119	0
2	LDA	B	402	16/16	0.91	0.34	93,101,119,122	0
2	LDA	B	408	16/16	0.91	0.51	90,101,152,154	0
2	LDA	C	404	16/16	0.92	0.24	78,102,130,131	0
2	LDA	B	404	10/16	0.93	0.32	87,88,93,94	0
2	LDA	B	401	16/16	0.93	0.28	65,94,129,132	0
2	LDA	C	403	16/16	0.93	0.29	82,90,101,101	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.