



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 05:49 pm GMT

PDB ID : 2O2Y  
Title : The crystal structure of *P. falciparum* enoyl acyl carrier protein reductase  
Authors : Muench, S.P.; Prigge, S.T.; McLeod, R.; Rafferty, J.B.; Kirisits, M.J.; Roberts, C.W.; Mui, E.J.; Rice, D.W.  
Deposited on : 2006-11-30  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 74%; height: 10px; background-color: green;"></div> <div style="width: 9%; height: 10px; background-color: yellow;"></div> <div style="width: 17%; height: 10px; background-color: grey;"></div> </div> </div> <div>74% 9% 17%</div> </div>
1	B	349	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 74%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: yellow;"></div> <div style="width: 17%; height: 10px; background-color: grey;"></div> </div> </div> <div>74% 8% 17%</div> </div>
1	C	349	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 75%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: yellow;"></div> <div style="width: 16%; height: 10px; background-color: grey;"></div> </div> </div> <div>75% 8% 16%</div> </div>
1	D	349	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; bottom: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 74%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: yellow;"></div> <div style="width: 17%; height: 10px; background-color: grey;"></div> </div> </div> <div>74% 8% 17%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9909 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-acyl carrier reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2288	1460	385	432	11			
1	B	290	Total	C	N	O	S	13	0	0
			2293	1463	385	434	11			
1	C	293	Total	C	N	O	S	18	0	0
			2310	1474	386	440	10			
1	D	288	Total	C	N	O	S	4	0	0
			2270	1451	378	430	11			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		
2	A	1	Total	Cl	0	0
			1	1		
2	D	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is TRICLOSAN (three-letter code: TCL) (formula: C<sub>12</sub>H<sub>7</sub>Cl<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	Cl	O	0	0
			17	12	3	2		
4	B	1	Total	C	Cl	O	0	0
			17	12	3	2		
4	C	1	Total	C	Cl	O	0	0
			17	12	3	2		
4	D	1	Total	C	Cl	O	0	0
			17	12	3	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	106	Total	O	0	0
			106	106		
6	B	130	Total	O	0	0
			130	130		
6	C	109	Total	O	0	0
			109	109		
6	D	150	Total	O	0	0
			150	150		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enoyl-acyl carrier reductase




- Molecule 1: Enoyl-acyl carrier reductase



- Molecule 1: Enoyl-acyl carrier reductase



● Molecule 1: Enoyl-acyl carrier reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.18Å 82.37Å 94.82Å 90.00° 90.77° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.89 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.7 (30.00-2.20) 97.7 (29.89-2.20)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.86 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.200 , 0.246 0.208 , 0.252	Depositor DCC
$R_{free}$ test set	3374 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 18.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.029 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9909	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TCL, SO4, NAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/2331	0.61	0/3147
1	B	0.61	3/2336 (0.1%)	0.77	6/3154 (0.2%)
1	C	0.61	5/2353 (0.2%)	0.79	8/3178 (0.3%)
1	D	0.58	1/2313 (0.0%)	0.70	3/3124 (0.1%)
All	All	0.57	9/9333 (0.1%)	0.72	17/12603 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	2
All	All	0	3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	60	LYS	CA-CB	-14.15	1.22	1.53
1	C	117	ASN	CG-OD1	-11.33	0.99	1.24
1	D	166	CYS	CB-SG	-10.70	1.64	1.82
1	C	42	LYS	CG-CD	-8.06	1.25	1.52
1	C	86	ALA	CA-CB	6.71	1.66	1.52
1	C	117	ASN	CG-ND2	6.47	1.49	1.32
1	B	133	LEU	CA-CB	5.87	1.67	1.53
1	B	68	MET	SD-CE	-5.36	1.47	1.77
1	C	68	MET	SD-CE	-5.21	1.48	1.77

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	68	MET	CG-SD-CE	20.00	132.19	100.20
1	D	81	MET	CG-SD-CE	15.30	124.68	100.20
1	B	81	MET	CG-SD-CE	13.58	121.92	100.20
1	B	198	MET	CG-SD-CE	12.95	120.92	100.20
1	C	53	ASN	OD1-CG-ND2	-10.96	96.68	121.90
1	B	68	MET	CG-SD-CE	10.16	116.46	100.20
1	C	42	LYS	CB-CG-CD	9.29	135.75	111.60
1	B	70	ILE	CG1-CB-CG2	-7.51	94.88	111.40
1	D	166	CYS	CA-CB-SG	7.03	126.66	114.00
1	C	53	ASN	CB-CG-OD1	6.22	134.04	121.60
1	D	235	ARG	NE-CZ-NH2	-6.02	117.29	120.30
1	C	86	ALA	N-CA-CB	-5.84	101.92	110.10
1	B	60	LYS	CB-CA-C	5.62	121.64	110.40
1	C	69	ILE	CG1-CB-CG2	-5.53	99.24	111.40
1	B	170	VAL	CG1-CB-CG2	-5.45	102.18	110.90
1	C	148	ARG	NE-CZ-NH2	5.19	122.90	120.30
1	C	148	ARG	NE-CZ-NH1	-5.17	117.72	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	133	LEU	Mainchain
1	C	117	ASN	Sidechain
1	C	53	ASN	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2288	0	2286	20	0
1	B	2293	0	2291	17	0
1	C	2310	0	2295	15	0
1	D	2270	0	2263	16	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	44	0	26	0	0
3	B	44	0	26	1	0
3	C	44	0	26	0	0
3	D	44	0	26	0	0
4	A	17	0	6	0	0
4	B	17	0	6	0	0
4	C	17	0	6	0	0
4	D	17	0	6	0	0
5	D	5	0	0	0	0
6	A	106	0	0	2	0
6	B	130	0	0	1	0
6	C	109	0	0	3	0
6	D	150	0	0	1	0
All	All	9909	0	9263	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:LEU:HD22	1:C:292:TYR:CE1	2.23	0.73
1:B:292:TYR:CD1	1:C:339:LEU:HD22	2.31	0.65
1:A:319:GLU:OE1	1:C:39:ARG:NH2	2.30	0.65
1:D:283:TYR:CE2	1:D:291:GLU:OE1	2.51	0.64
1:B:292:TYR:CE1	1:C:339:LEU:HD22	2.34	0.62
1:C:136:ALA:HB2	1:C:198:MET:SD	2.41	0.61
1:D:233:LYS:NZ	1:D:287:ASP:OD1	2.34	0.60
1:D:135:ASN:HD22	1:D:136:ALA:H	1.50	0.59
1:C:142:ASP:OD1	1:C:142:ASP:C	2.42	0.58
1:A:192:PRO:O	6:A:806:HOH:O	2.17	0.57
1:D:235:ARG:HB2	6:D:833:HOH:O	2.05	0.56
1:C:135:ASN:HD22	1:C:136:ALA:H	1.54	0.55
1:C:66:ASN:ND2	6:C:850:HOH:O	2.39	0.55
1:B:339:LEU:HD22	1:C:292:TYR:CD1	2.42	0.55
1:A:78:ILE:HG21	1:A:81:MET:HE2	1.89	0.55
1:B:135:ASN:HD22	1:B:136:ALA:H	1.54	0.55
1:A:288:TYR:CD1	1:D:343:ILE:HD12	2.42	0.54
1:D:169:PHE:CD2	1:D:172:ILE:HD11	2.44	0.53
1:D:233:LYS:HE3	1:D:238:THR:HG22	1.90	0.53
1:A:106:MET:HE3	1:A:107:LEU:CD2	2.40	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:ILE:HD12	1:A:45:PHE:CD1	2.44	0.52
1:D:109:ASN:HD22	1:D:114:ASP:HB3	1.76	0.51
1:D:15:ASP:HA	1:D:127:ASN:HD21	1.76	0.51
1:A:341:ASP:OD2	1:C:214:TYR:OH	2.27	0.50
1:A:78:ILE:HD13	1:A:81:MET:HE2	1.94	0.49
1:D:135:ASN:HD22	1:D:136:ALA:N	2.09	0.49
1:A:96:GLU:HA	1:A:96:GLU:OE1	2.13	0.48
1:A:87:SER:HB3	1:A:157:LYS:HE2	1.95	0.48
1:A:167:LYS:HD3	6:B:1525:HOH:O	2.14	0.47
1:B:169:PHE:O	1:B:173:MET:HG3	2.14	0.47
1:D:120:HIS:CD2	1:D:172:ILE:HG22	2.50	0.46
1:A:87:SER:CB	1:A:157:LYS:HE2	2.46	0.46
1:B:39:ARG:HD3	1:B:313:SER:OG	2.15	0.46
6:C:807:HOH:O	1:D:167:LYS:HD2	2.15	0.46
1:B:135:ASN:HD22	1:B:136:ALA:N	2.14	0.45
1:B:131:HIS:O	1:B:181:SER:HA	2.16	0.45
1:C:333:GLY:O	1:C:336:ILE:HG12	2.17	0.45
1:C:136:ALA:CB	1:C:198:MET:SD	3.05	0.45
1:C:237:ALA:HB1	1:C:286:ILE:HD13	1.99	0.45
1:A:78:ILE:HG21	1:A:81:MET:CE	2.45	0.45
1:A:292:TYR:OH	1:A:340:PRO:HG3	2.16	0.44
1:B:231:PRO:HA	3:B:1450:NAD:O7N	2.18	0.43
1:B:334:LEU:HD13	1:C:338:PHE:CZ	2.53	0.43
1:A:106:MET:CE	1:A:107:LEU:HD21	2.48	0.43
1:A:106:MET:HE3	1:A:107:LEU:HD23	2.01	0.43
1:B:109:ASN:HA	1:B:114:ASP:HB3	2.00	0.43
1:B:292:TYR:CE2	1:B:296:TYR:CD1	3.07	0.43
1:D:38:LYS:HG3	1:D:70:ILE:HG21	2.01	0.43
1:B:241:ASN:HB3	1:B:284:THR:HB	2.01	0.43
1:A:318:ARG:HD2	6:A:784:HOH:O	2.19	0.42
1:B:43:ILE:HG21	1:B:45:PHE:HE1	1.83	0.42
1:D:333:GLY:O	1:D:336:ILE:HG12	2.20	0.42
1:B:339:LEU:HD11	6:C:776:HOH:O	2.20	0.41
1:A:233:LYS:HG3	1:A:238:THR:CG2	2.51	0.41
1:D:131:HIS:O	1:D:181:SER:HA	2.19	0.41
1:B:182:LEU:HD23	1:B:227:ILE:HB	2.03	0.41
1:D:83:PRO:HB3	1:D:107:LEU:HD13	2.01	0.41
1:C:48:TRP:CG	1:C:50:PRO:HD2	2.56	0.40
1:A:131:HIS:HE1	1:A:161:SER:OG	2.04	0.40
1:A:140:GLN:HE22	1:A:241:ASN:N	2.19	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/349 (82%)	272 (95%)	13 (4%)	1 (0%)	43	48
1	B	286/349 (82%)	276 (96%)	9 (3%)	1 (0%)	43	48
1	C	289/349 (83%)	277 (96%)	10 (4%)	2 (1%)	24	23
1	D	284/349 (81%)	274 (96%)	10 (4%)	0	100	100
All	All	1145/1396 (82%)	1099 (96%)	42 (4%)	4 (0%)	43	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	72	LYS
1	C	14	GLU
1	B	72	LYS
1	A	240	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/307 (81%)	240 (97%)	8 (3%)	42	54
1	B	249/307 (81%)	243 (98%)	6 (2%)	52	65
1	C	249/307 (81%)	240 (96%)	9 (4%)	38	48
1	D	246/307 (80%)	239 (97%)	7 (3%)	47	59
All	All	992/1228 (81%)	962 (97%)	30 (3%)	44	56

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	SER
1	A	63	LYS
1	A	74	LYS
1	A	121	GLN
1	A	157	LYS
1	A	200	SER
1	A	301	GLN
1	A	306	THR
1	B	37	SER
1	B	135	ASN
1	B	149	LYS
1	B	172	ILE
1	B	241	ASN
1	B	334	LEU
1	C	75	LYS
1	C	77	ASN
1	C	102	LYS
1	C	135	ASN
1	C	142	ASP
1	C	170	VAL
1	C	300	ARG
1	C	302	LYS
1	C	345	ARG
1	D	37	SER
1	D	53	ASN
1	D	127	ASN
1	D	135	ASN
1	D	200	SER
1	D	235	ARG
1	D	342	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	100	ASN
1	A	109	ASN
1	A	127	ASN
1	A	131	HIS
1	A	140	GLN
1	A	301	GLN
1	B	117	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	135	ASN
1	B	176	GLN
1	B	215	HIS
1	B	301	GLN
1	C	66	ASN
1	C	101	ASN
1	C	135	ASN
1	C	140	GLN
1	C	221	ASN
1	D	53	ASN
1	D	58	ASN
1	D	61	ASN
1	D	100	ASN
1	D	109	ASN
1	D	127	ASN
1	D	135	ASN
1	D	140	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAD	A	650	-	40,48,48	1.66	4 (10%)	44,73,73	1.99	4 (9%)
4	TCL	A	705	-	18,18,18	1.77	2 (11%)	25,25,25	0.99	1 (4%)
3	NAD	B	1450	-	40,48,48	1.67	3 (7%)	44,73,73	1.80	2 (4%)
4	TCL	B	706	-	18,18,18	1.67	2 (11%)	25,25,25	1.13	2 (8%)
4	TCL	C	708	-	18,18,18	1.85	4 (22%)	25,25,25	1.11	1 (4%)
3	NAD	C	750	-	40,48,48	1.60	3 (7%)	44,73,73	2.05	5 (11%)
3	NAD	D	550	-	40,48,48	1.58	4 (10%)	44,73,73	2.21	6 (13%)
4	TCL	D	707	-	18,18,18	1.86	2 (11%)	25,25,25	0.98	1 (4%)
5	SO4	D	713	1	4,4,4	0.42	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	650	-	-	0/22/62/62	0/5/5/5
4	TCL	A	705	-	-	0/4/4/4	0/2/2/2
3	NAD	B	1450	-	-	0/22/62/62	0/5/5/5
4	TCL	B	706	-	-	0/4/4/4	0/2/2/2
4	TCL	C	708	-	-	0/4/4/4	0/2/2/2
3	NAD	C	750	-	-	0/22/62/62	0/5/5/5
3	NAD	D	550	-	-	0/22/62/62	0/5/5/5
4	TCL	D	707	-	-	0/4/4/4	0/2/2/2
5	SO4	D	713	1	-	0/0/0/0	0/0/0/0

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	650	NAD	O4B-C4B	-2.59	1.39	1.45
3	D	550	NAD	O4B-C4B	-2.50	1.39	1.45
4	C	708	TCL	C11-CL15	2.12	1.79	1.74
3	D	550	NAD	C2A-N1A	2.29	1.38	1.33
4	C	708	TCL	C2-CL14	2.37	1.79	1.74
3	C	750	NAD	C2A-N1A	2.63	1.38	1.33
3	A	650	NAD	C2A-N1A	2.63	1.38	1.33
3	B	1450	NAD	C2A-N1A	3.06	1.39	1.33
3	D	550	NAD	C2A-N3A	3.29	1.37	1.32
3	A	650	NAD	C2A-N3A	3.37	1.37	1.32
4	B	706	TCL	C6-C5	3.51	1.46	1.40

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	708	TCL	C6-C5	3.53	1.46	1.40
4	B	706	TCL	C8-C9	4.03	1.46	1.39
3	C	750	NAD	C2A-N3A	4.08	1.38	1.32
3	B	1450	NAD	C2A-N3A	4.18	1.38	1.32
4	A	705	TCL	C6-C5	4.20	1.47	1.40
4	C	708	TCL	C8-C9	4.33	1.47	1.39
4	A	705	TCL	C8-C9	4.36	1.47	1.39
4	D	707	TCL	C8-C9	4.39	1.47	1.39
4	D	707	TCL	C6-C5	4.88	1.48	1.40
3	D	550	NAD	O7N-C7N	7.43	1.39	1.24
3	C	750	NAD	O7N-C7N	7.50	1.39	1.24
3	B	1450	NAD	O7N-C7N	7.89	1.40	1.24
3	A	650	NAD	O7N-C7N	8.24	1.40	1.24

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	750	NAD	N3A-C2A-N1A	-11.34	119.16	128.86
3	A	650	NAD	N3A-C2A-N1A	-11.19	119.29	128.86
3	D	550	NAD	N3A-C2A-N1A	-10.62	119.77	128.86
3	B	1450	NAD	N3A-C2A-N1A	-9.98	120.32	128.86
3	D	550	NAD	C4B-O4B-C1B	-4.39	105.25	109.83
3	D	550	NAD	O7N-C7N-C3N	-3.77	114.90	119.62
3	D	550	NAD	C1B-N9A-C4A	-3.69	120.26	126.64
3	A	650	NAD	C4B-O4B-C1B	-3.59	106.09	109.83
3	C	750	NAD	C4B-O4B-C1B	-3.48	106.19	109.83
4	B	706	TCL	C8-C9-CL16	-2.76	116.17	119.42
3	C	750	NAD	O7N-C7N-C3N	-2.62	116.35	119.62
4	D	707	TCL	C1-C2-CL14	-2.23	116.30	119.15
3	A	650	NAD	O5B-C5B-C4B	-2.06	101.84	109.00
3	C	750	NAD	O2A-PA-O5B	2.04	117.23	107.75
3	A	650	NAD	C2A-N1A-C6A	2.05	122.24	118.75
4	C	708	TCL	C10-C9-CL16	2.09	121.79	118.48
4	A	705	TCL	C10-C9-CL16	2.39	122.27	118.48
3	B	1450	NAD	O2N-PN-O1N	2.48	124.76	112.14
3	D	550	NAD	O2N-PN-O1N	2.53	125.01	112.14
3	C	750	NAD	C3N-C7N-N7N	2.86	121.09	117.76
4	B	706	TCL	C10-C9-CL16	2.87	123.03	118.48
3	D	550	NAD	C3N-C7N-N7N	4.78	123.31	117.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1450	NAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	290/349 (83%)	-0.43	5 (1%) 70 68	17, 21, 25, 30	0
1	B	290/349 (83%)	-0.52	2 (0%) 87 86	17, 20, 25, 32	11 (3%)
1	C	293/349 (83%)	-0.37	5 (1%) 70 68	17, 21, 26, 33	15 (5%)
1	D	288/349 (82%)	-0.61	2 (0%) 87 86	17, 20, 25, 30	4 (1%)
All	All	1161/1396 (83%)	-0.48	14 (1%) 79 77	17, 20, 25, 33	30 (2%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	283	TYR	4.3
1	C	283	TYR	4.3
1	D	343	ILE	3.9
1	A	241	ASN	3.8
1	A	345	ARG	3.7
1	C	241	ASN	3.5
1	C	346	ASN	3.2
1	A	106	MET	2.8
1	C	13	ASN	2.5
1	A	73	ASP	2.5
1	D	344	TYR	2.3
1	C	102	LYS	2.2
1	B	72	LYS	2.2
1	B	241	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	SO4	D	713	5/5	0.85	0.27	46,47,48,49	0
3	NAD	A	650	44/44	0.95	0.11	20,21,24,24	0
3	NAD	C	750	44/44	0.96	0.09	17,21,24,26	0
4	TCL	A	705	17/17	0.96	0.10	21,22,22,24	0
4	TCL	D	707	17/17	0.97	0.09	21,23,26,27	0
3	NAD	D	550	44/44	0.97	0.10	20,22,25,28	0
3	NAD	B	1450	44/44	0.97	0.10	18,22,25,26	0
4	TCL	B	706	17/17	0.97	0.09	22,23,24,25	0
4	TCL	C	708	17/17	0.97	0.08	21,22,23,23	0
2	CL	C	712	1/1	0.99	0.04	23,23,23,23	0
2	CL	B	711	1/1	0.99	0.04	22,22,22,22	0
2	CL	A	710	1/1	1.00	0.06	14,14,14,14	0
2	CL	D	709	1/1	1.00	0.04	16,16,16,16	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.