



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 11:11 am GMT

PDB ID : 4O2X
Title : Structure of a malarial protein
Authors : AhYoung, A.P.; Koehl, A.; Cascio, D.; Egea, P.F.
Deposited on : 2013-12-17
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	507	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 12%, green 78%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 78% 12% 10% </div> </div>
1	B	507	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 12%, green 77%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 77% 12% 10% </div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, ATP-dependent Clp protease adaptor protein ClpS containing protein chimeric construct.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	457	Total	C	N	O	S	0	0	0
			3422	2202	555	658	7			
1	B	454	Total	C	N	O	S	0	1	0
			3411	2187	555	662	7			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	ASP	ALA	ENGINEERED MUTATION	UNP P0AEX9
A	84	LYS	ALA	ENGINEERED MUTATION	UNP P0AEX9
A	173	GLU	ALA	ENGINEERED MUTATION	UNP P0AEX9
A	174	ASN	ALA	ENGINEERED MUTATION	UNP P0AEX9
A	240	LYS	ALA	ENGINEERED MUTATION	UNP P0AEX9
A	360	GLU	ALA	ENGINEERED MUTATION	UNP P0AEX9
A	492	LEU	-	EXPRESSION TAG	UNP P0AEX9
A	493	VAL	-	EXPRESSION TAG	UNP P0AEX9
A	494	PRO	-	EXPRESSION TAG	UNP P0AEX9
A	495	ARG	-	EXPRESSION TAG	UNP P0AEX9
A	496	GLY	-	EXPRESSION TAG	UNP P0AEX9
A	497	SER	-	EXPRESSION TAG	UNP P0AEX9
A	498	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	499	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	500	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	501	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	502	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	503	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	504	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	505	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	506	HIS	-	EXPRESSION TAG	UNP P0AEX9
A	507	HIS	-	EXPRESSION TAG	UNP P0AEX9
B	83	ASP	ALA	ENGINEERED MUTATION	UNP P0AEX9
B	84	LYS	ALA	ENGINEERED MUTATION	UNP P0AEX9

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	173	GLU	ALA	ENGINEERED MUTATION	UNP P0AEX9
B	174	ASN	ALA	ENGINEERED MUTATION	UNP P0AEX9
B	240	LYS	ALA	ENGINEERED MUTATION	UNP P0AEX9
B	360	GLU	ALA	ENGINEERED MUTATION	UNP P0AEX9
B	492	LEU	-	EXPRESSION TAG	UNP Q8IEB2
B	493	VAL	-	EXPRESSION TAG	UNP Q8IEB2
B	494	PRO	-	EXPRESSION TAG	UNP Q8IEB2
B	495	ARG	-	EXPRESSION TAG	UNP Q8IEB2
B	496	GLY	-	EXPRESSION TAG	UNP Q8IEB2
B	497	SER	-	EXPRESSION TAG	UNP Q8IEB2
B	498	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	499	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	500	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	501	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	502	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	503	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	504	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	505	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	506	HIS	-	EXPRESSION TAG	UNP Q8IEB2
B	507	HIS	-	EXPRESSION TAG	UNP Q8IEB2


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein, ATP-dependent Clp protease adaptor protein ClpS containing protein chimeric construct



- Molecule 1: Maltose-binding periplasmic protein, ATP-dependent Clp protease adaptor protein ClpS containing protein chimeric construct



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	97.85Å 97.85Å 298.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	73.67 – 2.70 73.68 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.3 (73.67-2.70) 97.3 (73.68-2.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.69Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1555)	Depositor
R, R_{free}	0.185 , 0.239 0.189 , 0.240	Depositor DCC
R_{free} test set	3226 reflections (7.50%)	wwPDB-VP
Wilson B-factor (Å ²)	78.0	Xtriage
Anisotropy	0.163	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.064 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6833	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/3502	0.59	0/4784
1	B	0.43	0/3494	0.60	1/4776 (0.0%)
All	All	0.44	0/6996	0.60	1/9560 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	27	LYS	CB-CG-CD	-5.11	98.33	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3422	0	3268	37	0
1	B	3411	0	3232	31	0
All	All	6833	0	6500	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 68 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:PRO:HD2	1:A:225:MET:HE3	1.49	0.94
1:B:77:LEU:HB3	1:B:105:ILE:HD11	1.63	0.81
1:A:413:LEU:HB2	1:A:455:ILE:HD11	1.75	0.68
1:A:116:LEU:HD11	1:A:225:MET:HE2	1.81	0.63
1:A:458:THR:HG21	1:A:462:LYS:HD3	1.79	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/507 (89%)	425 (94%)	27 (6%)	1 (0%)	49	77
1	B	451/507 (89%)	423 (94%)	24 (5%)	4 (1%)	19	44
All	All	904/1014 (89%)	848 (94%)	51 (6%)	5 (1%)	27	54

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	174	ASN
1	B	415	ASN
1	B	418	ILE
1	A	63	TRP
1	B	99	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/418 (81%)	328 (97%)	9 (3%)	48	78
1	B	337/418 (81%)	322 (96%)	15 (4%)	30	60
All	All	674/836 (81%)	650 (96%)	24 (4%)	40	68

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	23	GLU
1	B	67	ARG
1	B	461	SER
1	B	42[A]	ASP
1	B	42[B]	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	457/507 (90%)	0.11	3 (0%) 87 88	51, 74, 124, 182	0
1	B	454/507 (89%)	0.13	3 (0%) 87 88	53, 79, 129, 183	0
All	All	911/1014 (89%)	0.12	6 (0%) 87 88	51, 76, 127, 183	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	100	TYR	3.8
1	B	68	PHE	3.3
1	A	175	GLY	2.6
1	B	173	GLU	2.2
1	A	93	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.