



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 06:55 pm GMT

PDB ID : 5O2A
Title : Fold Q98H
Authors : Koehnke, J.; Sikandar, A.
Deposited on : 2017-05-19
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5502 (1.90-1.90)
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>7% ..</div> </div> </div>
2	B	289	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>9% ..</div> </div> </div>
3	C	289	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>8% .</div> </div> </div>
4	D	289	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>8% ..</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 18413 atoms, of which 8879 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional protein Fold.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	285	Total	C	H	N	O	S	0	4	0
			4408	1375	2232	385	406	10			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P24186
A	1	GLY	-	expression tag	UNP P24186
A	98	HIS	GLN	engineered mutation	UNP P24186

- Molecule 2 is a protein called Bifunctional protein Fold.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	285	Total	C	H	N	O	S	0	3	0
			4395	1372	2226	385	402	10			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P24186
B	1	GLY	-	expression tag	UNP P24186
B	98	HIS	GLN	engineered mutation	UNP P24186

- Molecule 3 is a protein called Bifunctional protein Fold.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	284	Total	C	H	N	O	S	0	2	0
			4365	1361	2210	383	402	9			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP P24186
C	1	GLY	-	expression tag	UNP P24186
C	98	HIS	GLN	engineered mutation	UNP P24186

- Molecule 4 is a protein called Bifunctional protein FolD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	285	Total	C	H	N	O	S	0	1	0
			4369	1363	2211	385	402	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	MET	-	initiating methionine	UNP P24186
D	1	GLY	-	expression tag	UNP P24186
D	98	HIS	GLN	engineered mutation	UNP P24186


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	251	Total	O	0	0
			251	251		
5	B	203	Total	O	0	0
			203	203		
5	C	259	Total	O	0	0
			259	259		
5	D	163	Total	O	0	0
			163	163		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bifunctional protein Fold





- Molecule 2: Bifunctional protein Fold



- Molecule 3: Bifunctional protein Fold



- Molecule 4: Bifunctional protein Fold



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	100.11Å 79.39Å 101.78Å 90.00° 113.80° 90.00°	Depositor
Resolution (Å)	46.56 – 1.90 46.56 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.8 (46.56-1.90) 98.8 (46.56-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.30 (at 1.90Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.166 , 0.181 0.166 , 0.181	Depositor DCC
R_{free} test set	5718 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18413	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/2206	0.76	6/3006 (0.2%)
2	B	0.55	0/2193	0.70	3/2989 (0.1%)
3	C	0.61	1/2190 (0.0%)	0.74	5/2984 (0.2%)
4	D	0.48	1/2199 (0.0%)	0.73	3/2996 (0.1%)
All	All	0.56	2/8788 (0.0%)	0.73	17/11975 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	240	VAL	CB-CG1	-5.20	1.42	1.52
4	D	173	ARG	CD-NE	-5.07	1.37	1.46

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	173	ARG	NE-CZ-NH2	-18.88	110.86	120.30
1	A	173	ARG	NE-CZ-NH2	-12.89	113.85	120.30
4	D	173	ARG	NE-CZ-NH1	11.45	126.03	120.30
1	A	173	ARG	NE-CZ-NH1	10.44	125.52	120.30
1	A	139	ARG	NE-CZ-NH2	-9.38	115.61	120.30
2	B	99	LEU	C-N-CD	-8.38	102.16	120.60
3	C	51	VAL	CG1-CB-CG2	8.01	123.72	110.90
3	C	137	ARG	NE-CZ-NH1	7.95	124.28	120.30
3	C	137	ARG	NE-CZ-NH2	-6.96	116.82	120.30
4	D	173	ARG	CG-CD-NE	-6.53	98.08	111.80
3	C	173	ARG	NE-CZ-NH2	-5.94	117.33	120.30
2	B	139	ARG	NE-CZ-NH2	-5.93	117.33	120.30
3	C	51	VAL	CA-CB-CG1	5.75	119.53	110.90
1	A	130	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	177[A]	MET	CG-SD-CE	5.44	108.90	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	177[B]	MET	CG-SD-CE	5.44	108.90	100.20
2	B	196	LEU	CA-CB-CG	5.16	127.16	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2176	2232	2225	18	1
2	B	2169	2226	2228	19	1
3	C	2155	2210	2202	13	0
4	D	2158	2211	2213	11	0
5	A	251	0	0	6	2
5	B	203	0	0	7	0
5	C	259	0	0	4	2
5	D	163	0	0	2	0
All	All	9534	8879	8868	59	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:14:GLN:OE1	5:B:301:HOH:O	1.85	0.94
2:B:198:HIS:ND1	5:B:303:HOH:O	2.10	0.84
2:B:246:PHE:CD1	2:B:246:PHE:CZ	2.64	0.80
3:C:201:GLU:OE1	5:C:301:HOH:O	2.02	0.78
2:B:247:GLU:OE1	5:B:302:HOH:O	2.08	0.70
2:B:132[B]:CYS:SG	2:B:177[B]:MET:SD	2.92	0.67
2:B:195:ASN:ND2	5:B:305:HOH:O	2.28	0.67
2:B:81:GLU:OE1	5:B:304:HOH:O	2.13	0.66
5:A:356:HOH:O	2:B:133:GLN:HG3	1.96	0.65
1:A:248:ASP:HB3	5:A:309:HOH:O	1.96	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:123:PHE:HB3	5:B:306:HOH:O	2.02	0.59
4:D:69:ASP:OD1	5:D:301:HOH:O	2.16	0.59
4:D:40:LEU:HD22	4:D:42:GLY:O	2.05	0.56
1:A:132[B]:CYS:SG	2:B:132[B]:CYS:SG	3.04	0.55
1:A:47:SER:O	1:A:51:VAL:HG23	2.07	0.55
3:C:58[A]:CYS:SG	3:C:272:ILE:CD1	2.95	0.54
3:C:205:LEU:HD11	3:C:228:ILE:HD13	1.91	0.52
4:D:218:GLY:HA3	4:D:248:ASP:HB3	1.92	0.52
2:B:137:ARG:NH1	5:B:307:HOH:O	2.43	0.52
3:C:46:ALA:O	3:C:49:ILE:HD13	2.09	0.51
4:D:133:GLN:O	4:D:134:ARG:HG2	2.11	0.51
1:A:218:GLY:HA3	1:A:248:ASP:HB2	1.93	0.50
4:D:98:HIS:O	4:D:99:LEU:HB2	2.11	0.49
1:A:132[B]:CYS:SG	1:A:177[B]:MET:SD	3.09	0.49
1:A:40:LEU:HD11	1:A:51:VAL:HG22	1.94	0.48
1:A:40:LEU:HD21	1:A:51:VAL:HG22	1.96	0.47
2:B:236:GLU:O	2:B:237:ASN:CB	2.63	0.47
2:B:236:GLU:O	2:B:237:ASN:HB3	2.14	0.47
4:D:57:ALA:HB1	4:D:272:ILE:HD11	1.97	0.46
1:A:14:GLN:OE1	5:A:302:HOH:O	2.21	0.46
3:C:58[A]:CYS:SG	3:C:272:ILE:HD13	2.56	0.46
3:C:191:ARG:HG3	3:C:215:PHE:CZ	2.52	0.45
1:A:8:GLY:HA3	1:A:262:GLY:O	2.17	0.45
3:C:228:ILE:N	3:C:228:ILE:HD12	2.32	0.45
4:D:66:ARG:NH2	4:D:91:THR:OG1	2.50	0.44
3:C:226:ILE:HG22	3:C:228:ILE:HD11	1.98	0.44
3:C:170:ILE:HG12	5:C:306:HOH:O	2.17	0.44
1:A:177[A]:MET:HE1	2:B:173:ARG:HE	1.84	0.43
2:B:99:LEU:HD11	2:B:111:LEU:HD11	2.01	0.42
1:A:151:GLU:HG3	5:A:416:HOH:O	2.19	0.42
3:C:44:ASN:O	3:C:48:GLN:HG2	2.20	0.42
1:A:99:LEU:HD11	1:A:111:LEU:HD11	2.02	0.42
3:C:98:HIS:ND1	5:C:304:HOH:O	2.30	0.42
1:A:202:ASN:HB3	5:A:456:HOH:O	2.20	0.41
2:B:86:LEU:HD22	2:B:92:ILE:CD1	2.50	0.41
3:C:55:ARG:HD3	3:C:67:SER:OG	2.19	0.41
1:A:217:PRO:HG2	1:A:220:TRP:CE2	2.55	0.41
4:D:22:LYS:HD3	5:D:420:HOH:O	2.20	0.41
2:B:57:ALA:HB1	2:B:272:ILE:HD11	2.01	0.41
4:D:235:LEU:N	4:D:239:LYS:O	2.33	0.41
1:A:40:LEU:HD21	1:A:51:VAL:CG2	2.51	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:86:LEU:HD22	4:D:92:ILE:CD1	2.50	0.41
4:D:207:ILE:HD12	4:D:207:ILE:N	2.35	0.41
1:A:133:GLN:HB3	5:A:466:HOH:O	2.21	0.41
2:B:40:LEU:HD21	2:B:51:VAL:HG22	2.03	0.40
3:C:170:ILE:CG1	5:C:306:HOH:O	2.68	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:399:HOH:O	5:C:485:HOH:O[2_857]	1.91	0.29
5:A:316:HOH:O	5:C:494:HOH:O[1_455]	2.14	0.06
1:A:56:LYS:HZ2	2:B:156:ASP:OD2[2_747]	1.58	0.02
5:A:353:HOH:O	5:A:481:HOH:O[2_747]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/289 (99%)	279 (98%)	6 (2%)	0	100	100
2	B	284/289 (98%)	273 (96%)	10 (4%)	1 (0%)	36	25
3	C	283/289 (98%)	277 (98%)	6 (2%)	0	100	100
4	D	284/289 (98%)	271 (95%)	12 (4%)	1 (0%)	36	25
All	All	1136/1156 (98%)	1100 (97%)	34 (3%)	2 (0%)	49	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	237	ASN
4	D	99	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/235 (100%)	232 (98%)	4 (2%)	63	60
2	B	234/235 (100%)	228 (97%)	6 (3%)	49	42
3	C	234/236 (99%)	230 (98%)	4 (2%)	63	60
4	D	234/237 (99%)	225 (96%)	9 (4%)	36	26
All	All	938/943 (100%)	915 (98%)	23 (2%)	50	43

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LYS
1	A	65	SER
1	A	121	ASP
1	A	241	VAL
2	B	40	LEU
2	B	59	GLU
2	B	65	SER
2	B	69	ASP
2	B	121	ASP
2	B	219	ASP
3	C	51	VAL
3	C	60	GLU
3	C	65	SER
3	C	219	ASP
4	D	13	GLN
4	D	40	LEU
4	D	59	GLU
4	D	65	SER
4	D	69	ASP
4	D	121	ASP
4	D	181	LEU
4	D	234	ARG
4	D	248	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
4	D	283	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	MLY	A	194	1	10,10,11	1.05	1 (10%)	8,11,13	0.83	0
1	MLY	A	22	1	10,10,11	1.30	1 (10%)	8,11,13	1.04	0
2	MLY	B	212	2	10,10,11	0.83	0	8,11,13	1.04	0
2	MLY	B	222	2	10,10,11	0.88	0	8,11,13	1.09	0
3	MLY	C	4	3	10,10,11	1.14	1 (10%)	8,11,13	1.16	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	194	1	-	0/7/9/11	0/0/0/0
1	MLY	A	22	1	-	0/7/9/11	0/0/0/0
2	MLY	B	212	2	-	0/7/9/11	0/0/0/0
2	MLY	B	222	2	-	0/7/9/11	0/0/0/0
3	MLY	C	4	3	-	0/7/9/11	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	194	MLY	CA-C	2.41	1.53	1.50
3	C	4	MLY	CA-C	3.00	1.54	1.50
1	A	22	MLY	CA-C	3.69	1.55	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	4	MLY	CD-CE-NZ	-2.09	108.15	113.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/289 (97%)	-0.29	3 (1%) 80 83	19, 28, 53, 78	0
2	B	283/289 (97%)	-0.14	6 (2%) 63 66	19, 31, 55, 90	0
3	C	283/289 (97%)	-0.23	3 (1%) 80 83	19, 27, 47, 63	0
4	D	285/289 (98%)	0.03	15 (5%) 26 30	23, 37, 63, 118	0
All	All	1134/1156 (98%)	-0.16	27 (2%) 59 62	19, 30, 56, 118	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	238	GLY	6.8
4	D	237	ASN	6.4
1	A	286	GLN	4.9
4	D	239	LYS	4.8
4	D	236	GLU	4.0
1	A	285	PRO	3.8
2	B	123	PHE	3.7
2	B	236	GLU	3.5
4	D	285	PRO	3.4
4	D	240	VAL	3.3
4	D	235	LEU	3.2
4	D	1	GLY	3.2
2	B	239	LYS	3.1
2	B	237	ASN	3.0
1	A	284	ASP	2.9
4	D	31	LEU	2.8
4	D	13	GLN	2.5
3	C	285	PRO	2.5
4	D	123	PHE	2.5
3	C	284	ASP	2.5
4	D	28	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	234	ARG	2.4
2	B	285	PRO	2.4
3	C	57	ALA	2.3
4	D	45	PRO	2.2
4	D	241	VAL	2.2
2	B	235	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MLY	B	212	11/12	0.94	0.10	26,44,59,59	0
1	MLY	A	194	11/12	0.96	0.11	21,28,42,42	0
1	MLY	A	22	11/12	0.96	0.15	20,37,55,55	0
3	MLY	C	4	11/12	0.96	0.11	24,42,68,68	0
2	MLY	B	222	11/12	0.97	0.10	22,28,35,35	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.