



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 16, 2020 – 01:15 PM EDT

PDB ID : 6P5C  
Title : Bacillus Fragment DNA polymerase mutant I716M  
Authors : Wu, E.Y.  
Deposited on : 2019-05-30  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.8  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.8


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	9	 22% (red), 67% (green), 33% (yellow)
2	C	9	 11% (red), 100% (green)
3	A	581	 4% (red), 93% (green), 6% (yellow)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	904	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5188 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*CP\*GP\*AP\*TP\*CP\*AP\*CP\*GP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	9	Total	C	N	O	P	0	0	0
			179	86	34	51	8			

- Molecule 2 is a DNA chain called DNA (5'-D(P\*GP\*CP\*GP\*TP\*GP\*AP\*TP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	P	0	0	0
			187	88	35	55	9			

- Molecule 3 is a protein called DNA polymerase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	581	Total	C	N	O	S	0	0	0
			4658	2960	808	871	19			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	296	MET	-	initiating methionine	UNP D9N168
A	297	ALA	-	expression tag	UNP D9N168
A	598	ASP	ALA	conflict	UNP D9N168
A	713	VAL	PRO	conflict	UNP D9N168
A	716	MET	ILE	engineered mutation	UNP D9N168

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	O	0	0
			1	1		
5	C	8	Total	O	0	0
			8	8		
5	A	135	Total	O	0	0
			135	135		



- Molecule 1: DNA (5'-D(\*CP\*GP\*AP\*TP\*CP\*AP\*CP\*GP\*C)-3')



- 

- 
- The dot plot displays the frequency of 1000 random numbers. The x-axis represents the value of the random number, ranging from 0 to 1000. The y-axis represents the frequency of each value, ranging from 0 to 100. The plot shows a uniform distribution of points across the range, with a slight increase in frequency around the value 500.

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.22Å 93.52Å 105.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.34 – 2.20 28.32 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.4 (28.34-2.20) 96.4 (28.32-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.41 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.228 , 0.273 0.234 , 0.277	Depositor DCC
$R_{free}$ test set	2134 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtriage
Anisotropy	0.358	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 30.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	5188	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	0.55	0/200	0.92	1/306 (0.3%)
2	C	0.52	0/209	0.80	0/321
3	A	0.71	0/4742	0.85	3/6407 (0.0%)
All	All	0.70	0/5151	0.86	4/7034 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	843	ARG	NE-CZ-NH2	-6.39	117.10	120.30
1	B	27	DC	O5'-P-OP2	-6.22	100.10	105.70
3	A	859	ARG	NE-CZ-NH1	5.83	123.21	120.30
3	A	831	GLU	CB-CA-C	5.04	120.48	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	179	0	102	1	0
2	C	187	0	102	0	0
3	A	4658	0	4705	22	0
4	A	20	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	135	0	0	3	0
5	B	1	0	0	0	0
5	C	8	0	0	0	0
All	All	5188	0	4909	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:530:SER:HB2	3:A:533:GLN:HB2	1.74	0.69
3:A:587:TYR:O	3:A:591:LEU:HB2	1.94	0.66
3:A:561:LEU:O	3:A:571:VAL:HG11	2.00	0.62
3:A:591:LEU:O	3:A:595:VAL:HG23	1.99	0.61
3:A:692:VAL:HG21	3:A:701:MET:CE	2.35	0.57
3:A:296:MET:O	3:A:296:MET:HE3	2.12	0.49
3:A:534:LEU:HD11	3:A:574:ILE:HD13	1.94	0.49
3:A:692:VAL:HG21	3:A:701:MET:HE2	1.95	0.48
3:A:843:ARG:HD2	5:A:1127:HOH:O	2.14	0.47
3:A:692:VAL:HG21	3:A:701:MET:HE1	1.97	0.47
3:A:649:ILE:O	3:A:868:TYR:HA	2.16	0.46
1:B:22:DG:H4'	1:B:23:DA:OP1	2.16	0.45
3:A:440:GLU:N	3:A:441:PRO:CD	2.79	0.45
3:A:764:THR:HA	3:A:769:ARG:O	2.17	0.45
3:A:467:ARG:NH2	5:A:1013:HOH:O	2.51	0.44
3:A:459:ARG:HA	3:A:459:ARG:NE	2.33	0.44
3:A:503:MET:CE	3:A:636:ILE:HD13	2.48	0.43
3:A:692:VAL:HB	3:A:696:GLU:HB2	2.00	0.43
3:A:296:MET:CE	3:A:296:MET:O	2.67	0.43
3:A:680:ASP:OD1	3:A:680:ASP:C	2.57	0.43
3:A:812:ASN:ND2	5:A:1010:HOH:O	2.48	0.43
3:A:551:LYS:HA	3:A:551:LYS:HD2	1.90	0.43
3:A:296:MET:SD	4:A:903:SO4:O4	2.77	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	579/581 (100%)	558 (96%)	20 (4%)	1 (0%)	49 57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	628	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	496/497 (100%)	480 (97%)	16 (3%)	42 54

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	349	GLU
3	A	517	ARG
3	A	532	LYS
3	A	551	LYS
3	A	557	SER
3	A	591	LEU
3	A	593	LYS
3	A	716	MET
3	A	718	ASP
3	A	738	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	A	770	ARG
3	A	776	ILE
3	A	779	ARG
3	A	812	ASN
3	A	831	GLU
3	A	876	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	A	901	-	4,4,4	0.34	0	6,6,6	0.29	0
4	SO4	A	902	-	4,4,4	0.29	0	6,6,6	0.10	0
4	SO4	A	903	-	4,4,4	0.28	0	6,6,6	0.17	0
4	SO4	A	904	-	4,4,4	0.32	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	9/9 (100%)	0.48	2 (22%) 0 0	24, 37, 66, 87	0
2	C	9/9 (100%)	-0.19	1 (11%) 5 4	21, 34, 49, 56	0
3	A	581/581 (100%)	-0.05	21 (3%) 42 41	12, 25, 64, 112	0
All	All	599/599 (100%)	-0.04	24 (4%) 38 36	12, 25, 65, 112	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	552	THR	8.8
3	A	719	TYR	7.3
3	A	433	ALA	5.0
3	A	434	LYS	4.7
3	A	551	LYS	4.6
3	A	520	GLU	4.5
3	A	432	GLY	4.3
1	B	21	DC	4.2
3	A	553	GLY	3.6
3	A	729	ARG	3.5
3	A	531	PRO	3.5
3	A	523	GLY	3.4
3	A	431	LYS	3.2
3	A	550	THR	2.9
3	A	525	GLU	2.8
3	A	319	VAL	2.8
3	A	549	LYS	2.6
3	A	448	VAL	2.6
3	A	535	GLY	2.4
1	B	22	DG	2.2
2	C	14	DG	2.2
3	A	548	LYS	2.2
3	A	723	GLN	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	A	532	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	A	902	5/5	0.67	0.34	75,99,104,106	0
4	SO4	A	904	5/5	0.77	0.50	113,114,123,130	0
4	SO4	A	901	5/5	0.88	0.15	49,49,56,59	0
4	SO4	A	903	5/5	0.98	0.10	32,33,34,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.