



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2019 – 12:26 AM EDT

PDB ID : 6PF6  
Title : Crystal structure of TS-DHFR from *Cryptosporidium hominis* in complex with NADPH, FdUMP and 2-(4-((2-amino-4-oxo-4,7-dihydro-3H-pyrrolo[2,3-d]pyrimidin-5-yl)methyl)benzamido)terephthalic acid  
Authors : Czyzyk, D.L.; Valhondo, M.; Jorgensen, W.L.; Anderson, K.S.  
Deposited on : 2019-06-21  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.4  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4155 (2.50-2.50)
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	<div> <div>3%</div> <div>87%</div> <div>11%</div> <div>.</div> </div>
1	B	290	<div> <div>2%</div> <div>92%</div> <div>7%</div> <div>.</div> </div>
1	C	290	<div> <div>3%</div> <div>91%</div> <div>8%</div> <div>.</div> </div>
1	D	290	<div> <div>6%</div> <div>92%</div> <div>6%</div> <div>.</div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidylate synthase,Thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	285	Total	C	N	O	S	0	0	0
			2221	1424	384	400	13			
1	A	283	Total	C	N	O	S	0	0	0
			2197	1407	375	404	11			
1	C	285	Total	C	N	O	S	0	0	0
			2221	1423	380	405	13			
1	D	283	Total	C	N	O	S	0	0	0
			2194	1405	372	406	11			

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula:  $C_9H_{13}N_2O_8P$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	D	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is 2-({4-[(2-amino-4-oxo-4,7-dihydro-3H-pyrrolo[2,3-d]pyrimidin-5-yl)methyl]benzene-1-carbonyl}amino)benzene-1,4-dicarboxylic acid (three-letter code: OEJ) (formula: C<sub>22</sub>H<sub>17</sub>N<sub>5</sub>O<sub>6</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			33	22	5	6		
3	C	1	Total	C	N	O	0	0
			33	22	5	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	6	Total	O	0	0
			6	6		
4	A	5	Total	O	0	0
			5	5		
4	C	6	Total	O	0	0
			6	6		
4	D	5	Total	O	0	0
			5	5		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Thymidylate synthase, Thymidylate synthase

Chain B: 





- Molecule 1: Thymidylate synthase, Thymidylate synthase

Chain A: 





- Molecule 1: Thymidylate synthase, Thymidylate synthase

Chain C: 



- Molecule 1: Thymidylate synthase, Thymidylate synthase

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	152.22Å 152.22Å 107.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 2.50 48.13 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.13-2.50) 99.7 (48.13-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.18	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.16 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.15.2_3472: 000)	Depositor
R, $R_{free}$	0.225 , 0.264 0.225 , 0.265	Depositor DCC
$R_{free}$ test set	2187 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.7	Xtriage
Anisotropy	0.389	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9001	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OEJ, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/2254	0.42	0/3067
1	B	0.25	0/2278	0.42	0/3096
1	C	0.25	0/2278	0.42	0/3097
1	D	0.25	0/2251	0.42	0/3066
All	All	0.25	0/9061	0.42	0/12326

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2197	0	2086	17	0
1	B	2221	0	2133	12	0
1	C	2221	0	2115	11	0
1	D	2194	0	2070	11	0
2	A	20	0	11	0	0
2	B	20	0	11	0	0
2	C	20	0	11	0	0
2	D	20	0	11	2	0
3	B	33	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	33	0	0	0	0
4	A	5	0	0	0	0
4	B	6	0	0	0	0
4	C	6	0	0	0	0
4	D	5	0	0	0	0
All	All	9001	0	8448	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:ARG:NH2	1:A:126:ARG:O	2.30	0.65
1:C:30:GLU:HG3	1:C:74:LEU:HD22	1.81	0.61
1:B:196:HIS:HB2	1:B:212:LEU:HD11	1.85	0.59
1:A:140:ARG:NH2	1:A:289:ASP:OD1	2.37	0.58
1:C:140:ARG:NH2	1:C:289:ASP:OD1	2.36	0.58
1:D:214:GLN:HB3	1:D:252:LEU:HD23	1.87	0.56
1:A:214:GLN:HB3	1:A:252:LEU:HD23	1.90	0.54
1:A:5:GLY:HA2	1:A:32:GLN:NE2	2.24	0.53
1:D:115:ARG:NH2	1:D:126:ARG:O	2.42	0.53
1:A:30:GLU:HG3	1:A:74:LEU:HD22	1.91	0.53
1:B:184:PRO:HD2	1:A:142:PHE:CE1	2.44	0.53
1:B:214:GLN:HB3	1:B:252:LEU:HD23	1.90	0.53
1:A:196:HIS:HB2	1:A:212:LEU:HD11	1.90	0.52
1:D:256:HIS:CE1	2:D:600:UMP:HO3'	2.25	0.52
1:A:3:VAL:HA	1:A:31:LEU:HD12	1.92	0.52
1:D:196:HIS:HB2	1:D:212:LEU:HD11	1.91	0.52
1:A:285:VAL:HG13	1:A:290:ASP:HB2	1.92	0.51
1:A:182:TRP:HB2	1:A:197:ALA:HB1	1.91	0.51
1:C:196:HIS:CD2	1:C:196:HIS:H	2.29	0.51
1:C:196:HIS:HB2	1:C:212:LEU:HD11	1.94	0.49
1:D:55:THR:HB	1:D:256:HIS:HB2	1.94	0.49
1:B:140:ARG:HH11	1:B:161:LEU:HD23	1.77	0.49
1:B:140:ARG:NH1	1:B:161:LEU:HD23	2.28	0.49
1:B:142:PHE:CE1	1:A:184:PRO:HD2	2.47	0.49
1:C:268:GLN:HB2	1:C:307:ILE:HD12	1.94	0.49
1:D:50:ARG:HH21	2:D:600:UMP:H5'	1.79	0.48
1:D:74:LEU:HD12	1:D:224:PRO:HB3	1.96	0.48
1:D:115:ARG:NH1	1:D:119:ASP:OD1	2.47	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:285:VAL:HG13	1:C:290:ASP:HB2	1.98	0.46
1:B:140:ARG:NH2	1:B:289:ASP:OD1	2.48	0.46
1:B:30:GLU:HG3	1:B:74:LEU:HD22	1.98	0.46
1:B:251:THR:HG21	1:A:251:THR:HG21	1.99	0.45
1:A:5:GLY:HA2	1:A:32:GLN:HE21	1.81	0.44
1:D:285:VAL:HG13	1:D:290:ASP:HB2	2.00	0.44
1:A:69:ASP:OD2	1:A:278:LYS:NZ	2.50	0.43
1:C:214:GLN:HB3	1:C:252:LEU:HD23	1.99	0.43
1:B:55:THR:HB	1:B:256:HIS:HB2	2.00	0.43
1:A:55:THR:HB	1:A:256:HIS:HB2	2.01	0.43
1:A:68:ARG:NH1	1:A:247:ASP:OD1	2.45	0.43
1:B:274:ARG:HD2	1:B:302:ASN:O	2.19	0.42
1:C:55:THR:HB	1:C:256:HIS:HB2	2.01	0.42
1:B:115:ARG:NH1	1:B:119:ASP:OD1	2.53	0.42
1:A:193:PRO:HA	1:A:194:PRO:HD3	1.92	0.41
1:C:193:PRO:HA	1:C:194:PRO:HD3	1.96	0.41
1:D:30:GLU:HG3	1:D:74:LEU:HD22	2.01	0.41
1:D:140:ARG:HH11	1:D:161:LEU:HD23	1.85	0.41
1:C:126:ARG:NH2	1:C:189:LEU:O	2.46	0.41
1:C:37:ILE:HD12	1:C:265:LEU:HD13	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/290 (97%)	268 (95%)	13 (5%)	0	100	100
1	B	283/290 (98%)	275 (97%)	8 (3%)	0	100	100
1	C	283/290 (98%)	274 (97%)	9 (3%)	0	100	100
1	D	281/290 (97%)	272 (97%)	9 (3%)	0	100	100
All	All	1128/1160 (97%)	1089 (96%)	39 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/252 (90%)	225 (99%)	2 (1%)	81	93
1	B	230/252 (91%)	229 (100%)	1 (0%)	92	98
1	C	229/252 (91%)	227 (99%)	2 (1%)	81	93
1	D	226/252 (90%)	226 (100%)	0	100	100
All	All	912/1008 (90%)	907 (100%)	5 (0%)	90	97

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	50	ARG
1	A	50	ARG
1	A	263	GLU
1	C	42	ARG
1	C	50	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	211	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	UMP	A	600	-	17,21,21	1.69	3 (17%)	22,31,31	1.52	3 (13%)
2	UMP	B	600	-	17,21,21	1.70	3 (17%)	22,31,31	1.52	4 (18%)
3	OEJ	B	601	-	31,36,36	1.62	5 (16%)	35,52,52	2.27	6 (17%)
2	UMP	C	600	-	17,21,21	1.49	2 (11%)	22,31,31	1.49	2 (9%)
3	OEJ	C	601	-	31,36,36	1.63	5 (16%)	35,52,52	2.28	6 (17%)
2	UMP	D	600	-	17,21,21	1.66	3 (17%)	22,31,31	1.59	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMP	A	600	-	-	1/6/22/22	0/2/2/2
2	UMP	B	600	-	-	2/6/22/22	0/2/2/2
3	OEJ	B	601	-	-	1/12/20/20	0/4/4/4
2	UMP	C	600	-	-	3/6/22/22	0/2/2/2
3	OEJ	C	601	-	-	0/12/20/20	0/4/4/4
2	UMP	D	600	-	-	2/6/22/22	0/2/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	601	OEJ	C20-C21	4.72	1.52	1.47

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	OEJ	C20-C21	4.40	1.51	1.47
2	A	600	UMP	C4-N3	4.04	1.40	1.33
2	C	600	UMP	C4-N3	4.02	1.40	1.33
2	B	600	UMP	C4-N3	3.98	1.40	1.33
2	D	600	UMP	C4-N3	3.83	1.39	1.33
2	B	600	UMP	P-OP1	3.65	1.62	1.50
2	D	600	UMP	P-OP1	3.60	1.62	1.50
2	A	600	UMP	P-OP1	3.57	1.62	1.50
3	B	601	OEJ	C17-C22	3.40	1.50	1.47
3	C	601	OEJ	C17-C22	3.17	1.50	1.47
3	C	601	OEJ	C4-N2	3.16	1.41	1.35
3	B	601	OEJ	C4-N2	3.14	1.41	1.35
2	C	600	UMP	C6-N1	2.76	1.39	1.35
2	A	600	UMP	C6-N1	2.71	1.39	1.35
2	B	600	UMP	C6-N1	2.69	1.39	1.35
2	D	600	UMP	C6-N1	2.51	1.39	1.35
3	B	601	OEJ	C4-N4	2.30	1.38	1.33
3	C	601	OEJ	C4-N4	2.22	1.38	1.33
3	B	601	OEJ	C2-C1	-2.10	1.37	1.43
3	C	601	OEJ	C2-C1	-2.09	1.37	1.43

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	601	OEJ	C2-C3-N2	-10.03	115.11	124.08
3	B	601	OEJ	C2-C3-N2	-9.72	115.39	124.08
3	C	601	OEJ	C3-N2-C4	4.64	122.67	116.06
3	B	601	OEJ	C3-N2-C4	4.55	122.53	116.06
3	B	601	OEJ	N4-C4-N1	4.08	124.51	117.77
2	D	600	UMP	C5-C4-N3	-3.93	114.48	123.28
2	A	600	UMP	C5-C4-N3	-3.89	114.56	123.28
2	C	600	UMP	C5-C4-N3	-3.86	114.63	123.28
2	B	600	UMP	C5-C4-N3	-3.82	114.73	123.28
3	C	601	OEJ	C3-C2-C1	3.81	117.16	115.04
3	B	601	OEJ	N4-C4-N2	-3.80	111.30	117.25
3	C	601	OEJ	N4-C4-N1	3.80	124.04	117.77
2	D	600	UMP	OP2-P-O5'	3.66	116.47	106.73
3	C	601	OEJ	N4-C4-N2	-3.62	111.59	117.25
3	B	601	OEJ	C3-C2-C1	3.53	117.01	115.04
2	A	600	UMP	OP2-P-O5'	3.44	115.88	106.73
2	C	600	UMP	O5'-P-OP1	3.41	116.05	106.47
2	B	600	UMP	OP2-P-O5'	3.36	115.67	106.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	OEJ	N1-C4-N2	-2.27	124.18	127.25
2	D	600	UMP	OP3-P-OP2	2.22	116.19	107.57
2	B	600	UMP	OP3-P-OP1	-2.13	102.13	110.53
2	A	600	UMP	OP3-P-OP1	-2.13	102.14	110.53
3	C	601	OEJ	N1-C4-N2	-2.13	124.37	127.25
2	D	600	UMP	OP3-P-OP1	-2.11	102.21	110.53
2	B	600	UMP	OP3-P-OP2	2.04	115.50	107.57

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	600	UMP	O4'-C4'-C5'-O5'
2	D	600	UMP	C3'-C4'-C5'-O5'
2	C	600	UMP	O4'-C4'-C5'-O5'
2	C	600	UMP	C3'-C4'-C5'-O5'
2	B	600	UMP	O4'-C4'-C5'-O5'
2	C	600	UMP	C5'-O5'-P-OP2
2	A	600	UMP	O4'-C4'-C5'-O5'
2	B	600	UMP	C3'-C4'-C5'-O5'
3	B	601	OEJ	C5-C7-C8-C9

There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	600	UMP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand OEJ B 601



## Ligand UMP C 600





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	283/290 (97%)	0.39	10 (3%) 44 47	55, 67, 80, 97	0
1	B	285/290 (98%)	0.29	5 (1%) 68 71	55, 67, 87, 98	0
1	C	285/290 (98%)	0.42	10 (3%) 44 47	59, 71, 95, 102	0
1	D	283/290 (97%)	0.53	18 (6%) 19 20	59, 74, 91, 106	0
All	All	1136/1160 (97%)	0.41	43 (3%) 40 43	55, 69, 89, 106	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	305	PRO	5.4
1	A	4	ALA	5.1
1	A	306	THR	4.5
1	C	121	LEU	4.4
1	D	306	THR	4.4
1	C	113	GLY	4.3
1	B	313	VAL	4.1
1	A	3	VAL	4.0
1	C	123	PHE	4.0
1	D	131	LEU	3.3
1	B	52	GLY	3.2
1	A	282	LEU	3.1
1	D	267	ILE	2.8
1	B	106	VAL	2.8
1	D	106	VAL	2.8
1	A	195	CYS	2.7
1	D	110	ASP	2.7
1	A	2	PRO	2.7
1	D	117	PHE	2.7
1	D	109	TRP	2.6
1	C	117	PHE	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	114	SER	2.5
1	C	253	GLY	2.4
1	A	281	ILE	2.4
1	C	221	LEU	2.3
1	D	239	HIS	2.3
1	B	285	VAL	2.3
1	D	98	ALA	2.3
1	D	285	VAL	2.2
1	D	120	SER	2.2
1	D	2	PRO	2.2
1	C	126	ARG	2.2
1	D	240	ILE	2.2
1	A	286	GLU	2.2
1	D	286	GLU	2.2
1	C	285	VAL	2.1
1	A	305	PRO	2.1
1	C	187	LEU	2.1
1	A	290	ASP	2.1
1	B	129	GLY	2.1
1	D	199	CYS	2.1
1	D	129	GLY	2.1
1	D	111	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	OEJ	C	601	33/33	0.82	0.22	76,82,88,89	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	OEJ	B	601	33/33	0.88	0.21	66,71,81,82	0
2	UMP	B	600	20/20	0.93	0.21	61,66,69,73	0
2	UMP	C	600	20/20	0.95	0.17	65,73,80,80	0
2	UMP	A	600	20/20	0.96	0.16	61,67,71,72	0
2	UMP	D	600	20/20	0.97	0.17	66,70,74,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around OEJ C 601:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OEJ B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UMP B 600:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around UMP C 600:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UMP A 600:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.