



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 11:31 pm GMT

PDB ID : 3PM9
Title : Crystal structure of a Putative dehydrogenase (RPA1076) from Rhodopseudomonas palustris CGA009 at 2.57 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2010-11-16
Resolution : 2.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3182 (2.60-2.56)
Clashscore	122126	3541 (2.60-2.56)
Ramachandran outliers	120053	3489 (2.60-2.56)
Sidechain outliers	120020	3489 (2.60-2.56)
RSRZ outliers	108989	3120 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div></div> </div> <div></div> </div>
1	B	476	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>10%</div> <div></div> </div> <div></div> </div>
1	C	476	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div></div> </div> <div></div> </div>
1	D	476	<div> <div>0%</div> <div> <div></div> <div>87%</div> <div>10%</div> <div></div> </div> <div></div> </div>
1	E	476	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div></div> </div> <div></div> </div>
1	F	476	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>10%</div> <div></div> </div> <div></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNL	A	477	-	-	X	-
3	UNL	B	477	-	-	X	-
3	UNL	C	477	-	-	X	-
3	UNL	D	477	-	-	X	-
3	UNL	E	477	-	-	X	-
3	UNL	F	477	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 22350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative oxidoreductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	Se	0	3	0
			3487	2208	600	671	2	6			
1	B	465	Total	C	N	O	S	Se	0	4	0
			3493	2212	602	671	2	6			
1	C	465	Total	C	N	O	S	Se	0	5	0
			3499	2215	602	674	2	6			
1	D	465	Total	C	N	O	S	Se	0	5	0
			3501	2216	609	668	2	6			
1	E	465	Total	C	N	O	S	Se	0	6	0
			3509	2220	608	673	2	6			
1	F	465	Total	C	N	O	S	Se	0	4	0
			3498	2213	605	672	2	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	leader sequence	UNP Q6NAV4
B	0	GLY	-	leader sequence	UNP Q6NAV4
C	0	GLY	-	leader sequence	UNP Q6NAV4
D	0	GLY	-	leader sequence	UNP Q6NAV4
E	0	GLY	-	leader sequence	UNP Q6NAV4
F	0	GLY	-	leader sequence	UNP Q6NAV4

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is UNKNOWN LIGAND (three-letter code: UNL) (formula:).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	O	0	0
			6	6		
3	E	1	Total	O	0	0
			6	6		
3	B	1	Total	O	0	0
			6	6		
3	C	1	Total	O	0	0
			6	6		
3	A	1	Total	O	0	0
			6	6		
3	F	1	Total	O	0	0
			6	6		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	C	1	Total	O	P	0	0
			5	4	1		
4	C	1	Total	O	P	0	0
			5	4	1		
4	C	1	Total	O	P	0	0
			5	4	1		
4	D	1	Total	O	P	0	0
			5	4	1		
4	D	1	Total	O	P	0	0
			5	4	1		
4	D	1	Total	O	P	0	0
			5	4	1		
4	E	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	O	P	0	0
			5	4	1		
4	E	1	Total	O	P	0	0
			5	4	1		
4	F	1	Total	O	P	0	0
			5	4	1		
4	F	1	Total	O	P	0	0
			5	4	1		
4	F	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	203	Total	O	0	0
			203	203		
5	B	168	Total	O	0	0
			168	168		
5	C	126	Total	O	0	0
			126	126		
5	D	143	Total	O	0	0
			143	143		
5	E	145	Total	O	0	0
			145	145		
5	F	134	Total	O	0	0
			134	134		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Putative oxidoreductase




• Molecule 1: Putative oxidoreductase



• Molecule 1: Putative oxidoreductase





• Molecule 1: Putative oxidoreductase







• Molecule 1: Putative oxidoreductase



• Molecule 1: Putative oxidoreductase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	146.09Å 250.73Å 251.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.36 – 2.57 49.36 – 2.57	Depositor EDS
% Data completeness (in resolution range)	98.9 (49.36-2.57) 98.9 (49.36-2.57)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.193 , 0.227 0.190 , 0.220	Depositor DCC
R_{free} test set	7270 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	43.7	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	22350	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.3365e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, UNL, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	1/3549 (0.0%)	0.64	1/4814 (0.0%)
1	B	0.66	0/3558	0.62	0/4826
1	C	0.69	1/3567 (0.0%)	0.61	0/4838
1	D	0.66	0/3569	0.63	0/4840
1	E	0.67	1/3580 (0.0%)	0.63	0/4855
1	F	0.67	1/3563 (0.0%)	0.62	0/4833
All	All	0.67	4/21386 (0.0%)	0.63	1/29006 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	407	GLU	CG-CD	6.02	1.60	1.51
1	C	61	GLU	CG-CD	5.90	1.60	1.51
1	A	61	GLU	CG-CD	5.73	1.60	1.51
1	E	72	GLU	CG-CD	5.73	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	LEU	CA-CB-CG	7.31	132.11	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3487	0	3506	29	0
1	B	3493	0	3516	31	0
1	C	3499	0	3520	27	0
1	D	3501	0	3539	29	0
1	E	3509	0	3540	24	0
1	F	3498	0	3522	30	0
2	A	53	0	31	3	0
2	B	53	0	31	3	0
2	C	53	0	31	4	0
2	D	53	0	31	3	0
2	E	53	0	31	2	0
2	F	53	0	31	2	0
3	A	6	0	0	3	0
3	B	6	0	0	4	0
3	C	6	0	0	4	0
3	D	6	0	0	4	0
3	E	6	0	0	3	0
3	F	6	0	0	2	0
4	A	15	0	0	0	0
4	B	15	0	0	0	0
4	C	15	0	0	1	0
4	D	15	0	0	1	0
4	E	15	0	0	0	0
4	F	15	0	0	0	0
5	A	203	0	0	3	0
5	B	168	0	0	2	0
5	C	126	0	0	2	0
5	D	143	0	0	1	0
5	E	145	0	0	2	0
5	F	134	0	0	0	0
All	All	22350	0	21329	194	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:477:UNL:O1	3:F:477:UNL:O3	1.64	1.15
3:B:477:UNL:O2	3:B:477:UNL:O3	1.66	1.12

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:477:UNL:O3	3:C:477:UNL:O2	1.68	1.11
3:D:477:UNL:O3	3:D:477:UNL:O4	1.78	0.99
3:A:477:UNL:O6	3:A:477:UNL:O5	1.85	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/476 (98%)	453 (97%)	13 (3%)	0	100	100
1	B	467/476 (98%)	454 (97%)	12 (3%)	1 (0%)	49	72
1	C	468/476 (98%)	456 (97%)	12 (3%)	0	100	100
1	D	468/476 (98%)	450 (96%)	16 (3%)	2 (0%)	36	59
1	E	469/476 (98%)	456 (97%)	11 (2%)	2 (0%)	36	59
1	F	467/476 (98%)	453 (97%)	12 (3%)	2 (0%)	36	59
All	All	2805/2856 (98%)	2722 (97%)	76 (3%)	7 (0%)	43	72

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	45	ASN
1	E	45	ASN
1	F	146	GLN
1	D	45	ASN
1	D	384	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/372 (97%)	357 (99%)	5 (1%)	69	86
1	B	363/372 (98%)	359 (99%)	4 (1%)	76	89
1	C	365/372 (98%)	363 (100%)	2 (0%)	90	96
1	D	365/372 (98%)	362 (99%)	3 (1%)	83	93
1	E	366/372 (98%)	360 (98%)	6 (2%)	65	83
1	F	364/372 (98%)	362 (100%)	2 (0%)	90	96
All	All	2185/2232 (98%)	2163 (99%)	22 (1%)	81	90

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	234	PHE
1	D	234	PHE
1	F	234	PHE
1	C	237	LEU
1	D	143	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	154	ASN
1	B	364	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 30 ligands modelled in this entry, 6 are unknown - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	476	-	51,58,58	1.66	8 (15%)	57,89,89	2.29	11 (19%)
4	PO4	A	478	-	4,4,4	0.78	0	6,6,6	0.44	0
4	PO4	A	484	-	4,4,4	0.72	0	6,6,6	0.35	0
4	PO4	A	490	-	4,4,4	0.76	0	6,6,6	0.51	0
2	FAD	B	476	-	51,58,58	1.53	8 (15%)	57,89,89	2.21	13 (22%)
4	PO4	B	479	-	4,4,4	0.75	0	6,6,6	0.31	0
4	PO4	B	485	-	4,4,4	0.55	0	6,6,6	0.58	0
4	PO4	B	491	-	4,4,4	1.02	0	6,6,6	0.52	0
2	FAD	C	476	-	51,58,58	1.65	8 (15%)	57,89,89	2.24	10 (17%)
4	PO4	C	480	-	4,4,4	0.66	0	6,6,6	0.44	0
4	PO4	C	486	-	4,4,4	0.68	0	6,6,6	0.59	0
4	PO4	C	492	-	4,4,4	0.88	0	6,6,6	0.47	0
2	FAD	D	476	-	51,58,58	1.69	9 (17%)	57,89,89	2.23	11 (19%)
4	PO4	D	481	-	4,4,4	0.74	0	6,6,6	0.40	0
4	PO4	D	487	-	4,4,4	0.70	0	6,6,6	0.53	0
4	PO4	D	493	-	4,4,4	0.71	0	6,6,6	0.50	0
2	FAD	E	476	-	51,58,58	1.64	9 (17%)	57,89,89	2.20	11 (19%)
4	PO4	E	482	-	4,4,4	0.75	0	6,6,6	0.35	0
4	PO4	E	488	-	4,4,4	0.66	0	6,6,6	0.35	0
4	PO4	E	494	-	4,4,4	0.66	0	6,6,6	0.40	0
2	FAD	F	476	-	51,58,58	1.58	7 (13%)	57,89,89	2.26	10 (17%)
4	PO4	F	483	-	4,4,4	0.69	0	6,6,6	0.53	0
4	PO4	F	489	-	4,4,4	0.58	0	6,6,6	0.41	0
4	PO4	F	495	-	4,4,4	0.78	0	6,6,6	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	476	-	-	0/28/50/50	0/6/6/6
4	PO4	A	478	-	-	0/0/0/0	0/0/0/0
4	PO4	A	484	-	-	0/0/0/0	0/0/0/0
4	PO4	A	490	-	-	0/0/0/0	0/0/0/0
2	FAD	B	476	-	-	0/28/50/50	0/6/6/6
4	PO4	B	479	-	-	0/0/0/0	0/0/0/0
4	PO4	B	485	-	-	0/0/0/0	0/0/0/0
4	PO4	B	491	-	-	0/0/0/0	0/0/0/0
2	FAD	C	476	-	-	0/28/50/50	0/6/6/6
4	PO4	C	480	-	-	0/0/0/0	0/0/0/0
4	PO4	C	486	-	-	0/0/0/0	0/0/0/0
4	PO4	C	492	-	-	0/0/0/0	0/0/0/0
2	FAD	D	476	-	-	0/28/50/50	0/6/6/6
4	PO4	D	481	-	-	0/0/0/0	0/0/0/0
4	PO4	D	487	-	-	0/0/0/0	0/0/0/0
4	PO4	D	493	-	-	0/0/0/0	0/0/0/0
2	FAD	E	476	-	-	0/28/50/50	0/6/6/6
4	PO4	E	482	-	-	0/0/0/0	0/0/0/0
4	PO4	E	488	-	-	0/0/0/0	0/0/0/0
4	PO4	E	494	-	-	0/0/0/0	0/0/0/0
2	FAD	F	476	-	-	0/28/50/50	0/6/6/6
4	PO4	F	483	-	-	0/0/0/0	0/0/0/0
4	PO4	F	489	-	-	0/0/0/0	0/0/0/0
4	PO4	F	495	-	-	0/0/0/0	0/0/0/0

The worst 5 of 49 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	476	FAD	C2'-C3'	-3.32	1.47	1.53
2	E	476	FAD	C2'-C3'	-2.94	1.47	1.53
2	B	476	FAD	C2'-C3'	-2.75	1.48	1.53
2	D	476	FAD	C2'-C3'	-2.49	1.48	1.53
2	F	476	FAD	C2'-C3'	-2.46	1.48	1.53

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	476	FAD	N3A-C2A-N1A	-11.03	119.42	128.86
2	F	476	FAD	N3A-C2A-N1A	-10.68	119.72	128.86
2	A	476	FAD	N3A-C2A-N1A	-10.66	119.73	128.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	476	FAD	N3A-C2A-N1A	-10.51	119.87	128.86
2	D	476	FAD	N3A-C2A-N1A	-10.05	120.26	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	476	FAD	3	0
2	B	476	FAD	3	0
2	C	476	FAD	4	0
4	C	492	PO4	1	0
2	D	476	FAD	3	0
4	D	493	PO4	1	0
2	E	476	FAD	2	0
2	F	476	FAD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	459/476 (96%)	-0.01	10 (2%) 62 58	23, 35, 54, 72	0
1	B	459/476 (96%)	0.00	10 (2%) 62 58	23, 36, 56, 73	0
1	C	459/476 (96%)	0.04	11 (2%) 59 55	26, 39, 59, 77	0
1	D	459/476 (96%)	-0.03	7 (1%) 73 71	25, 37, 57, 75	0
1	E	459/476 (96%)	-0.03	11 (2%) 59 55	24, 36, 56, 72	0
1	F	459/476 (96%)	0.03	13 (2%) 53 49	24, 38, 59, 77	0
All	All	2754/2856 (96%)	-0.00	62 (2%) 60 56	23, 37, 57, 77	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	43	GLU	5.3
1	A	43	GLU	4.6
1	B	34	LEU	4.5
1	F	11	THR	4.3
1	B	45	ASN	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	UNL	A	477	6/-	0.78	0.33	36,52,57,67	0
4	PO4	C	492	5/5	0.86	0.27	40,40,41,41	5
3	UNL	E	477	6/-	0.87	0.29	25,50,56,57	0
3	UNL	D	477	6/-	0.87	0.29	29,48,51,53	0
3	UNL	F	477	6/-	0.88	0.34	30,49,54,55	0
3	UNL	B	477	6/-	0.89	0.24	38,44,50,56	0
3	UNL	C	477	6/-	0.90	0.22	33,47,51,54	0
4	PO4	D	493	5/5	0.90	0.24	40,40,41,41	5
4	PO4	A	490	5/5	0.90	0.27	35,35,36,37	5
4	PO4	B	491	5/5	0.91	0.22	29,29,30,31	5
4	PO4	E	494	5/5	0.92	0.23	43,43,44,44	5
4	PO4	C	486	5/5	0.92	0.34	33,34,34,34	5
4	PO4	D	487	5/5	0.92	0.30	41,42,43,43	5
4	PO4	F	489	5/5	0.92	0.35	42,42,43,43	5
4	PO4	E	488	5/5	0.92	0.34	36,37,38,38	5
4	PO4	B	485	5/5	0.94	0.39	35,35,35,36	5
4	PO4	A	484	5/5	0.94	0.31	32,32,33,34	5
4	PO4	F	495	5/5	0.95	0.23	43,44,44,44	5
4	PO4	E	482	5/5	0.96	0.13	50,51,51,52	0
2	FAD	C	476	53/53	0.97	0.12	28,35,45,48	0
2	FAD	D	476	53/53	0.97	0.13	28,33,44,46	0
2	FAD	F	476	53/53	0.97	0.13	26,33,45,48	0
4	PO4	C	480	5/5	0.97	0.14	51,51,52,53	0
2	FAD	E	476	53/53	0.97	0.13	26,32,43,45	0
2	FAD	B	476	53/53	0.98	0.13	25,32,43,45	0
4	PO4	F	483	5/5	0.98	0.14	46,46,47,48	0
4	PO4	B	479	5/5	0.98	0.17	49,49,50,50	0
2	FAD	A	476	53/53	0.98	0.14	24,29,42,44	0
4	PO4	A	478	5/5	0.98	0.12	53,54,54,55	0
4	PO4	D	481	5/5	0.99	0.13	54,54,54,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.