



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 06:01 am GMT

PDB ID : 3PMN
Title : ternary crystal structure of polymerase lambda variant with a GT mispair at the primer terminus with Mn²⁺ in the active site
Authors : Bebenek, K.; Pedersen, L.C.; Kunkel, T.A.
Deposited on : 2010-11-17
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div> <div>7%</div> <div> <div></div> <div>72%</div> <div>22%</div> <div>5%</div> </div> </div>
2	B	11	<div> <div>82%</div> <div>18%</div> </div>
3	C	6	<div> <div>100%</div> </div>
4	D	4	<div> <div>50%</div> <div>50%</div> </div>

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 3143 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	1	5	0
			2481	1566	450	454	11			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	463	LYS	SER	SEE REMARK 999	UNP Q9UGP5
A	464	GLY	GLN	SEE REMARK 999	UNP Q9UGP5
A	?	-	GLU	SEE REMARK 999	UNP Q9UGP5
A	?	-	GLU	SEE REMARK 999	UNP Q9UGP5
A	?	-	ASN	SEE REMARK 999	UNP Q9UGP5
A	?	-	GLY	SEE REMARK 999	UNP Q9UGP5
A	?	-	GLN	SEE REMARK 999	UNP Q9UGP5
A	470	GLU	GLN	SEE REMARK 999	UNP Q9UGP5
A	471	THR	GLN	SEE REMARK 999	UNP Q9UGP5

- Molecule 2 is a DNA chain called 5'-D(*CP*GP*GP*CP*CP*TP*TP*AP*CP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	11	Total	C	N	O	P	0	0	0
			220	106	38	66	10			


- Molecule 3 is a DNA chain called 5'-D(*CP*AP*GP*TP*AP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	6	Total	C	N	O	P	0	0	0
			122	59	25	33	5			

- Molecule 4 is a DNA chain called 5'-D(*GP*CP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	4	Total	C	N	O	P	1	0	0
			79	38	16	22	3			

- Molecule 5 is 2'-deoxy-5'-O-[(R)-hydroxy{[(S)-hydroxy(phosphonooxy)phosphoryl]methyl}phosphoryl]guanosine (three-letter code: 1GC) (formula: C₁₁H₁₈N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

- Molecule 7 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	3	Total	Mn	0	0
			3	3		

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	2	Total	Na	0	0
			2	2		

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	2	Total 2	Cl 2	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	146	Total 146	O 146	0	0
10	B	27	Total 27	O 27	0	0
10	C	21	Total 21	O 21	0	0
10	D	8	Total 8	O 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase lambda



- Molecule 2: 5'-D(*CP*GP*GP*CP*CP*TP*TP*AP*CP*TP*G)-3'



- Molecule 3: 5'-D(*CP*AP*GP*TP*AP*G)-3'



There are no outlier residues recorded for this chain.

- Molecule 4: 5'-D(*GP*CP*CP*G)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.39Å 62.42Å 139.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.27 – 2.20 25.27 – 2.20	Depositor EDS
% Data completeness (in resolution range)	89.0 (25.27-2.20) 89.1 (25.27-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.60 (at 2.19Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.213 , 0.243 0.219 , 0.246	Depositor DCC
R_{free} test set	1287 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3143	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, MN, 1GC, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2533	0.54	0/3417
2	B	0.33	0/245	0.75	0/376
3	C	0.34	0/137	0.72	0/210
4	D	0.23	0/88	0.68	0/134
All	All	0.32	0/3003	0.58	0/4137

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2481	0	2461	60	0
2	B	220	0	126	6	0
3	C	122	0	69	0	0
4	D	79	0	46	1	0
5	A	31	0	14	0	0
6	A	1	0	0	0	0
7	A	3	0	0	0	0
8	A	2	0	0	0	0
9	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	A	146	0	0	4	0
10	B	27	0	0	0	0
10	C	21	0	0	0	0
10	D	8	0	0	0	0
All	All	3143	0	2716	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:536:VAL:HG12	1:A:547:PRO:HA	1.26	1.12
1:A:252:HIS:HB3	1:A:292:PRO:HG3	1.72	0.70
1:A:557:LYS:HG2	1:A:567:TYR:CD2	2.30	0.67
1:A:464:GLY:HA3	2:B:9:DC:H5''	1.76	0.66
1:A:279:TYR:OH	1:A:312:LYS:HE3	1.97	0.65
1:A:557:LYS:HG3	10:A:604:HOH:O	1.98	0.64
1:A:261:GLU:HG2	1:A:283:ILE:HD13	1.79	0.62
1:A:464:GLY:CA	2:B:9:DC:H5''	2.29	0.61
1:A:290:HIS:ND1	1:A:291:LYS:N	2.51	0.59
1:A:431:LEU:HD23	1:A:432:ILE:N	2.18	0.58
1:A:443:ILE:O	1:A:444:PHE:C	2.43	0.57
1:A:266:ALA:O	1:A:270:GLN:HG3	2.05	0.57
4:D:3:DC:H2''	4:D:4:DG:C8	2.41	0.56
1:A:434:HIS:CD2	1:A:439:SER:HB2	2.40	0.56
1:A:462:VAL:CG1	2:B:9:DC:H4'	2.36	0.55
1:A:536:VAL:HG12	1:A:547:PRO:CA	2.18	0.55
1:A:464:GLY:HA3	2:B:9:DC:C5'	2.38	0.54
1:A:257:THR:O	1:A:261:GLU:HG3	2.08	0.54
1:A:325:LEU:HD12	10:A:634:HOH:O	2.08	0.54
1:A:462:VAL:HG13	2:B:9:DC:H4'	1.90	0.53
1:A:501:CYS:SG	1:A:531:ALA:HA	2.47	0.53
1:A:321:HIS:CE1	10:A:634:HOH:O	2.61	0.53
1:A:256:ILE:HD13	1:A:313:ILE:HG23	1.91	0.53
1:A:315:GLU:O	1:A:319:SER:HB2	2.09	0.53
1:A:251:ASN:OD1	1:A:287:LYS:HA	2.08	0.52
1:A:252:HIS:HB3	1:A:292:PRO:CG	2.39	0.52
1:A:313:ILE:O	1:A:317:LEU:HD13	2.10	0.51
1:A:449:ASP:O	1:A:453:GLN:HG3	2.10	0.51
1:A:513:ASN:O	1:A:517:ARG:HG3	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:THR:OG1	1:A:373:GLN:HG3	2.11	0.50
1:A:264:ALA:HB2	1:A:279:TYR:HB3	1.94	0.50
1:A:516:MET:HG2	1:A:527:LEU:CD2	2.43	0.48
1:A:471:THR:O	1:A:492:ILE:HA	2.13	0.48
1:A:434:HIS:O	1:A:496:TYR:HB2	2.13	0.48
1:A:293:VAL:HA	1:A:298:GLU:OE2	2.15	0.47
1:A:519:LEU:CD2	1:A:565:LEU:HD11	2.45	0.47
1:A:431:LEU:C	1:A:431:LEU:HD23	2.36	0.45
1:A:443:ILE:HG13	1:A:444:PHE:N	2.31	0.45
1:A:443:ILE:HG13	1:A:444:PHE:H	1.82	0.44
1:A:253:ASN:HB3	1:A:256:ILE:HD12	2.00	0.44
1:A:263:LEU:HD21	1:A:324:LYS:HG2	2.00	0.44
1:A:519:LEU:HD23	1:A:565:LEU:HD11	2.00	0.44
1:A:252:HIS:HB3	1:A:292:PRO:CD	2.48	0.43
1:A:295:SER:O	1:A:298:GLU:HB3	2.19	0.43
1:A:443:ILE:O	1:A:446:ARG:N	2.51	0.43
1:A:429:ASP:HB3	1:A:506:PHE:CZ	2.54	0.43
1:A:315:GLU:HG2	1:A:321:HIS:O	2.19	0.43
1:A:557:LYS:HG2	1:A:567:TYR:CE2	2.53	0.42
1:A:415:CYS:HA	1:A:419:ARG:HB2	2.01	0.42
1:A:289:PHE:CZ	1:A:302:ILE:HD11	2.54	0.42
1:A:407:ASN:HB3	1:A:410:LEU:HG	2.02	0.42
1:A:521:LYS:HE3	2:B:6:DT:OP1	2.20	0.42
1:A:534:THR:O	1:A:548:GLY:HA3	2.20	0.41
1:A:302:ILE:HA	1:A:303:PRO:HD3	1.93	0.41
1:A:359:SER:O	1:A:362:ASP:HB2	2.20	0.41
1:A:251:ASN:HD21	1:A:254:LEU:HA	1.86	0.41
1:A:440:HIS:O	1:A:443:ILE:HG12	2.20	0.41
1:A:311:GLU:O	1:A:314:ILE:HB	2.22	0.40
1:A:471:THR:HB	1:A:493:VAL:O	2.22	0.40
1:A:311:GLU:HB2	10:A:614:HOH:O	2.21	0.40
1:A:307:LYS:HG2	1:A:311:GLU:OE2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	312/329 (95%)	299 (96%)	12 (4%)	1 (0%)	43 48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	444	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	259/275 (94%)	257 (99%)	2 (1%)	83 91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	444	PHE
1	A	506	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	251	ASN
1	A	453	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	1GC	A	1	7,6	28,33,33	2.12	5 (17%)	31,52,52	2.28	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	1GC	A	1	7,6	-	0/15/34/34	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1	1GC	PB-O3B	-6.66	1.50	1.58
5	A	1	1GC	PG-O3B	-5.39	1.51	1.60
5	A	1	1GC	PA-O1A	-3.03	1.49	1.56
5	A	1	1GC	PB-O2B	-2.94	1.49	1.56
5	A	1	1GC	C6-N1	3.96	1.40	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1	1GC	C5-C6-N1	-8.41	111.51	123.47
5	A	1	1GC	C2-N3-C4	-2.70	112.01	115.16
5	A	1	1GC	N3-C2-N1	-2.34	123.98	127.41
5	A	1	1GC	PG-O3B-PB	-2.32	124.25	132.40
5	A	1	1GC	O1A-PA-O2A	2.51	118.35	110.04
5	A	1	1GC	O2B-PB-O1B	2.57	118.58	110.04
5	A	1	1GC	O2G-PG-O1G	2.92	119.15	107.59
5	A	1	1GC	C6-N1-C2	6.18	124.95	116.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/329 (94%)	0.24	22 (7%) 16 15	20, 38, 79, 93	0
2	B	11/11 (100%)	-0.81	0 100 100	27, 30, 45, 48	0
3	C	6/6 (100%)	-0.94	0 100 100	21, 23, 24, 30	0
4	D	4/4 (100%)	-0.67	0 100 100	37, 38, 41, 43	0
All	All	333/350 (95%)	0.18	22 (6%) 18 17	20, 38, 79, 93	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	249	ALA	6.9
1	A	319	SER	6.4
1	A	290	HIS	5.9
1	A	547	PRO	4.8
1	A	250	THR	4.5
1	A	430	VAL	3.5
1	A	491	ILE	3.2
1	A	289	PHE	2.9
1	A	536	VAL	2.9
1	A	453	GLN	2.9
1	A	254	LEU	2.7
1	A	489	LEU	2.5
1	A	575[A]	TRP	2.4
1	A	534	THR	2.4
1	A	431	LEU	2.3
1	A	449	ASP	2.2
1	A	344	ALA	2.2
1	A	470	GLU	2.1
1	A	252	HIS	2.1
1	A	535	ALA	2.1
1	A	492	ILE	2.1
1	A	313	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	NA	A	10	1/1	0.68	0.15	67,67,67,67	0
9	CL	A	8	1/1	0.91	0.07	55,55,55,55	0
9	CL	A	9	1/1	0.96	0.08	54,54,54,54	0
6	MG	A	576	1/1	0.96	0.08	17,17,17,17	0
7	MN	A	7	1/1	0.97	0.03	85,85,85,85	0
5	1GC	A	1	31/31	0.97	0.10	20,23,25,26	0
8	NA	A	5	1/1	0.98	0.10	24,24,24,24	0
7	MN	A	6	1/1	1.00	0.04	45,45,45,45	0
7	MN	A	2	1/1	1.00	0.08	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.