



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 19, 2019 – 10:49 AM EDT

PDB ID : 6PSM  
Title : Crystal structure of PsS1\_19B C77S in complex with kappa-neocarrabiose  
Authors : Hettle, A.G.; Boraston, A.B.  
Deposited on : 2019-07-12  
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.4  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2220 (1.96-1.96)
Clashscore	122126	2333 (1.96-1.96)
Ramachandran outliers	120053	2314 (1.96-1.96)
Sidechain outliers	120020	2314 (1.96-1.96)
RSRZ outliers	108989	2174 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	 91% • 5%
1	B	469	 91% • 5%
1	C	469	 91% 5% •
1	D	469	 91% • 5%
1	E	469	 92% • •

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	469	<div><div></div><div>92%</div><div></div><div>• 5%</div></div>

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 22255 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called exo-4S-kappa carrageenan S1 sulfatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	2	0
			3537	2259	594	674	10			
1	B	447	Total	C	N	O	S	0	0	0
			3526	2254	593	670	9			
1	C	448	Total	C	N	O	S	0	0	0
			3540	2260	599	672	9			
1	D	444	Total	C	N	O	S	0	1	0
			3505	2240	591	666	8			
1	E	448	Total	C	N	O	S	0	1	0
			3545	2264	599	673	9			
1	F	447	Total	C	N	O	S	0	0	1
			3522	2249	593	671	9			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Ca	0	0
			1	1		
2	E	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		
2	F	1	Total	Ca	0	0
			1	1		

- Molecule 3 is 4-O-sulfo-beta-D-galactopyranose (three-letter code: G4S) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>9</sub>S) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			16	6	9	1		
3	B	1	Total	C	O	S	0	0
			16	6	9	1		
3	C	1	Total	C	O	S	0	0
			16	6	9	1		
3	D	1	Total	C	O	S	0	0
			16	6	9	1		
3	E	1	Total	C	O	S	0	0
			16	6	9	1		
3	F	1	Total	C	O	S	0	0
			16	6	9	1		

- Molecule 4 is 3,6-anhydro-D-galactose (three-letter code: 9RN) (formula: C<sub>6</sub>H<sub>10</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		
4	C	1	Total	C	O	0	0
			10	6	4		
4	D	1	Total	C	O	0	0
			10	6	4		
4	E	1	Total	C	O	0	0
			10	6	4		
4	F	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	Cl	0	0
			1	1		
5	E	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		
5	C	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		
5	F	1	Total	Cl	0	0
			1	1		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Zn	0	0
			2	2		
6	D	2	Total	Zn	0	0
			2	2		
6	E	2	Total	Zn	0	0
			2	2		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	C	1	Total	C	O	0	0
			4	2	2		
7	D	1	Total	C	O	0	0
			4	2	2		
7	E	1	Total	C	O	0	0
			4	2	2		
7	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	140	Total	O	0	0
			140	140		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	121	Total 121	O 121	0	0
8	C	163	Total 163	O 163	0	0
8	D	131	Total 131	O 131	0	0
8	E	145	Total 145	O 145	0	0
8	F	186	Total 186	O 186	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: exo-4S-kappa carrageenan S1 sulfatase




- Molecule 1: exo-4S-kappa carrageenan S1 sulfatase



- Molecule 1: exo-4S-kappa carrageenan S1 sulfatase



- Molecule 1: exo-4S-kappa carrageenan S1 sulfatase





• Molecule 1: exo-4S-kappa carrageenan S1 sulfatase

Chain E: 92%



• Molecule 1: exo-4S-kappa carrageenan S1 sulfatase

Chain F: 92% 5%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.05Å 93.05Å 300.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.19 – 1.95 48.15 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.19-1.95) 99.7 (48.15-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.06 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, $R_{free}$	0.192 , 0.212 0.198 , 0.216	Depositor DCC
$R_{free}$ test set	10434 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.8	Xtriage
Anisotropy	0.703	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 17.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	0.110 for -h,-k,l 0.114 for h,-h-k,-l 0.178 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	22255	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 9RN, CL, CA, EDO, G4S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/3630	0.76	0/4909
1	B	0.67	0/3616	0.76	0/4889
1	C	0.68	0/3630	0.77	0/4904
1	D	0.67	0/3594	0.76	0/4856
1	E	0.69	0/3635	0.76	0/4912
1	F	0.69	0/3611	0.78	0/4882
All	All	0.68	0/21716	0.77	0/29352

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3537	0	3383	10	0
1	B	3526	0	3374	12	0
1	C	3540	0	3412	16	0
1	D	3505	0	3359	11	0
1	E	3545	0	3406	12	0
1	F	3522	0	3370	13	0
2	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	16	0	10	0	0
3	B	16	0	10	0	0
3	C	16	0	10	0	0
3	D	16	0	10	0	0
3	E	16	0	10	0	0
3	F	16	0	10	0	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
4	C	10	0	0	0	0
4	D	10	0	0	0	0
4	E	10	0	0	0	0
4	F	10	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
6	A	2	0	0	0	0
6	D	2	0	0	0	0
6	E	2	0	0	0	0
7	A	4	0	6	0	0
7	C	4	0	6	0	0
7	D	4	0	6	0	0
7	E	4	0	6	0	0
7	F	4	0	6	0	0
8	A	140	0	0	1	0
8	B	121	0	0	0	0
8	C	163	0	0	0	0
8	D	131	0	0	0	0
8	E	145	0	0	1	0
8	F	186	0	0	0	0
All	All	22255	0	20394	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:248:ASN:HB3	1:F:130:GLU:OE1	1.76	0.85
1:B:85:MET:HE1	1:B:123:THR:HB	1.75	0.67
1:B:85:MET:CE	1:B:123:THR:HB	2.28	0.63
1:D:85:MET:CE	1:D:123:THR:HB	2.29	0.63
1:C:85:MET:CE	1:C:123:THR:HB	2.28	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/469 (95%)	433 (97%)	14 (3%)	0	100	100
1	B	445/469 (95%)	432 (97%)	13 (3%)	0	100	100
1	C	446/469 (95%)	429 (96%)	17 (4%)	0	100	100
1	D	441/469 (94%)	426 (97%)	15 (3%)	0	100	100
1	E	447/469 (95%)	433 (97%)	14 (3%)	0	100	100
1	F	443/469 (94%)	428 (97%)	15 (3%)	0	100	100
All	All	2669/2814 (95%)	2581 (97%)	88 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/398 (92%)	360 (99%)	5 (1%)	69	65
1	B	361/398 (91%)	355 (98%)	6 (2%)	63	58
1	C	367/398 (92%)	363 (99%)	4 (1%)	76	73
1	D	362/398 (91%)	357 (99%)	5 (1%)	69	65
1	E	366/398 (92%)	363 (99%)	3 (1%)	83	82
1	F	363/398 (91%)	359 (99%)	4 (1%)	76	73
All	All	2184/2388 (92%)	2157 (99%)	27 (1%)	74	70

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	321	PHE
1	D	95	TYR
1	F	291	ASN
1	C	388	ASN
1	A	434	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 35 ligands modelled in this entry, 18 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	G4S	A	502	2,4	16,16,16	1.35	1 (6%)	19,24,24	0.66	0
4	9RN	A	503	3	11,11,12	0.53	0	15,16,18	1.35	3 (20%)
7	EDO	A	507	-	3,3,3	0.21	0	2,2,2	0.29	0
3	G4S	B	503	2,4	16,16,16	0.87	1 (6%)	19,24,24	0.76	1 (5%)
4	9RN	B	504	3	11,11,12	0.50	0	15,16,18	1.25	2 (13%)
7	EDO	C	503	-	3,3,3	0.30	0	2,2,2	0.61	0
3	G4S	C	504	2,4	16,16,16	1.35	1 (6%)	19,24,24	0.68	0
4	9RN	C	505	3	11,11,12	0.50	0	15,16,18	1.26	3 (20%)
7	EDO	D	505	-	3,3,3	0.20	0	2,2,2	0.19	0
3	G4S	D	506	2,4	16,16,16	0.90	1 (6%)	19,24,24	0.77	0
4	9RN	D	507	3	11,11,12	0.47	0	15,16,18	1.20	2 (13%)
7	EDO	E	505	-	3,3,3	0.30	0	2,2,2	0.12	0
3	G4S	E	506	2,4	16,16,16	0.90	1 (6%)	19,24,24	0.78	0
4	9RN	E	507	3	11,11,12	0.46	0	15,16,18	1.37	2 (13%)
7	EDO	F	503	-	3,3,3	0.13	0	2,2,2	0.42	0
3	G4S	F	504	2,4	16,16,16	1.31	1 (6%)	19,24,24	0.64	0
4	9RN	F	505	3	11,11,12	0.49	0	15,16,18	1.27	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G4S	A	502	2,4	-	0/7/27/27	0/1/1/1
4	9RN	A	503	3	-	-	0/3/2/2
7	EDO	A	507	-	-	0/1/1/1	-
3	G4S	B	503	2,4	-	0/7/27/27	0/1/1/1
4	9RN	B	504	3	-	-	0/3/2/2
7	EDO	C	503	-	-	0/1/1/1	-
3	G4S	C	504	2,4	-	0/7/27/27	0/1/1/1
4	9RN	C	505	3	-	-	0/3/2/2
7	EDO	D	505	-	-	0/1/1/1	-
3	G4S	D	506	2,4	-	0/7/27/27	0/1/1/1
4	9RN	D	507	3	-	-	0/3/2/2

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	E	505	-	-	0/1/1/1	-
3	G4S	E	506	2,4	-	0/7/27/27	0/1/1/1
4	9RN	E	507	3	-	-	0/3/2/2
7	EDO	F	503	-	-	0/1/1/1	-
3	G4S	F	504	2,4	-	0/7/27/27	0/1/1/1
4	9RN	F	505	3	-	-	0/3/2/2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	G4S	O7-S	4.41	1.62	1.44
3	C	504	G4S	O8-S	4.38	1.62	1.44
3	F	504	G4S	O7-S	4.34	1.62	1.44
3	D	506	G4S	O9-S	2.20	1.62	1.49
3	E	506	G4S	O9-S	2.12	1.62	1.49

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	507	9RN	O5-C5-C6	-3.27	108.47	113.33
4	D	507	9RN	C2-C3-C4	-3.12	106.12	113.85
4	F	505	9RN	C2-C3-C4	-3.09	106.21	113.85
4	A	503	9RN	O5-C5-C6	-3.01	108.87	113.33
4	B	504	9RN	O5-C5-C6	-2.98	108.91	113.33

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand G4S A 502




## Ligand G4S B 503




## Ligand G4S C 504




## Ligand G4S D 506



## Ligand G4S E 506





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	447/469 (95%)	-0.40	1 (0%) 94 97	11, 19, 32, 47	0
1	B	447/469 (95%)	-0.32	1 (0%) 94 97	11, 21, 36, 47	0
1	C	448/469 (95%)	-0.41	4 (0%) 84 89	11, 18, 32, 61	0
1	D	444/469 (94%)	-0.34	2 (0%) 90 94	13, 21, 38, 57	0
1	E	448/469 (95%)	-0.38	2 (0%) 92 95	13, 20, 34, 48	0
1	F	447/469 (95%)	-0.44	0 100 100	11, 17, 29, 56	0
All	All	2681/2814 (95%)	-0.38	10 (0%) 92 95	11, 19, 35, 61	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	50	THR	5.0
1	C	183	HIS	3.9
1	E	26	ALA	3.7
1	C	26	ALA	2.9
1	C	180	GLY	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	D	501	1/1	0.90	0.15	37,37,37,37	0
7	EDO	F	503	4/4	0.93	0.11	24,28,28,30	0
7	EDO	D	505	4/4	0.94	0.15	27,29,30,31	0
7	EDO	C	503	4/4	0.95	0.14	19,23,23,23	0
4	9RN	C	505	10/11	0.95	0.08	15,16,16,17	0
7	EDO	E	505	4/4	0.95	0.14	24,28,28,31	0
4	9RN	E	507	10/11	0.96	0.07	15,17,18,20	0
4	9RN	D	507	10/11	0.96	0.09	18,19,20,20	0
2	CA	A	501	1/1	0.96	0.13	26,26,26,26	1
4	9RN	B	504	10/11	0.96	0.09	16,16,17,18	0
3	G4S	A	502	16/16	0.97	0.10	16,17,20,23	0
4	9RN	A	503	10/11	0.97	0.08	15,17,17,18	0
3	G4S	C	504	16/16	0.97	0.08	15,17,20,20	0
7	EDO	A	507	4/4	0.97	0.10	23,25,25,26	0
3	G4S	E	506	16/16	0.97	0.08	16,18,22,22	0
4	9RN	F	505	10/11	0.98	0.07	14,15,16,17	0
3	G4S	B	503	16/16	0.98	0.07	17,19,22,23	0
5	CL	A	504	1/1	0.98	0.06	20,20,20,20	0
3	G4S	F	504	16/16	0.98	0.08	13,15,17,17	0
2	CA	F	501	1/1	0.98	0.06	27,27,27,27	0
3	G4S	D	506	16/16	0.98	0.07	20,23,25,27	0
5	CL	F	502	1/1	0.99	0.07	15,15,15,15	0
2	CA	B	501	1/1	0.99	0.16	27,27,27,27	1
2	CA	E	501	1/1	0.99	0.10	24,24,24,24	1
2	CA	C	501	1/1	0.99	0.10	28,28,28,28	1
6	ZN	A	506	1/1	1.00	0.04	23,23,23,23	0
6	ZN	A	505	1/1	1.00	0.05	22,22,22,22	0
6	ZN	D	504	1/1	1.00	0.04	22,22,22,22	0
5	CL	C	502	1/1	1.00	0.05	17,17,17,17	0
6	ZN	E	504	1/1	1.00	0.05	21,21,21,21	0
5	CL	D	502	1/1	1.00	0.06	22,22,22,22	0
5	CL	E	502	1/1	1.00	0.05	17,17,17,17	0
6	ZN	E	503	1/1	1.00	0.04	22,22,22,22	0
5	CL	B	502	1/1	1.00	0.06	21,21,21,21	0
6	ZN	D	503	1/1	1.00	0.06	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around G4S A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around G4S C 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around G4S E 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around G4S B 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around G4S F 504:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around G4S D 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.