



# Full wwPDB X-ray Structure Validation Report ⓘ

May 5, 2020 – 01:43 PM EDT

PDB ID : 4PVB  
Title : Crystal structure of Aminopeptidase N in complex with the phosphonic acid analogue of leucine (D-(S)-LeuP)  
Authors : Nocek, B.; Vassiliou, S.; Berlicki, L.; Mulligan, R.; Mucha, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2014-03-16  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.10.1  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.10.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	867	

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 7259 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	864	Total	C	N	O	S	Se	0	1	0
			6864	4352	1186	1303	5	18			

- Molecule 2 is [(1S)-1-amino-3-methylbutyl]phosphonic acid (three-letter code: 2WW) (formula: C<sub>5</sub>H<sub>14</sub>NO<sub>3</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			10	5	1	3	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	351	Total	O	0	3
			354	354		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Aminopeptidase N



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	224.26 Å   224.26 Å   57.94 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	32.00 – 2.10 32.37 – 2.10	Depositor EDS
% Data completeness (in resolution range)	85.9 (32.00-2.10) 85.9 (32.37-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.46 (at 2.10 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.177 , 0.233 0.191 , 0.243	Depositor DCC
$R_{free}$ test set	2753 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7259	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN, PO4, 2WW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.86	1/7001 (0.0%)	0.89	4/9463 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	711	GLU	CD-OE1	5.18	1.31	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	52	ASP	CB-CG-OD1	-5.96	112.93	118.30
1	A	277	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	85	ARG	NE-CZ-NH2	-5.53	117.54	120.30
1	A	85	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6864	0	6716	97	0
2	A	10	0	13	2	0
3	A	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	25	0	0	0	0
5	A	5	0	0	0	0
6	A	354	0	0	12	0
All	All	7259	0	6729	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:453:ASN:HD21	1:A:520:THR:HG23	1.38	0.88
1:A:655:ASN:OD1	1:A:679:ARG:CD	2.28	0.82
1:A:508:THR:H	1:A:512:GLN:HE22	1.28	0.82
1:A:655:ASN:OD1	1:A:679:ARG:HD2	1.82	0.80
1:A:453:ASN:ND2	1:A:520:THR:HG23	1.98	0.77
1:A:322:ARG:HG2	1:A:322:ARG:HH11	1.50	0.76
1:A:438:GLN:HE22	1:A:473:GLN:H	1.34	0.74
1:A:322:ARG:HG2	1:A:322:ARG:NH1	2.03	0.73
1:A:31:GLN:H	1:A:31:GLN:NE2	1.89	0.70
1:A:255:ASN:O	1:A:781:ARG:NH2	2.24	0.69
1:A:31:GLN:H	1:A:31:GLN:HE21	1.40	0.69
1:A:553:ASP:O	1:A:559:ARG:HD2	1.94	0.67
1:A:459:VAL:CG1	1:A:507:LEU:HD22	2.26	0.65
1:A:450:LEU:HD12	1:A:454:ILE:O	1.97	0.64
1:A:62:ILE:HD11	1:A:78:ILE:HD13	1.78	0.64
1:A:174:ASP:OD2	1:A:178:LYS:NZ	2.31	0.62
1:A:386:MSE:HE1	1:A:435:TRP:CD2	2.35	0.61
1:A:459:VAL:HG12	1:A:507:LEU:HD22	1.80	0.61
1:A:483:LEU:HD23	1:A:525:PRO:HA	1.82	0.60
1:A:508:THR:H	1:A:512:GLN:NE2	1.98	0.60
1:A:386:MSE:HE1	1:A:435:TRP:CE2	2.36	0.60
1:A:203:ARG:NH2	6:A:1411:HOH:O	2.20	0.60
1:A:256:MSE:HG3	2:A:1001:2WW:H6	1.85	0.58
1:A:588:GLU:HG3	6:A:1101:HOH:O	2.02	0.58
1:A:322:ARG:CG	1:A:322:ARG:HH11	2.17	0.57
1:A:347:HIS:HB2	1:A:374:MSE:HE2	1.87	0.57
1:A:588:GLU:CG	6:A:1101:HOH:O	2.53	0.57
1:A:374:MSE:CE	1:A:378:GLU:HG3	2.37	0.55
1:A:31:GLN:N	1:A:31:GLN:HE21	2.04	0.55
1:A:781:ARG:NE	6:A:1403:HOH:O	2.41	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:738:SER:O	1:A:771:LYS:HE3	2.10	0.51
1:A:115:GLN:OE1	2:A:1001:2WW:H3	2.11	0.51
1:A:427:ILE:HG12	1:A:428:ASN:N	2.24	0.50
1:A:100:LYS:NZ	6:A:1377:HOH:O	2.43	0.50
1:A:115:GLN:NE2	6:A:1129:HOH:O	2.43	0.50
1:A:205:VAL:HG11	1:A:230:MSE:HE1	1.94	0.50
1:A:661:GLN:HG3	1:A:675:TRP:CH2	2.47	0.50
1:A:16:TYR:CD1	1:A:42:PRO:HA	2.46	0.49
1:A:605:ASN:HD22	1:A:606:ALA:N	2.11	0.49
1:A:481:VAL:HG12	1:A:527:LEU:HD23	1.95	0.49
1:A:115:GLN:NE2	1:A:115:GLN:HA	2.28	0.49
1:A:697:VAL:HG22	1:A:704:MSE:HE1	1.95	0.49
1:A:148:TYR:O	1:A:168:HIS:HE1	1.96	0.49
1:A:831[A]:ASN:ND2	6:A:1393:HOH:O	2.46	0.48
1:A:605:ASN:HD22	1:A:606:ALA:H	1.60	0.48
1:A:655:ASN:OD1	1:A:679:ARG:HD3	2.09	0.48
1:A:375:THR:O	1:A:379:LYS:HB3	2.14	0.47
1:A:459:VAL:HG11	1:A:507:LEU:HD22	1.95	0.47
1:A:661:GLN:HB3	1:A:675:TRP:CE2	2.49	0.47
1:A:45:VAL:CG1	1:A:83:SER:OG	2.63	0.47
1:A:309:TRP:HB3	1:A:368:MSE:SE	2.65	0.47
1:A:483:LEU:HD23	1:A:525:PRO:CA	2.45	0.47
1:A:62:ILE:HG12	1:A:67:ALA:HB2	1.97	0.46
1:A:797:GLU:O	1:A:836:HIS:HE1	1.97	0.46
1:A:119:GLU:O	1:A:122:ARG:HG2	2.15	0.46
1:A:206:LYS:HD3	6:A:1380:HOH:O	2.16	0.46
1:A:182:LEU:O	1:A:261:ASN:HB3	2.16	0.46
1:A:509:GLU:HG3	1:A:512:GLN:HG3	1.98	0.46
1:A:553:ASP:OD1	1:A:554:SER:N	2.49	0.46
1:A:860:GLU:O	1:A:864:LYS:HG2	2.16	0.46
1:A:509:GLU:HG2	1:A:512:GLN:NE2	2.31	0.45
1:A:630:LEU:HD13	1:A:837:ARG:HH21	1.82	0.45
1:A:53:GLY:HA3	1:A:126:PHE:CD2	2.52	0.45
1:A:338:ILE:HD11	1:A:557:PHE:HA	1.97	0.45
1:A:648:LEU:HB3	1:A:649:PRO:HD3	1.99	0.45
1:A:52:ASP:O	1:A:126:PHE:HA	2.18	0.44
1:A:250:ALA:HB2	1:A:272:VAL:HG11	2.00	0.44
1:A:509:GLU:CG	1:A:512:GLN:HE21	2.30	0.44
1:A:664:GLN:HA	1:A:666:TYR:CZ	2.53	0.44
1:A:842:LYS:NZ	6:A:1393:HOH:O	2.51	0.44
1:A:374:MSE:HE1	1:A:378:GLU:HG3	2.01	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:GLY:O	1:A:114:THR:HA	2.18	0.43
1:A:230:MSE:HG2	1:A:295:TYR:CZ	2.53	0.43
1:A:60:VAL:CG1	1:A:69:TYR:CE2	3.02	0.42
1:A:755:ARG:HG3	6:A:1341:HOH:O	2.18	0.42
1:A:459:VAL:HG12	1:A:507:LEU:CD2	2.48	0.42
1:A:588:GLU:HG2	6:A:1101:HOH:O	2.19	0.42
1:A:366:GLU:HB2	1:A:860:GLU:CD	2.40	0.42
1:A:198:THR:OG1	1:A:204:ASN:ND2	2.53	0.42
1:A:531:PHE:CZ	1:A:534:PRO:HA	2.54	0.41
1:A:598:ILE:HG23	1:A:654:LEU:HD22	2.02	0.41
1:A:767:LEU:HD21	1:A:784:ILE:HD11	2.02	0.41
1:A:265:ASN:HD22	1:A:265:ASN:N	2.18	0.41
1:A:509:GLU:CG	1:A:512:GLN:NE2	2.83	0.41
1:A:9:LYS:NZ	6:A:1257:HOH:O	2.53	0.41
1:A:178:LYS:HB2	1:A:179:PRO:HD2	2.01	0.41
1:A:444:LEU:HD22	1:A:535:VAL:HG21	2.01	0.41
1:A:509:GLU:H	1:A:512:GLN:HE21	1.69	0.41
1:A:62:ILE:HD11	1:A:78:ILE:CD1	2.45	0.41
1:A:653:GLU:O	1:A:657:GLN:HG3	2.21	0.41
1:A:349:PHE:HB2	1:A:350:PRO:HD3	2.03	0.41
1:A:539:TYR:CD2	1:A:540:PRO:HD2	2.56	0.41
1:A:314:LEU:HD21	1:A:412:VAL:HG23	2.02	0.41
1:A:100:LYS:HB2	1:A:815:PHE:HB2	2.03	0.41
1:A:726:ARG:C	1:A:726:ARG:HD3	2.41	0.41
1:A:38:LEU:O	1:A:87:THR:HA	2.22	0.40
1:A:586:LYS:O	1:A:587:HIS:C	2.59	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	863/867 (100%)	838 (97%)	24 (3%)	1 (0%)	53	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	830	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	730/715 (102%)	704 (96%)	26 (4%)	38	39

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	41	GLU
1	A	83	SER
1	A	84	GLU
1	A	164	SER
1	A	177	SER
1	A	219	VAL
1	A	261	ASN
1	A	270	LYS
1	A	289	SER
1	A	322	ARG
1	A	323	ASP
1	A	352	ASP
1	A	462	THR
1	A	496	GLN
1	A	499	ARG
1	A	509	GLU
1	A	517	GLU
1	A	520	THR
1	A	596	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	605	ASN
1	A	679	ARG
1	A	755	ARG
1	A	822	ARG
1	A	856	LYS
1	A	857	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	115	GLN
1	A	168	HIS
1	A	204	ASN
1	A	261	ASN
1	A	265	ASN
1	A	346	GLN
1	A	347	HIS
1	A	438	GLN
1	A	453	ASN
1	A	512	GLN
1	A	605	ASN
1	A	732	GLN
1	A	818	GLN
1	A	836	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	A	1006	-	4,4,4	0.47	0	6,6,6	0.38	0
4	SO4	A	1003	-	4,4,4	0.22	0	6,6,6	0.75	0
4	SO4	A	1007	-	4,4,4	0.57	0	6,6,6	0.43	0
4	SO4	A	1004	-	4,4,4	0.37	0	6,6,6	0.73	0
2	2WW	A	1001	3	8,9,9	4.15	3 (37%)	8,13,13	2.69	3 (37%)
4	SO4	A	1005	-	4,4,4	0.31	0	6,6,6	0.51	0
5	PO4	A	1008	-	4,4,4	0.70	0	6,6,6	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2WW	A	1001	3	-	6/9/10/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	2WW	P11-C3	9.87	1.93	1.84
2	A	1001	2WW	P11-O11	-4.96	1.47	1.54
2	A	1001	2WW	P11-O12	-3.37	1.49	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	2WW	C1-C3-N12	5.07	122.15	110.17
2	A	1001	2WW	O12-P11-O11	3.85	118.00	107.64
2	A	1001	2WW	O12-P11-O13	3.22	121.54	113.45

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1001	2WW	C1-C3-P11-O11
2	A	1001	2WW	C1-C3-P11-O12
2	A	1001	2WW	C1-C3-P11-O13
2	A	1001	2WW	C2-C1-C3-N12
2	A	1001	2WW	C3-C1-C2-C5
2	A	1001	2WW	C3-C1-C2-C4

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	2WW	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	846/867 (97%)	-0.24	14 (1%) 70 74	18, 30, 56, 78	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	581	GLY	3.7
1	A	852	GLU	3.4
1	A	451	LYS	3.3
1	A	182	LEU	3.3
1	A	500	ALA	2.8
1	A	452	ASN	2.5
1	A	582	VAL	2.5
1	A	181	TYR	2.4
1	A	583	GLU	2.4
1	A	453	ASN	2.2
1	A	584	LEU	2.1
1	A	586	LYS	2.1
1	A	578	LEU	2.0
1	A	449	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	A	1002	1/1	0.93	0.05	34,34,34,34	1
4	SO4	A	1007	5/5	0.93	0.23	59,60,65,68	0
4	SO4	A	1006	5/5	0.94	0.21	61,65,73,74	0
4	SO4	A	1005	5/5	0.94	0.26	59,61,67,70	0
2	2WW	A	1001	10/10	0.96	0.14	27,30,34,39	0
4	SO4	A	1004	5/5	0.97	0.08	53,53,60,71	0
5	PO4	A	1008	5/5	0.97	0.06	40,47,55,58	0
4	SO4	A	1003	5/5	0.99	0.08	32,36,39,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.