



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 19, 2019 – 10:34 AM EDT

PDB ID : 2Q04  
Title : Crystal structure of acetoin utilization protein (ZP\_00540088.1) from *Exiguobacterium sibiricum* 255-15 at 2.33 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-05-18  
Resolution : 2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.3.2  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.3.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1763 (2.36-2.32)
Clashscore	122126	1858 (2.36-2.32)
Ramachandran outliers	120053	1834 (2.36-2.32)
Sidechain outliers	120020	1835 (2.36-2.32)
RSRZ outliers	108989	1737 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>32%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>
1	B	211	<div> <div>27%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>•</div> </div> </div>
1	C	211	<div> <div>35%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>••</div> </div> </div>
1	D	211	<div> <div>19%</div> <div> <div></div> <div>89%</div> <div>10%</div> </div> </div>
1	E	211	<div> <div>16%</div> <div> <div></div> <div>89%</div> <div>10%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	211	<div><div>35%</div><div>89%</div><div>6% . .</div></div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACY	E	212	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetoin utilization protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	Se	0	2	0
			1691	1092	284	307	1	7			
1	B	207	Total	C	N	O	S	Se	0	2	0
			1645	1058	282	298	1	6			
1	C	208	Total	C	N	O	S	Se	0	2	0
			1608	1030	272	297	1	8			
1	D	210	Total	C	N	O	S	Se	0	1	0
			1678	1083	280	306	1	8			
1	E	210	Total	C	N	O	S	Se	0	0	0
			1660	1072	277	302	1	8			
1	F	204	Total	C	N	O	S	Se	0	1	0
			1560	997	267	288	1	7			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q41BL0
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
A	120	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
A	125	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
A	155	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
A	158	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
A	159	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
A	183	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
A	208	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
B	0	GLY	-	LEADER SEQUENCE	UNP Q41BL0
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
B	120	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
B	125	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
B	155	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
B	158	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
B	159	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
B	183	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	208	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	0	GLY	-	LEADER SEQUENCE	UNP Q41BL0
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	120	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	125	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	155	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	158	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	159	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	183	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
C	208	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	0	GLY	-	LEADER SEQUENCE	UNP Q41BL0
D	1	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	120	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	125	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	155	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	158	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	159	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	183	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
D	208	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	0	GLY	-	LEADER SEQUENCE	UNP Q41BL0
E	1	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	120	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	125	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	155	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	158	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	159	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	183	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
E	208	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	0	GLY	-	LEADER SEQUENCE	UNP Q41BL0
F	1	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	120	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	125	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	155	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	158	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	159	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	183	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0
F	208	MSE	MET	MODIFIED RESIDUE	UNP Q41BL0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
-----	-------	----------	-------	---------	---------

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total 1 Ca 1	0	0
2	E	1	Total 1 Ca 1	0	0
2	B	1	Total 1 Ca 1	0	0
2	C	1	Total 1 Ca 1	0	0
2	A	2	Total 2 Ca 2	0	0
2	F	1	Total 1 Ca 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total 4 C 2 O 2	0	0
3	A	1	Total 4 C 2 O 2	0	0
3	A	1	Total 4 C 2 O 2	0	0
3	C	1	Total 4 C 2 O 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	57	Total	O	0	0
			57	57		
5	B	45	Total	O	0	0
			45	45		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	23	Total 23	O 23	0	0
5	D	57	Total 57	O 57	0	0
5	E	40	Total 40	O 40	0	0
5	F	18	Total 18	O 18	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetoin utilization protein




- Molecule 1: Acetoin utilization protein




- Molecule 1: Acetoin utilization protein





- Molecule 1: Acetoin utilization protein







• Molecule 1: Acetoin utilization protein



• Molecule 1: Acetoin utilization protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.87Å 128.58Å 108.43Å 90.00° 108.43° 90.00°	Depositor
Resolution (Å)	29.34 – 2.33 29.33 – 2.33	Depositor EDS
% Data completeness (in resolution range)	93.7 (29.34-2.33) 93.7 (29.33-2.33)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.2.0005, PHENIX	Depositor
R, $R_{free}$	0.229 , 0.260 0.235 , 0.267	Depositor DCC
$R_{free}$ test set	3696 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.1	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 51.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	10133	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.81 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0849e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACY, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/1734	0.42	0/2344
1	B	0.61	0/1689	0.42	0/2286
1	C	0.53	0/1650	0.42	0/2235
1	D	0.74	1/1718 (0.1%)	0.44	0/2319
1	E	0.65	0/1698	0.44	0/2298
1	F	0.54	0/1598	0.43	0/2168
All	All	0.62	1/10087 (0.0%)	0.43	0/13650

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	158	MSE	SE-CE	5.33	2.26	1.95

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1691	0	1622	15	0
1	B	1645	0	1586	12	0
1	C	1608	0	1486	14	0
1	D	1678	0	1633	18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1660	0	1597	15	0
1	F	1560	0	1440	11	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	12	0	18	1	0
3	C	4	0	6	1	0
3	D	8	0	12	0	0
3	F	4	0	6	0	0
4	A	4	0	3	1	0
4	E	8	0	6	2	0
4	F	4	0	3	0	0
5	A	57	0	0	0	0
5	B	45	0	0	0	0
5	C	23	0	0	0	0
5	D	57	0	0	1	0
5	E	40	0	0	1	0
5	F	18	0	0	0	0
All	All	10133	0	9418	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:125:MSE:SE	1:D:125:MSE:CE	2.15	1.44
1:D:158:MSE:SE	1:D:158:MSE:CE	2.26	1.33
1:E:155:MSE:SE	1:E:183:MSE:HE1	2.20	0.92
1:B:157:LYS:NZ	1:F:150:ASP:OD2	2.12	0.82
1:A:178:PRO:HG2	1:D:149:TRP:CH2	2.22	0.75

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/211 (100%)	206 (98%)	5 (2%)	0	100	100
1	B	207/211 (98%)	201 (97%)	4 (2%)	2 (1%)	17	16
1	C	208/211 (99%)	196 (94%)	10 (5%)	2 (1%)	17	16
1	D	209/211 (99%)	202 (97%)	7 (3%)	0	100	100
1	E	208/211 (99%)	205 (99%)	3 (1%)	0	100	100
1	F	203/211 (96%)	198 (98%)	5 (2%)	0	100	100
All	All	1246/1266 (98%)	1208 (97%)	34 (3%)	4 (0%)	43	49

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	207	PHE
1	C	35	ASP
1	C	36	PRO
1	B	205	ARG

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/170 (101%)	171 (99%)	1 (1%)	87	93
1	B	167/170 (98%)	167 (100%)	0	100	100
1	C	154/170 (91%)	151 (98%)	3 (2%)	60	72

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	173/170 (102%)	172 (99%)	1 (1%)	87	93
1	E	169/170 (99%)	167 (99%)	2 (1%)	74	84
1	F	150/170 (88%)	146 (97%)	4 (3%)	48	58
All	All	985/1020 (97%)	974 (99%)	11 (1%)	76	85

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	115	LEU
1	E	109	GLN
1	F	20	ILE
1	C	204	ARG
1	F	13	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	110	GLN
1	C	160	ASN
1	D	160	ASN
1	E	160	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 7 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	A	213	-	3,3,3	0.53	0	2,2,2	0.28	0
3	EDO	A	214	-	3,3,3	0.48	0	2,2,2	0.27	0
3	EDO	A	215	-	3,3,3	0.65	0	2,2,2	0.22	0
4	ACY	A	216	-	1,3,3	1.23	0	0,3,3	0.00	-
3	EDO	C	212	-	3,3,3	0.54	0	2,2,2	0.19	0
3	EDO	D	212	-	3,3,3	0.43	0	2,2,2	0.25	0
3	EDO	D	213	-	3,3,3	0.53	0	2,2,2	0.36	0
4	ACY	E	212	-	1,3,3	0.61	0	0,3,3	0.00	-
4	ACY	E	213	2	1,3,3	2.29	1 (100%)	0,3,3	0.00	-
3	EDO	F	212	-	3,3,3	0.49	0	2,2,2	0.31	0
4	ACY	F	213	-	1,3,3	1.28	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	213	-	-	0/1/1/1	-
3	EDO	A	214	-	-	1/1/1/1	-
3	EDO	A	215	-	-	0/1/1/1	-
3	EDO	C	212	-	-	0/1/1/1	-
3	EDO	D	212	-	-	0/1/1/1	-
3	EDO	D	213	-	-	1/1/1/1	-
3	EDO	F	212	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	213	ACY	CH3-C	2.29	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	F	212	EDO	O1-C1-C2-O2
3	D	213	EDO	O1-C1-C2-O2
3	A	214	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	214	EDO	1	0
3	C	212	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	203/211 (96%)	1.58	67 (33%) 0 0	52, 57, 67, 82	0
1	B	200/211 (94%)	1.58	57 (28%) 0 0	51, 57, 69, 76	0
1	C	201/211 (95%)	1.70	74 (36%) 0 0	50, 57, 66, 83	0
1	D	202/211 (95%)	1.20	40 (19%) 1 2	47, 57, 71, 90	0
1	E	202/211 (95%)	1.03	33 (16%) 1 3	48, 57, 67, 85	0
1	F	197/211 (93%)	1.87	74 (37%) 0 0	51, 57, 64, 85	0
All	All	1205/1266 (95%)	1.49	345 (28%) 0 0	47, 57, 69, 90	0

The worst 5 of 345 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	146	LEU	8.7
1	C	40	ALA	7.7
1	F	69	ASN	7.2
1	F	27	GLN	7.2
1	F	56	ALA	7.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	214	4/4	0.78	0.34	76,79,80,81	0
3	EDO	A	213	4/4	0.83	0.29	82,83,83,83	0
4	ACY	F	213	4/4	0.84	0.34	88,88,88,89	0
2	CA	C	211	1/1	0.87	0.10	73,73,73,73	0
4	ACY	E	213	4/4	0.89	0.21	56,56,57,58	0
3	EDO	C	212	4/4	0.89	0.22	62,65,67,68	0
4	ACY	A	216	4/4	0.91	0.22	79,79,80,80	0
3	EDO	D	213	4/4	0.91	0.19	44,45,49,53	0
2	CA	E	211	1/1	0.94	0.10	51,51,51,51	0
4	ACY	E	212	4/4	0.95	0.14	48,49,49,50	0
3	EDO	F	212	4/4	0.95	0.07	69,71,73,75	0
3	EDO	D	212	4/4	0.96	0.10	37,40,43,48	0
2	CA	F	211	1/1	0.96	0.17	68,68,68,68	0
3	EDO	A	215	4/4	0.97	0.10	38,48,48,55	0
2	CA	A	212	1/1	0.97	0.04	68,68,68,68	0
2	CA	D	211	1/1	0.98	0.04	39,39,39,39	0
2	CA	A	211	1/1	0.98	0.04	48,48,48,48	0
2	CA	B	211	1/1	0.99	0.03	48,48,48,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.