



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 09:10 pm GMT

PDB ID : 3QA0
Title : Crystal structure of the apo-form of human CK2 alpha at pH 6.5
Authors : Battistutta, R.; Ranchio, A.; Papinutto, E.
Deposited on : 2011-01-10
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4155 (2.50-2.50)
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>16%</div> <div>••</div> </div> </div>
1	B	336	<div> <div>4%</div> <div> <div></div> <div>79%</div> <div>15%</div> <div>•••</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	0	2	0
			2805	1795	495	504	11			
1	B	330	Total	C	N	O	S	0	2	0
			2805	1795	495	504	11			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	119	Total	O	0	0
			119	119		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	110	Total 110	O 110	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Casein kinase II subunit alpha



• Molecule 1: Casein kinase II subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	127.84Å 127.84Å 124.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.17 – 2.50 57.17 – 2.50	Depositor EDS
% Data completeness (in resolution range)	87.2 (57.17-2.50) 87.2 (57.17-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.207 , 0.249 0.205 , 0.245	Depositor DCC
R_{free} test set	1576 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	44.5	Xtriage
Anisotropy	0.428	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for -h,-l,-k 0.009 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5894	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.71 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0135e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2880	0.73	1/3895 (0.0%)
1	B	0.67	0/2880	0.76	4/3895 (0.1%)
All	All	0.66	0/5760	0.75	5/7790 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	280	ARG	NE-CZ-NH1	-5.59	117.51	120.30
1	B	105	VAL	N-CA-CB	5.58	123.78	111.50
1	B	107	ARG	N-CA-C	5.33	125.40	111.00
1	A	128	LEU	CA-CB-CG	5.04	126.90	115.30
1	B	195	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2805	0	2747	43	1
1	B	2805	0	2747	69	1
2	A	15	0	0	0	0
2	B	20	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	12	0	18	6	0
3	B	8	0	12	1	0
4	A	119	0	0	14	0
4	B	110	0	0	10	0
All	All	5894	0	5524	112	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:ASP:HB3	1:B:163:MET:CE	1.74	1.15
1:B:120:ASP:HB3	1:B:163:MET:HE1	1.28	1.14
1:A:278:ARG:H	3:A:341:EDO:H22	1.23	1.03
1:A:169:ARG:HD3	4:A:396:HOH:O	1.67	0.95
1:B:104:PRO:O	1:B:105:VAL:HG22	1.68	0.94
1:B:124:LEU:HA	1:B:125:TYR:CB	1.98	0.94
1:B:124:LEU:HA	1:B:125:TYR:HB3	1.47	0.93
1:A:310:GLN:HG2	4:A:402:HOH:O	1.71	0.90
1:B:124:LEU:CA	1:B:125:TYR:HB3	2.02	0.89
1:B:123:GLN:O	1:B:125:TYR:HB3	1.75	0.87
1:A:278:ARG:N	3:A:341:EDO:H22	1.88	0.87
1:A:8:ARG:HD2	4:A:368:HOH:O	1.77	0.84
1:B:124:LEU:HD22	1:B:124:LEU:N	1.94	0.82
1:A:118:ASN:HB3	4:A:435:HOH:O	1.80	0.80
1:B:62:ASN:HB2	4:B:422:HOH:O	1.81	0.80
1:A:89:ARG:HD2	4:A:393:HOH:O	1.85	0.75
1:B:326:THR:HG21	4:B:428:HOH:O	1.87	0.74
1:A:21:ARG:NH1	4:A:460:HOH:O	2.20	0.74
1:A:68:LYS:HE2	3:A:340:EDO:C1	2.18	0.74
1:B:124:LEU:H	1:B:124:LEU:HD22	1.53	0.72
1:B:96:THR:HB	1:B:114:GLU:HG2	1.69	0.72
1:B:104:PRO:O	1:B:105:VAL:CG2	2.37	0.71
1:A:326:THR:HG22	4:A:412:HOH:O	1.92	0.69
1:B:120:ASP:HB3	1:B:163:MET:HE2	1.69	0.69
1:B:123:GLN:C	1:B:125:TYR:HB3	2.12	0.69
1:B:299:ASP:OD2	1:B:303:LYS:HE2	1.95	0.66
1:B:120:ASP:CB	1:B:163:MET:CE	2.64	0.66
1:B:244:ARG:HD3	4:B:387:HOH:O	1.95	0.66
1:B:123:GLN:O	1:B:125:TYR:CB	2.43	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:ASN:C	1:B:58:ASN:HD22	1.99	0.65
1:B:120:ASP:CB	1:B:163:MET:HE1	2.17	0.64
1:B:89:ARG:HD2	4:B:395:HOH:O	1.98	0.64
1:B:310:GLN:HG2	4:B:365:HOH:O	1.96	0.63
1:A:58:ASN:C	1:A:58:ASN:HD22	2.02	0.63
1:B:283:ARG:NH1	4:B:347:HOH:O	2.32	0.62
1:A:299:ASP:OD2	1:A:303:LYS:HE2	2.02	0.60
1:A:278:ARG:H	3:A:341:EDO:C2	2.08	0.59
1:A:89:ARG:HB3	4:A:393:HOH:O	2.01	0.59
1:B:124:LEU:HD23	1:B:124:LEU:O	2.03	0.59
1:B:121:PHE:O	1:B:125:TYR:CD1	2.55	0.59
1:B:107:ARG:HG3	1:B:108:THR:N	2.17	0.59
1:B:124:LEU:HB2	1:B:126:GLN:HG3	1.84	0.59
1:B:124:LEU:CB	1:B:126:GLN:HG3	2.33	0.58
1:B:73:VAL:HG23	4:B:391:HOH:O	2.04	0.57
1:B:121:PHE:HD2	1:B:162:VAL:HG11	1.70	0.56
1:B:36:GLN:OE1	3:B:341:EDO:H11	2.05	0.56
1:A:68:LYS:HE2	3:A:340:EDO:H12	1.88	0.55
1:B:120:ASP:CB	1:B:163:MET:HE2	2.35	0.54
1:B:168:HIS:CD2	4:B:344:HOH:O	2.59	0.54
1:B:36:GLN:OE1	1:B:104:PRO:HD3	2.08	0.54
1:B:124:LEU:HA	1:B:125:TYR:CD2	2.42	0.54
1:A:170:LYS:NZ	4:A:418:HOH:O	2.41	0.54
1:B:104:PRO:C	1:B:106:SER:H	2.10	0.53
1:B:64:LYS:O	1:B:115:HIS:HB2	2.08	0.53
1:A:326:THR:CG2	4:A:412:HOH:O	2.52	0.53
1:A:260:LYS:NZ	1:A:308:ASP:OD2	2.38	0.52
1:A:134:ARG:HG2	1:A:323:TYR:CZ	2.45	0.51
1:B:73:VAL:HG21	1:B:77:LYS:HE3	1.92	0.51
1:A:89:ARG:HD3	4:B:385:HOH:O	2.10	0.51
1:A:244:ARG:HD3	4:A:400:HOH:O	2.11	0.50
1:A:134:ARG:HD3	1:A:323:TYR:O	2.12	0.50
1:A:86:GLU:HG2	1:A:89:ARG:NH2	2.27	0.49
1:B:122:LYS:HG3	1:B:159:PRO:HB2	1.94	0.48
1:A:290:GLN:HG2	4:A:427:HOH:O	2.12	0.48
1:B:121:PHE:O	1:B:125:TYR:HD1	1.96	0.48
1:B:10:ARG:HD2	1:B:310:GLN:O	2.13	0.48
1:A:196:TYR:CD2	1:A:231:PRO:HG3	2.49	0.47
1:A:10:ARG:HD2	1:A:310:GLN:O	2.14	0.47
1:B:124:LEU:H	1:B:124:LEU:HD13	1.79	0.47
1:B:86:GLU:HG2	1:B:89:ARG:NH2	2.29	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:TYR:CG	1:B:125:TYR:O	2.67	0.47
1:B:124:LEU:HA	1:B:125:TYR:CG	2.50	0.46
1:A:102:LYS:CE	4:A:446:HOH:O	2.62	0.46
1:B:124:LEU:N	1:B:125:TYR:HB3	2.29	0.46
1:B:124:LEU:O	1:B:124:LEU:CD2	2.64	0.46
1:B:165:ASP:OD2	1:B:168:HIS:HD2	1.98	0.46
1:A:136:TYR:O	1:A:140:ILE:HG13	2.14	0.46
1:A:35:ASN:O	1:A:38:ASP:HB2	2.16	0.46
1:A:25:ASP:OD1	3:A:342:EDO:O2	2.27	0.45
1:B:96:THR:HB	1:B:114:GLU:CG	2.44	0.45
1:B:257:TYR:CE1	1:B:308:ASP:HA	2.52	0.44
1:B:119:THR:HG23	1:B:164:ILE:O	2.18	0.44
1:B:321:HIS:CG	1:B:322:PRO:HD2	2.53	0.44
1:B:64:LYS:HG2	1:B:115:HIS:CD2	2.53	0.44
1:A:165:ASP:OD2	1:A:168:HIS:HD2	2.00	0.44
1:A:124:LEU:C	1:A:126:GLN:H	2.20	0.43
1:B:171:LEU:O	1:B:172:ARG:NH1	2.47	0.43
1:B:125:TYR:C	1:B:125:TYR:CD2	2.92	0.43
1:B:121:PHE:CD2	1:B:162:VAL:HG11	2.51	0.43
1:A:95:ILE:HB	1:A:174:ILE:HG22	2.00	0.43
1:B:58:ASN:C	1:B:58:ASN:ND2	2.71	0.43
1:A:73:VAL:HG21	1:A:77:LYS:HE3	2.00	0.42
1:B:124:LEU:CB	1:B:126:GLN:HE21	2.32	0.42
1:B:134:ARG:HG2	1:B:323:TYR:CZ	2.54	0.42
1:B:103:ASP:OD2	1:B:106:SER:HB2	2.19	0.42
1:B:49:LYS:HE3	1:B:49:LYS:HB2	1.75	0.42
1:B:77:LYS:HD3	1:B:80:ARG:NH2	2.35	0.42
1:B:105:VAL:C	1:B:107:ARG:H	2.23	0.42
1:A:257:TYR:CE1	1:A:308:ASP:HA	2.55	0.42
1:A:77:LYS:HD3	1:A:80:ARG:NH2	2.35	0.42
1:B:95:ILE:HB	1:B:174:ILE:HG22	2.01	0.42
1:B:316:ARG:HA	1:B:316:ARG:HD2	1.85	0.41
1:A:102:LYS:HE2	4:A:446:HOH:O	2.18	0.41
1:A:124:LEU:C	1:A:126:GLN:N	2.73	0.41
1:A:237:ASP:C	1:A:237:ASP:OD1	2.58	0.41
1:A:322:PRO:HA	1:A:325:TYR:CD1	2.56	0.41
1:B:107:ARG:HG3	1:B:108:THR:HG23	2.01	0.41
1:A:132:ASP:OD1	1:A:169:ARG:NH1	2.53	0.41
1:B:120:ASP:HA	1:B:163:MET:HA	2.03	0.41
1:A:196:TYR:HA	1:A:241:GLN:NE2	2.36	0.41
1:B:117:ASN:HB2	4:B:438:HOH:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:LYS:O	1:A:110:ALA:HA	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:TYR:OH	1:B:32:GLU:OE2[2_565]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/336 (98%)	314 (95%)	16 (5%)	0	100	100
1	B	330/336 (98%)	311 (94%)	18 (6%)	1 (0%)	43	64
All	All	660/672 (98%)	625 (95%)	34 (5%)	1 (0%)	49	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	105	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/308 (99%)	291 (95%)	15 (5%)	27	50
1	B	306/308 (99%)	280 (92%)	26 (8%)	12	22
All	All	612/616 (99%)	571 (93%)	41 (7%)	18	34

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	40	GLN
1	A	49	LYS
1	A	58	ASN
1	A	89	ARG
1	A	118	ASN
1	A	119	THR
1	A	122	LYS
1	A	258	ILE
1	A	259	ASP
1	A	265	LEU
1	A	271	ASP
1	A	273	LEU
1	A	288	GLU
1	A	330	ASP
1	B	2	SER
1	B	21	ARG
1	B	32	GLU
1	B	36	GLN
1	B	40	GLN
1	B	49	LYS
1	B	58	ASN
1	B	77	LYS
1	B	102	LYS
1	B	106	SER
1	B	107	ARG
1	B	108	THR
1	B	114	GLU
1	B	119	THR
1	B	120	ASP
1	B	121	PHE
1	B	122	LYS
1	B	124	LEU
1	B	125	TYR
1	B	153	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	258	ILE
1	B	259	ASP
1	B	265	LEU
1	B	271	ASP
1	B	273	LEU
1	B	288	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	ASN
1	A	118	ASN
1	A	168	HIS
1	A	331	GLN
1	B	58	ASN
1	B	115	HIS
1	B	126	GLN
1	B	168	HIS
1	B	234	HIS
1	B	331	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	337	-	4,4,4	0.25	0	6,6,6	0.48	0
2	SO4	A	338	-	4,4,4	0.17	0	6,6,6	0.44	0
2	SO4	A	339	-	4,4,4	0.21	0	6,6,6	0.18	0
3	EDO	A	340	-	3,3,3	0.48	0	2,2,2	0.52	0
3	EDO	A	341	-	3,3,3	0.63	0	2,2,2	0.34	0
3	EDO	A	342	-	3,3,3	0.34	0	2,2,2	1.03	0
2	SO4	B	337	-	4,4,4	0.20	0	6,6,6	0.38	0
2	SO4	B	338	-	4,4,4	0.26	0	6,6,6	0.47	0
2	SO4	B	339	-	4,4,4	0.22	0	6,6,6	0.15	0
2	SO4	B	340	-	4,4,4	0.24	0	6,6,6	0.47	0
3	EDO	B	341	-	3,3,3	0.48	0	2,2,2	0.45	0
3	EDO	B	342	-	3,3,3	0.51	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	337	-	-	0/0/0/0	0/0/0/0
2	SO4	A	338	-	-	0/0/0/0	0/0/0/0
2	SO4	A	339	-	-	0/0/0/0	0/0/0/0
3	EDO	A	340	-	-	0/1/1/1	0/0/0/0
3	EDO	A	341	-	-	0/1/1/1	0/0/0/0
3	EDO	A	342	-	-	0/1/1/1	0/0/0/0
2	SO4	B	337	-	-	0/0/0/0	0/0/0/0
2	SO4	B	338	-	-	0/0/0/0	0/0/0/0
2	SO4	B	339	-	-	0/0/0/0	0/0/0/0
2	SO4	B	340	-	-	0/0/0/0	0/0/0/0
3	EDO	B	341	-	-	0/1/1/1	0/0/0/0
3	EDO	B	342	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	340	EDO	2	0
3	A	341	EDO	3	0
3	A	342	EDO	1	0
3	B	341	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/336 (98%)	-0.02	6 (1%) 68 71	25, 40, 74, 90	0
1	B	330/336 (98%)	0.09	15 (4%) 33 36	22, 41, 88, 148	0
All	All	660/672 (98%)	0.04	21 (3%) 47 51	22, 40, 77, 148	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	121	PHE	7.4
1	B	125	TYR	4.5
1	B	118	ASN	4.5
1	B	104	PRO	4.3
1	A	329	LYS	3.4
1	A	120	ASP	3.2
1	B	329	LYS	2.8
1	B	271	ASP	2.8
1	B	116	VAL	2.8
1	B	325	TYR	2.7
1	A	271	ASP	2.5
1	A	2	SER	2.5
1	B	120	ASP	2.4
1	B	115	HIS	2.3
1	A	128	LEU	2.3
1	B	330	ASP	2.2
1	B	107	ARG	2.2
1	B	131	TYR	2.1
1	B	105	VAL	2.1
1	A	275	ARG	2.0
1	B	160	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	B	341	4/4	0.77	0.24	51,53,56,59	0
3	EDO	A	340	4/4	0.88	0.18	56,57,57,58	0
3	EDO	A	341	4/4	0.90	0.15	41,45,47,50	0
3	EDO	A	342	4/4	0.92	0.28	42,43,46,48	0
3	EDO	B	342	4/4	0.93	0.20	47,50,53,54	0
2	SO4	A	339	5/5	0.93	0.17	92,93,93,94	0
2	SO4	A	337	5/5	0.94	0.16	63,64,65,65	0
2	SO4	B	340	5/5	0.94	0.13	69,69,69,70	0
2	SO4	B	338	5/5	0.94	0.15	68,68,70,70	0
2	SO4	A	338	5/5	0.95	0.15	80,80,80,81	0
2	SO4	B	339	5/5	0.95	0.12	87,87,87,88	0
2	SO4	B	337	5/5	0.97	0.15	43,44,44,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.