



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 07:05 pm GMT

PDB ID : 3QB7  
Title : Interleukin-4 mutant RGA bound to cytokine receptor common gamma  
Authors : Bates, D.L.; Junttila, I.S.; Creusot, R.J.; Moraga, I.; Lupardus, P.; Fathman, C.G.; Paul, W.E.; Garcia, K.C.  
Deposited on : 2011-01-12  
Resolution : 3.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1379 (3.28-3.20)
Clashscore	122126	1510 (3.28-3.20)
Ramachandran outliers	120053	1485 (3.28-3.20)
Sidechain outliers	120020	1484 (3.28-3.20)
RSRZ outliers	108989	1331 (3.28-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	<div> <div>%</div> <div> <div></div> <div>58%</div> <div>35%</div> <div>7%</div> </div> </div>
1	B	132	<div> <div></div> <div>69%</div> <div>25%</div> <div>• 5%</div> </div>
2	C	203	<div> <div>%</div> <div> <div></div> <div>58%</div> <div>36%</div> <div>6%</div> </div> </div>
2	D	203	<div> <div>4%</div> <div> <div></div> <div>55%</div> <div>35%</div> <div>10%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5337 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	123	Total	C	N	O	S	0	1	0
			1015	639	186	183	7			
1	B	125	Total	C	N	O	S	0	1	0
			1025	644	187	187	7			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	ALA	-	EXPRESSION TAG	UNP D4HNR6
A	-1	ASP	-	EXPRESSION TAG	UNP D4HNR6
A	0	PRO	-	EXPRESSION TAG	UNP D4HNR6
A	117	ARG	LYS	ENGINEERED MUTATION	UNP D4HNR6
A	118	VAL	THR	ENGINEERED MUTATION	UNP D4HNR6
A	121	GLN	ARG	ENGINEERED MUTATION	UNP D4HNR6
A	122	SER	GLU	ENGINEERED MUTATION	UNP D4HNR6
A	124	TRP	TYR	ENGINEERED MUTATION	UNP D4HNR6
A	125	PHE	SER	ENGINEERED MUTATION	UNP D4HNR6
A	128	GLY	SER	ENGINEERED MUTATION	UNP D4HNR6
A	129	ALA	SER	ENGINEERED MUTATION	UNP D4HNR6
B	-2	ALA	-	EXPRESSION TAG	UNP D4HNR6
B	-1	ASP	-	EXPRESSION TAG	UNP D4HNR6
B	0	PRO	-	EXPRESSION TAG	UNP D4HNR6
B	117	ARG	LYS	ENGINEERED MUTATION	UNP D4HNR6
B	118	VAL	THR	ENGINEERED MUTATION	UNP D4HNR6
B	121	GLN	ARG	ENGINEERED MUTATION	UNP D4HNR6
B	122	SER	GLU	ENGINEERED MUTATION	UNP D4HNR6
B	124	TRP	TYR	ENGINEERED MUTATION	UNP D4HNR6
B	125	PHE	SER	ENGINEERED MUTATION	UNP D4HNR6
B	128	GLY	SER	ENGINEERED MUTATION	UNP D4HNR6
B	129	ALA	SER	ENGINEERED MUTATION	UNP D4HNR6


- Molecule 2 is a protein called Cytokine receptor common subunit gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	191	Total	C	N	O	S	5	1	0
			1635	1042	293	292	8			
2	D	183	Total	C	N	O	S	0	0	0
			1568	1001	280	279	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	30	ALA	-	EXPRESSION TAG	UNP P31785
C	31	ASP	-	EXPRESSION TAG	UNP P31785
C	32	PRO	-	EXPRESSION TAG	UNP P31785
C	53	GLN	ASN	ENGINEERED MUTATION	UNP P31785
D	30	ALA	-	EXPRESSION TAG	UNP P31785
D	31	ASP	-	EXPRESSION TAG	UNP P31785
D	32	PRO	-	EXPRESSION TAG	UNP P31785
D	53	GLN	ASN	ENGINEERED MUTATION	UNP P31785

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin 4



- Molecule 1: Interleukin 4




- Molecule 2: Cytokine receptor common subunit gamma



- Molecule 2: Cytokine receptor common subunit gamma





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.97Å 105.88Å 121.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.51 – 3.25 48.51 – 3.25	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.51-3.25) 99.4 (48.51-3.25)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 3.25Å)	Xtriage
Refinement program	PHENIX dev_613	Depositor
R, $R_{free}$	0.231 , 0.292 0.232 , 0.287	Depositor DCC
$R_{free}$ test set	799 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.0	Xtriage
Anisotropy	0.385	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 68.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	5337	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.23	0/1033	0.40	0/1387
1	B	0.24	0/1044	0.42	0/1405
2	C	0.23	0/1692	0.45	1/2301 (0.0%)
2	D	0.23	0/1617	0.44	0/2194
All	All	0.23	0/5386	0.43	1/7287 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	206	ASN	C-N-CD	-6.11	107.16	120.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1015	0	1032	36	0
1	B	1025	0	1037	31	0
2	C	1635	0	1545	62	0
2	D	1568	0	1471	50	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	42	0	38	2	0
4	D	42	0	38	5	0
All	All	5337	0	5161	175	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 175 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:205:PHE:CZ	2:C:209:CYS:HB2	1.98	0.98
1:A:69:THR:HB	4:D:233:NAG:H82	1.47	0.95
1:B:22:THR:HG22	1:B:25:THR:HG23	1.49	0.91
2:C:188:LEU:HD12	2:C:189:PRO:HD2	1.60	0.82
2:C:131:ILE:HD11	2:C:214:HIS:HB3	1.61	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/132 (91%)	118 (98%)	2 (2%)	0	100	100
1	B	124/132 (94%)	124 (100%)	0	0	100	100
2	C	188/203 (93%)	179 (95%)	9 (5%)	0	100	100
2	D	173/203 (85%)	168 (97%)	5 (3%)	0	100	100
All	All	605/670 (90%)	589 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/118 (97%)	113 (99%)	1 (1%)	81	91
1	B	115/118 (98%)	114 (99%)	1 (1%)	81	91
2	C	185/195 (95%)	183 (99%)	2 (1%)	76	89
2	D	176/195 (90%)	176 (100%)	0	100	100
All	All	590/626 (94%)	586 (99%)	4 (1%)	85	93

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	CYS
1	B	56	TYR
2	C	93	CYS
2	C	171	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	72	GLN
2	C	66	HIS
2	C	206	ASN
1	B	59	HIS
2	C	154	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	130	-	4,4,4	0.17	0	6,6,6	0.08	0
3	SO4	B	130	-	4,4,4	0.17	0	6,6,6	0.08	0
4	NAG	C	233	2,4	14,14,15	0.93	1 (7%)	17,19,21	5.46	5 (29%)
4	NAG	C	234	4	14,14,15	0.45	0	17,19,21	1.03	1 (5%)
4	NAG	C	300	2	14,14,15	0.50	0	17,19,21	0.88	1 (5%)
4	NAG	D	233	2,4	14,14,15	0.77	1 (7%)	17,19,21	5.21	5 (29%)
4	NAG	D	234	4	14,14,15	0.45	0	17,19,21	0.83	1 (5%)
4	NAG	D	300	2	14,14,15	0.53	0	17,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	130	-	-	0/0/0/0	0/0/0/0
3	SO4	B	130	-	-	0/0/0/0	0/0/0/0
4	NAG	C	233	2,4	-	0/6/23/26	0/1/1/1
4	NAG	C	234	4	-	0/6/23/26	0/1/1/1
4	NAG	C	300	2	-	0/6/23/26	0/1/1/1
4	NAG	D	233	2,4	-	0/6/23/26	0/1/1/1
4	NAG	D	234	4	-	0/6/23/26	0/1/1/1
4	NAG	D	300	2	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	233	NAG	O5-C1	-2.70	1.39	1.43
4	D	233	NAG	O5-C1	-2.25	1.40	1.43

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	233	NAG	C1-O5-C5	-2.70	108.47	112.19
4	C	233	NAG	C1-O5-C5	-2.59	108.63	112.19
4	C	233	NAG	C6-C5-C4	-2.36	107.41	112.99
4	D	233	NAG	C2-N2-C7	-2.15	119.81	122.94
4	D	233	NAG	C6-C5-C4	-2.05	108.14	112.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	233	NAG	1	0
4	C	234	NAG	2	0
4	D	233	NAG	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	123/132 (93%)	-0.05	1 (0%) 86 79	47, 83, 135, 161	0
1	B	125/132 (94%)	-0.17	0 100 100	58, 82, 123, 149	0
2	C	191/203 (94%)	0.15	3 (1%) 72 62	48, 85, 143, 179	1 (0%)
2	D	183/203 (90%)	0.40	9 (4%) 29 20	58, 99, 155, 187	0
All	All	622/670 (92%)	0.12	13 (2%) 63 52	47, 87, 145, 187	1 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	64	THR	3.1
2	C	184	HIS	2.5
2	D	65	LEU	2.4
2	C	96	GLN	2.4
2	D	111	GLN	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	C	300	14/15	0.81	0.12	104,113,120,122	0
3	SO4	A	130	5/5	0.84	0.28	130,131,135,137	0
4	NAG	D	234	14/15	0.85	0.22	107,128,141,142	0
4	NAG	D	300	14/15	0.87	0.13	110,114,121,122	0
4	NAG	C	234	14/15	0.88	0.18	90,101,117,120	0
4	NAG	C	233	14/15	0.94	0.17	88,98,107,110	0
3	SO4	B	130	5/5	0.95	0.20	96,97,101,103	0
4	NAG	D	233	14/15	0.96	0.16	53,65,87,87	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.