



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 30, 2019 – 02:09 PM EST

PDB ID : 6QB5
Title : Crystal structure of the N-terminal region of human cohesin subunit STAG1
Authors : Newman, J.A.; Katis, V.L.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.; Bountra, C.; Gileadi, O.
Deposited on : 2018-12-20
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	rb-20031633
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20031633


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	9172 (2.04-2.00)
Clashscore	122126	10355 (2.04-2.00)
Ramachandran outliers	120053	10237 (2.04-2.00)
Sidechain outliers	120020	10236 (2.04-2.00)
RSRZ outliers	108989	8961 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	339	<div> <div>%</div> <div>85% 11%</div> </div>
1	B	339	<div> <div>%</div> <div>87% 6% 7%</div> </div>
1	C	339	<div> <div>%</div> <div>86% 6% 7%</div> </div>
1	D	339	<div> <div>%</div> <div>84% 6% 10%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10628 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cohesin subunit SA-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	315	Total	C	N	O	S	0	1	0
			2571	1637	434	480	20			
1	A	303	Total	C	N	O	S	0	0	0
			2414	1540	404	450	20			
1	B	316	Total	C	N	O	S	0	0	0
			2549	1623	432	474	20			
1	D	305	Total	C	N	O	S	0	0	0
			2466	1569	416	461	20			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	82	SER	-	expression tag	UNP Q8WVM7
C	83	MET	-	expression tag	UNP Q8WVM7
C	84	GLY	-	expression tag	UNP Q8WVM7
C	85	GLY	-	expression tag	UNP Q8WVM7
A	82	SER	-	expression tag	UNP Q8WVM7
A	83	MET	-	expression tag	UNP Q8WVM7
A	84	GLY	-	expression tag	UNP Q8WVM7
A	85	GLY	-	expression tag	UNP Q8WVM7
B	82	SER	-	expression tag	UNP Q8WVM7
B	83	MET	-	expression tag	UNP Q8WVM7
B	84	GLY	-	expression tag	UNP Q8WVM7
B	85	GLY	-	expression tag	UNP Q8WVM7
D	82	SER	-	expression tag	UNP Q8WVM7
D	83	MET	-	expression tag	UNP Q8WVM7
D	84	GLY	-	expression tag	UNP Q8WVM7
D	85	GLY	-	expression tag	UNP Q8WVM7

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Na 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	169	Total 169	O 169	0	0
3	A	152	Total 152	O 152	0	0
3	B	148	Total 148	O 148	0	0
3	D	158	Total 158	O 158	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.61Å 124.98Å 231.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.00 – 2.02 85.00 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.5 (85.00-2.02) 99.7 (85.00-2.02)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.02Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.218 , 0.245 0.218 , 0.245	Depositor DCC
R_{free} test set	6239 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.644	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10628	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/2455	0.36	0/3313
1	B	0.23	0/2591	0.35	0/3492
1	C	0.23	0/2615	0.35	0/3523
1	D	0.23	0/2507	0.35	0/3381
All	All	0.23	0/10168	0.35	0/13709

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2414	0	2369	5	0
1	B	2549	0	2522	8	0
1	C	2571	0	2550	11	0
1	D	2466	0	2433	7	0
2	D	1	0	0	0	0
3	A	152	0	0	0	0
3	B	148	0	0	0	0
3	C	169	0	0	1	0
3	D	158	0	0	1	0
All	All	10628	0	9874	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:300:TYR:OH	1:D:301:ARG:NH2	2.31	0.63
1:A:121:ILE:HG12	1:A:206:LEU:HD22	1.81	0.62
1:C:227:MET:HB3	1:C:314:GLU:HG2	1.83	0.61
1:D:227:MET:HB3	1:D:314:GLU:HG2	1.83	0.60
1:B:227:MET:HB3	1:B:314:GLU:HG2	1.83	0.59
1:C:135:ARG:NH1	1:C:137:GLU:OE2	2.38	0.57
1:A:227:MET:HB3	1:A:314:GLU:HG2	1.87	0.57
1:C:114:ASP:OD2	1:C:140:ARG:NH1	2.41	0.53
1:A:330:SER:O	1:A:333:LYS:NZ	2.39	0.52
1:B:121:ILE:HD11	1:B:202:THR:HG22	1.92	0.52
1:A:394:ILE:O	1:A:398:THR:HG23	2.11	0.51
1:C:312:ILE:HG13	1:C:338:THR:HG21	1.92	0.51
1:D:146:GLU:HG3	1:D:149:ARG:HH22	1.78	0.49
1:C:149:ARG:NH2	3:C:503:HOH:O	2.35	0.49
1:C:121:ILE:HG12	1:C:206:LEU:HD22	1.95	0.48
1:C:93:LEU:HD21	1:C:95:LYS:HE3	1.95	0.48
1:D:158:ASP:OD1	1:D:216:ARG:NE	2.47	0.47
1:B:109:TYR:CZ	1:B:113:ARG:HG2	2.51	0.46
1:D:251:TYR:HA	1:D:270:LEU:HD13	1.97	0.45
1:C:184:VAL:HG22	1:C:187:ARG:HH22	1.81	0.45
1:D:349:LYS:NZ	3:D:602:HOH:O	2.32	0.45
1:C:334:TYR:O	1:C:338:THR:OG1	2.31	0.43
1:C:230:MET:HG2	1:C:318:TRP:CE2	2.55	0.42
1:B:263:ALA:HB3	1:B:266:ARG:HB3	2.02	0.42
1:D:315:ILE:HA	1:D:318:TRP:CE3	2.55	0.42
1:B:98:MET:HA	1:B:101:VAL:HG12	2.02	0.41
1:B:170:GLN:HG2	1:B:171:TRP:CD1	2.56	0.41
1:B:361:ARG:HA	1:B:364:PHE:CD1	2.56	0.41
1:C:351:LEU:HD11	1:C:381:MET:HE2	2.03	0.41
1:B:230:MET:HG2	1:B:318:TRP:CZ2	2.56	0.40
1:A:315:ILE:HA	1:A:318:TRP:CE3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/339 (88%)	293 (98%)	6 (2%)	0	100	100
1	B	312/339 (92%)	307 (98%)	5 (2%)	0	100	100
1	C	312/339 (92%)	307 (98%)	5 (2%)	0	100	100
1	D	301/339 (89%)	296 (98%)	4 (1%)	1 (0%)	43	38
All	All	1224/1356 (90%)	1203 (98%)	20 (2%)	1 (0%)	53	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	192	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/306 (85%)	256 (98%)	4 (2%)	67	71
1	B	275/306 (90%)	270 (98%)	5 (2%)	62	64
1	C	281/306 (92%)	279 (99%)	2 (1%)	85	89
1	D	269/306 (88%)	263 (98%)	6 (2%)	55	56
All	All	1085/1224 (89%)	1068 (98%)	17 (2%)	65	68

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	172	LYS
1	C	322	TYR
1	A	99	GLN
1	A	137	GLU
1	A	214	GLN
1	A	322	TYR
1	B	161	ASP
1	B	201	ASP
1	B	249	ARG
1	B	269	LEU
1	B	322	TYR
1	D	137	GLU
1	D	268	GLU
1	D	269	LEU
1	D	272	GLN
1	D	322	TYR
1	D	364	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/339 (89%)	-0.07	4 (1%) 77 77	28, 43, 80, 94	0
1	B	316/339 (93%)	-0.04	2 (0%) 89 89	28, 47, 74, 86	0
1	C	315/339 (92%)	0.06	5 (1%) 72 71	28, 42, 74, 100	0
1	D	305/339 (89%)	0.06	4 (1%) 77 77	29, 43, 81, 104	0
All	All	1239/1356 (91%)	0.00	15 (1%) 79 78	28, 44, 77, 104	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	169	PRO	5.3
1	C	170	GLN	3.8
1	C	158	ASP	3.4
1	D	267	LEU	3.3
1	D	269	LEU	3.0
1	A	169	PRO	2.7
1	A	246	ASN	2.7
1	D	270	LEU	2.7
1	C	168	GLY	2.6
1	B	401	LEU	2.6
1	A	162	TYR	2.5
1	A	364	PHE	2.5
1	C	337[A]	TRP	2.1
1	D	248	GLN	2.1
1	B	169	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NA	D	501	1/1	0.91	0.12	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.