



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2018 – 03:41 pm GMT

PDB ID : 1QWJ
Title : The Crystal Structure of Murine CMP-5-N-Acetylneuraminic Acid Synthetase
Authors : Krapp, S.; Muenster-Kuehnel, A.K.; Kaiser, J.T.; Huber, R.; Tiralongo, J.; Gerardy-Schahn, R.; Jacob, U.
Deposited on : 2003-09-02
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2792 (2.80-2.80)
Clashscore	122126	3209 (2.80-2.80)
Ramachandran outliers	120053	3158 (2.80-2.80)
Sidechain outliers	120020	3160 (2.80-2.80)
RSRZ outliers	108989	2726 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	<div> <div>53%</div> <div>42%</div> <div>.</div> </div>
1	B	229	<div> <div>59%</div> <div>35%</div> <div>6%</div> </div>
1	C	229	<div> <div>%</div> <div>51%</div> <div>42%</div> <div>7%</div> </div>
1	D	229	<div> <div>10%</div> <div>50%</div> <div>45%</div> <div>5%</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 7611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cytidine monophospho-N-acetylneuraminic acid synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	37	0	0
			1831	1160	329	337	5			
1	B	229	Total	C	N	O	S	22	0	0
			1840	1165	330	340	5			
1	C	228	Total	C	N	O	S	118	0	0
			1831	1160	329	337	5			
1	D	228	Total	C	N	O	S	130	0	0
			1831	1160	329	337	5			

- Molecule 2 is CYTIDINE-5'-MONOPHOSPHATE-5-N-ACETYLNEURAMINIC ACID (three-letter code: NCC) (formula: C₂₀H₃₁N₄O₁₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			41	20	4	16	1		
2	C	1	Total	C	N	O	P	0	0
			41	20	4	16	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	P	0	0
			41	20	4	16	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	52	Total	O	0	0
			52	52		
3	B	59	Total	O	0	0
			59	59		
3	C	26	Total	O	0	0
			26	26		
3	D	18	Total	O	0	0
			18	18		

● Molecule 1: cytidine monophospho-N-acetylneuraminic acid synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.05Å 79.94Å 170.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 42.56 – 2.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.80) 99.4 (42.56-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.69Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.240 , 0.304 0.232 , 0.352	Depositor DCC
R_{free} test set	2429 reflections (8.52%)	wwPDB-VP
Wilson B-factor (Å ²)	46.1	Xtriage
Anisotropy	0.143	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 78.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	7611	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NCC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/1875	0.63	0/2537
1	B	0.42	0/1884	0.68	1/2549 (0.0%)
1	C	0.36	0/1875	0.58	0/2537
1	D	0.36	0/1875	0.58	0/2537
All	All	0.39	0/7509	0.62	1/10160 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	206	ASP	N-CA-C	-5.50	96.16	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	186	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1831	0	1794	90	0
1	B	1840	0	1800	79	0
1	C	1831	0	1794	91	0
1	D	1831	0	1794	101	0
2	A	41	0	29	2	0
2	C	41	0	29	0	0
2	D	41	0	29	0	0
3	A	52	0	0	5	0
3	B	59	0	0	9	0
3	C	26	0	0	2	0
3	D	18	0	0	0	0
All	All	7611	0	7269	347	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 347 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:HIS:HD2	1:B:94:GLU:H	1.12	0.98
1:C:228:LEU:HD23	1:C:228:LEU:H	1.31	0.93
1:D:177:ARG:H	1:D:193:ASN:ND2	1.67	0.93
1:D:43:LEU:H	1:D:84:GLN:HE21	1.11	0.92
1:B:92:HIS:CD2	1:B:94:GLU:H	1.93	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/229 (99%)	208 (92%)	14 (6%)	4 (2%)	9	30
1	B	227/229 (99%)	200 (88%)	18 (8%)	9 (4%)	3	10
1	C	226/229 (99%)	194 (86%)	23 (10%)	9 (4%)	3	10
1	D	226/229 (99%)	189 (84%)	27 (12%)	10 (4%)	3	9
All	All	905/916 (99%)	791 (87%)	82 (9%)	32 (4%)	4	13

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	54	LYS
1	A	185	VAL
1	A	207	GLY
1	B	179	SER
1	B	185	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/195 (100%)	179 (92%)	15 (8%)	14	38
1	B	195/195 (100%)	179 (92%)	16 (8%)	12	35
1	C	194/195 (100%)	169 (87%)	25 (13%)	5	14
1	D	194/195 (100%)	179 (92%)	15 (8%)	14	38
All	All	777/780 (100%)	706 (91%)	71 (9%)	10	30

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	46	LEU
1	C	122	ASP
1	D	193	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	56	ILE
1	C	113	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	148	HIS
1	C	42	HIS
1	D	175	GLN
1	B	153	GLN
1	B	195	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NCC	A	2002	-	36,43,43	1.90	7 (19%)	41,65,65	1.19	3 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NCC	C	3002	-	36,43,43	1.71	8 (22%)	41,65,65	1.09	2 (4%)
2	NCC	D	4002	-	36,43,43	1.80	8 (22%)	41,65,65	1.11	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NCC	A	2002	-	-	0/23/70/70	0/3/3/3
2	NCC	C	3002	-	-	0/23/70/70	0/3/3/3
2	NCC	D	4002	-	-	0/23/70/70	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3002	NCC	C19-C16	2.04	1.55	1.53
2	A	2002	NCC	C6-C5	2.12	1.42	1.38
2	A	2002	NCC	O4'-C1'	2.16	1.44	1.41
2	D	4002	NCC	O4'-C1'	2.20	1.44	1.41
2	D	4002	NCC	C6-N1	2.25	1.38	1.35

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4002	NCC	C16-N5-C17	-4.13	112.95	123.23
2	C	3002	NCC	C16-N5-C17	-3.94	113.43	123.23
2	A	2002	NCC	C16-N5-C17	-3.54	114.44	123.23
2	A	2002	NCC	N4-C4-N3	2.03	119.88	116.56
2	D	4002	NCC	N4-C4-N3	2.06	119.92	116.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2002	NCC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/229 (99%)	-0.46	1 (0%) 92 90	15, 32, 58, 95	8 (3%)
1	B	227/229 (99%)	-0.48	0 100 100	11, 30, 62, 112	3 (1%)
1	C	218/229 (95%)	-0.13	3 (1%) 75 69	20, 49, 93, 111	12 (5%)
1	D	219/229 (95%)	0.37	23 (10%) 6 3	25, 58, 104, 121	16 (7%)
All	All	891/916 (97%)	-0.18	27 (3%) 50 40	11, 41, 91, 121	39 (4%)

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	51	GLY	5.1
1	C	95	ILE	4.4
1	D	234	ALA	4.4
1	D	120	SER	4.2
1	D	166	SER	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NCC	D	4002	41/41	0.79	0.29	64,73,92,93	41
2	NCC	C	3002	41/41	0.84	0.26	63,73,82,86	41
2	NCC	A	2002	41/41	0.96	0.17	15,29,38,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.