



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 16, 2019 – 06:26 AM EDT

PDB ID : 2QWZ
Title : CRYSTAL STRUCTURE OF A PUTATIVE THIOESTERASE
(TM1040_1390) FROM SILICIBACTER SP. TM1040 AT 2.15 Å RES-
OLUTION
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2007-08-10
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.3.2
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.3.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1287 (2.16-2.16)
Clashscore	122126	1390 (2.16-2.16)
Ramachandran outliers	120053	1368 (2.16-2.16)
Sidechain outliers	120020	1367 (2.16-2.16)
RSRZ outliers	108989	1262 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	<div> <div>3%</div> <div>79%</div> <div>11%</div> <div>9%</div> </div>
1	B	159	<div> <div>3%</div> <div>77%</div> <div>13%</div> <div>9%</div> </div>
1	C	159	<div> <div>12%</div> <div>72%</div> <div>14%</div> <div>13%</div> </div>
1	D	159	<div> <div>14%</div> <div>68%</div> <div>17%</div> <div>14%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	141	-	-	X	-
2	ACT	B	141	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylacetic acid degradation-related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	144	Total	C	N	O	Se	0	0	0
			1068	683	178	201	6			
1	B	144	Total	C	N	O	Se	0	0	0
			1060	677	177	200	6			
1	C	138	Total	C	N	O	Se	0	1	0
			987	629	165	187	6			
1	D	137	Total	C	N	O	Se	0	1	0
			994	635	167	186	6			

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	LEADER SEQUENCE	UNP Q1GGU3
A	-17	GLY	-	LEADER SEQUENCE	UNP Q1GGU3
A	-16	SER	-	LEADER SEQUENCE	UNP Q1GGU3
A	-15	ASP	-	LEADER SEQUENCE	UNP Q1GGU3
A	-14	LYS	-	LEADER SEQUENCE	UNP Q1GGU3
A	-13	ILE	-	LEADER SEQUENCE	UNP Q1GGU3
A	-12	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
A	-11	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
A	-10	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
A	-9	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
A	-8	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
A	-7	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
A	-6	GLU	-	LEADER SEQUENCE	UNP Q1GGU3
A	-5	ASN	-	LEADER SEQUENCE	UNP Q1GGU3
A	-4	LEU	-	LEADER SEQUENCE	UNP Q1GGU3
A	-3	TYR	-	LEADER SEQUENCE	UNP Q1GGU3
A	-2	PHE	-	LEADER SEQUENCE	UNP Q1GGU3
A	-1	GLN	-	LEADER SEQUENCE	UNP Q1GGU3
A	0	GLY	-	LEADER SEQUENCE	UNP Q1GGU3
B	-18	MSE	-	LEADER SEQUENCE	UNP Q1GGU3
B	-17	GLY	-	LEADER SEQUENCE	UNP Q1GGU3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	LEADER SEQUENCE	UNP Q1GGU3
B	-15	ASP	-	LEADER SEQUENCE	UNP Q1GGU3
B	-14	LYS	-	LEADER SEQUENCE	UNP Q1GGU3
B	-13	ILE	-	LEADER SEQUENCE	UNP Q1GGU3
B	-12	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
B	-11	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
B	-10	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
B	-9	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
B	-8	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
B	-7	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
B	-6	GLU	-	LEADER SEQUENCE	UNP Q1GGU3
B	-5	ASN	-	LEADER SEQUENCE	UNP Q1GGU3
B	-4	LEU	-	LEADER SEQUENCE	UNP Q1GGU3
B	-3	TYR	-	LEADER SEQUENCE	UNP Q1GGU3
B	-2	PHE	-	LEADER SEQUENCE	UNP Q1GGU3
B	-1	GLN	-	LEADER SEQUENCE	UNP Q1GGU3
B	0	GLY	-	LEADER SEQUENCE	UNP Q1GGU3
C	-18	MSE	-	LEADER SEQUENCE	UNP Q1GGU3
C	-17	GLY	-	LEADER SEQUENCE	UNP Q1GGU3
C	-16	SER	-	LEADER SEQUENCE	UNP Q1GGU3
C	-15	ASP	-	LEADER SEQUENCE	UNP Q1GGU3
C	-14	LYS	-	LEADER SEQUENCE	UNP Q1GGU3
C	-13	ILE	-	LEADER SEQUENCE	UNP Q1GGU3
C	-12	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
C	-11	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
C	-10	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
C	-9	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
C	-8	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
C	-7	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
C	-6	GLU	-	LEADER SEQUENCE	UNP Q1GGU3
C	-5	ASN	-	LEADER SEQUENCE	UNP Q1GGU3
C	-4	LEU	-	LEADER SEQUENCE	UNP Q1GGU3
C	-3	TYR	-	LEADER SEQUENCE	UNP Q1GGU3
C	-2	PHE	-	LEADER SEQUENCE	UNP Q1GGU3
C	-1	GLN	-	LEADER SEQUENCE	UNP Q1GGU3
C	0	GLY	-	LEADER SEQUENCE	UNP Q1GGU3
D	-18	MSE	-	LEADER SEQUENCE	UNP Q1GGU3
D	-17	GLY	-	LEADER SEQUENCE	UNP Q1GGU3
D	-16	SER	-	LEADER SEQUENCE	UNP Q1GGU3
D	-15	ASP	-	LEADER SEQUENCE	UNP Q1GGU3
D	-14	LYS	-	LEADER SEQUENCE	UNP Q1GGU3
D	-13	ILE	-	LEADER SEQUENCE	UNP Q1GGU3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	-12	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
D	-11	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
D	-10	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
D	-9	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
D	-8	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
D	-7	HIS	-	LEADER SEQUENCE	UNP Q1GGU3
D	-6	GLU	-	LEADER SEQUENCE	UNP Q1GGU3
D	-5	ASN	-	LEADER SEQUENCE	UNP Q1GGU3
D	-4	LEU	-	LEADER SEQUENCE	UNP Q1GGU3
D	-3	TYR	-	LEADER SEQUENCE	UNP Q1GGU3
D	-2	PHE	-	LEADER SEQUENCE	UNP Q1GGU3
D	-1	GLN	-	LEADER SEQUENCE	UNP Q1GGU3
D	0	GLY	-	LEADER SEQUENCE	UNP Q1GGU3

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		
4	B	61	Total	O	0	0
			61	61		
4	C	25	Total	O	0	0
			25	25		
4	D	21	Total	O	0	0
			21	21		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phenylacetic acid degradation-related protein




- Molecule 1: Phenylacetic acid degradation-related protein



- Molecule 1: Phenylacetic acid degradation-related protein



- Molecule 1: Phenylacetic acid degradation-related protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	95.31Å 95.31Å 132.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.23 – 2.15 28.23 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.23-2.15) 100.0 (28.23-2.15)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, R_{free}	0.200 , 0.252 0.204 , 0.249	Depositor DCC
R_{free} test set	1926 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.512	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 60.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4343	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.16 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.0382e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/1081	0.85	2/1456 (0.1%)
1	B	0.82	2/1073 (0.2%)	0.88	0/1446
1	C	0.61	0/1001	0.75	3/1354 (0.2%)
1	D	0.60	0/1009	0.81	2/1363 (0.1%)
All	All	0.70	2/4164 (0.0%)	0.83	7/5619 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	-3	TYR	CD1-CE1	7.55	1.50	1.39
1	B	113	VAL	CB-CG1	5.02	1.63	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	90	ARG	NE-CZ-NH1	8.64	124.62	120.30
1	D	90	ARG	NE-CZ-NH2	-7.57	116.52	120.30
1	C	90	ARG	NE-CZ-NH1	7.50	124.05	120.30
1	C	129	ARG	NE-CZ-NH2	-6.87	116.86	120.30
1	C	90	ARG	NE-CZ-NH2	-6.74	116.93	120.30
1	A	129	ARG	NE-CZ-NH2	-6.43	117.09	120.30
1	A	62	ASP	CB-CG-OD1	5.53	123.28	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1068	0	1059	16	0
1	B	1060	0	1042	19	0
1	C	987	0	963	12	0
1	D	994	0	976	14	0
2	A	4	0	3	4	0
2	B	4	0	3	2	0
2	C	8	0	6	1	0
2	D	4	0	3	0	0
3	A	12	0	16	1	0
3	B	12	0	16	0	0
3	C	6	0	8	1	0
3	D	12	0	16	0	0
4	A	65	0	0	2	0
4	B	61	0	0	3	0
4	C	25	0	0	0	0
4	D	21	0	0	0	0
All	All	4343	0	4111	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:155:HOH:O	1:C:123[A]:MSE:HE3	1.80	0.81
1:C:117:LEU:HD23	1:C:129:ARG:HB3	1.69	0.73
1:D:32:LYS:O	1:D:72:HIS:NE2	2.21	0.73
1:B:17:VAL:HG12	1:B:18:PHE:CE1	2.27	0.70
1:B:81:THR:HG21	1:B:132:MSE:HE1	1.74	0.70
1:C:32:LYS:O	1:C:72:HIS:NE2	2.24	0.68
1:B:103:ARG:HH12	2:C:141:ACT:H1	1.59	0.66
1:D:117:LEU:HD23	1:D:129:ARG:HB3	1.80	0.63
1:A:52:VAL:HG23	4:A:201:HOH:O	1.97	0.63
1:D:32:LYS:O	1:D:72:HIS:CE1	2.52	0.62
1:C:73:LEU:HB3	1:C:77:ALA:HB2	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:61:ALA:HA	1:B:116:ILE:HD13	1.83	0.61
1:B:116:ILE:CG2	1:B:130:SER:HB3	2.32	0.60
1:D:116:ILE:HD13	1:D:118:LEU:HD11	1.85	0.58
1:D:35:ILE:HG22	1:D:102:ALA:HB3	1.88	0.56
1:B:81:THR:CG2	1:B:132:MSE:CE	2.84	0.56
1:A:40:ASN:HB2	4:A:145:HOH:O	2.07	0.54
1:C:35:ILE:HG22	1:C:102:ALA:HB3	1.89	0.54
1:D:52:VAL:HG11	1:D:98:LEU:HD22	1.89	0.53
1:D:17:VAL:HG12	1:D:18:PHE:CD1	2.43	0.53
1:D:3:LEU:HD22	1:D:71:ALA:HB1	1.90	0.53
1:B:81:THR:HG21	1:B:132:MSE:CE	2.39	0.52
1:A:118:LEU:HD12	1:A:118:LEU:N	2.23	0.52
1:B:35:ILE:HG22	1:B:102:ALA:HB3	1.91	0.52
1:C:106:LYS:HZ1	3:C:143:GOL:H12	1.74	0.51
1:B:40:ASN:HB2	4:B:159:HOH:O	2.10	0.51
1:A:99:LEU:HD12	3:A:143:GOL:H32	1.94	0.50
1:D:17:VAL:HG12	1:D:18:PHE:CE1	2.47	0.50
1:A:116:ILE:CG2	1:A:130:SER:HB3	2.42	0.49
1:A:35:ILE:HG22	1:A:102:ALA:HB3	1.95	0.49
1:A:102:ALA:HB2	1:A:116:ILE:HD12	1.94	0.49
1:B:17:VAL:HG12	1:B:18:PHE:CD1	2.49	0.48
1:B:81:THR:CG2	1:B:132:MSE:HE1	2.43	0.48
1:B:90:ARG:HD2	2:B:141:ACT:CH3	2.42	0.48
1:A:81:THR:HG21	1:A:132:MSE:HE1	1.96	0.48
1:B:90:ARG:HD2	2:B:141:ACT:H1	1.95	0.47
1:A:53:SER:HB3	2:A:141:ACT:H1	1.95	0.47
1:A:46:LEU:CD2	2:A:141:ACT:H3	2.44	0.47
1:B:52:VAL:HG23	4:B:168:HOH:O	2.14	0.47
1:B:116:ILE:HG23	1:B:130:SER:HB3	1.97	0.46
1:C:61:ALA:HA	1:C:116:ILE:HD13	1.98	0.46
1:A:53:SER:HB3	2:A:141:ACT:CH3	2.46	0.46
1:A:56:SER:O	1:A:59:ALA:HB3	2.16	0.45
1:C:4:VAL:O	1:C:4:VAL:HG23	2.17	0.45
1:D:93:GLU:HB2	1:D:123[B]:MSE:HE3	1.98	0.45
1:A:81:THR:CG2	1:A:132:MSE:CE	2.95	0.44
1:A:46:LEU:HD21	2:A:141:ACT:H3	2.00	0.44
1:D:24:GLU:HG2	1:D:39:LEU:HD13	2.00	0.43
1:A:118:LEU:N	1:A:118:LEU:CD1	2.82	0.43
1:B:57:MSE:HG2	1:B:98:LEU:HD21	2.00	0.43
1:C:52:VAL:HG13	1:C:56:SER:HB2	2.01	0.43
1:C:32:LYS:O	1:C:72:HIS:CE1	2.72	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ALA:HA	1:A:116:ILE:HD13	2.01	0.42
1:B:10:LEU:HD23	1:B:27:ILE:HD13	2.01	0.42
1:B:100:GLY:HA3	1:B:116:ILE:HD11	2.01	0.42
1:D:73:LEU:HB3	1:D:77:ALA:HB2	2.02	0.42
1:C:57:MSE:HG2	1:C:98:LEU:HD21	2.02	0.42
1:B:81:THR:HG23	1:B:132:MSE:CE	2.50	0.41
1:D:84:ALA:HB1	1:D:132:MSE:CE	2.51	0.41
1:C:39:LEU:C	1:C:39:LEU:HD23	2.40	0.41
1:D:37:MSE:HE3	1:D:63:VAL:HG13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/159 (89%)	136 (96%)	6 (4%)	0	100	100
1	B	142/159 (89%)	138 (97%)	4 (3%)	0	100	100
1	C	137/159 (86%)	133 (97%)	4 (3%)	0	100	100
1	D	136/159 (86%)	129 (95%)	6 (4%)	1 (1%)	24	16
All	All	557/636 (88%)	536 (96%)	20 (4%)	1 (0%)	49	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	22	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/124 (88%)	107 (98%)	2 (2%)	62	66
1	B	107/124 (86%)	106 (99%)	1 (1%)	81	85
1	C	97/124 (78%)	92 (95%)	5 (5%)	25	21
1	D	99/124 (80%)	94 (95%)	5 (5%)	26	21
All	All	412/496 (83%)	399 (97%)	13 (3%)	42	41

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ILE
1	A	129	ARG
1	B	129	ARG
1	C	5	PHE
1	C	40	ASN
1	C	86	LEU
1	C	89	MSE
1	C	129	ARG
1	D	6	ASP
1	D	98	LEU
1	D	111	LEU
1	D	120	SER
1	D	129	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	141	-	1,3,3	2.14	1 (100%)	0,3,3	0.00	-
3	GOL	A	142	-	5,5,5	0.46	0	5,5,5	0.48	0
3	GOL	A	143	-	5,5,5	0.60	0	5,5,5	0.57	0
2	ACT	B	141	-	1,3,3	0.97	0	0,3,3	0.00	-
3	GOL	B	142	-	5,5,5	0.22	0	5,5,5	0.43	0
3	GOL	B	143	-	5,5,5	0.47	0	5,5,5	0.44	0
2	ACT	C	141	-	1,3,3	1.66	0	0,3,3	0.00	-
2	ACT	C	142	-	1,3,3	1.86	0	0,3,3	0.00	-
3	GOL	C	143	-	5,5,5	0.41	0	5,5,5	0.70	0
2	ACT	D	141	-	1,3,3	1.63	0	0,3,3	0.00	-
3	GOL	D	142	-	5,5,5	0.53	0	5,5,5	0.44	0
3	GOL	D	143	-	5,5,5	0.37	0	5,5,5	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	142	-	-	4/4/4/4	-
3	GOL	A	143	-	-	2/4/4/4	-
3	GOL	B	142	-	-	2/4/4/4	-
3	GOL	B	143	-	-	4/4/4/4	-
3	GOL	C	143	-	-	2/4/4/4	-
3	GOL	D	142	-	-	1/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	143	-	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	141	ACT	CH3-C	2.14	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	143	GOL	C1-C2-C3-O3
3	C	143	GOL	C1-C2-C3-O3
3	C	143	GOL	O2-C2-C3-O3
3	B	142	GOL	O1-C1-C2-O2
3	B	142	GOL	O1-C1-C2-C3
3	A	142	GOL	O1-C1-C2-C3
3	B	143	GOL	O1-C1-C2-C3
3	A	142	GOL	O2-C2-C3-O3
3	D	142	GOL	O1-C1-C2-C3
3	A	142	GOL	C1-C2-C3-O3
3	D	143	GOL	C1-C2-C3-O3
3	A	143	GOL	O2-C2-C3-O3
3	B	143	GOL	O1-C1-C2-O2
3	D	143	GOL	O2-C2-C3-O3
3	B	143	GOL	O2-C2-C3-O3
3	A	142	GOL	O1-C1-C2-O2
3	D	143	GOL	O1-C1-C2-O2
3	B	143	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	143	GOL	1	0
3	C	143	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/159 (86%)	0.17	4 (2%) 51 60	20, 34, 69, 73	0
1	B	138/159 (86%)	0.21	5 (3%) 42 50	20, 34, 70, 76	0
1	C	132/159 (83%)	0.79	19 (14%) 2 3	29, 44, 73, 78	0
1	D	132/159 (83%)	0.70	23 (17%) 1 1	28, 42, 73, 81	0
All	All	540/636 (84%)	0.46	51 (9%) 8 12	20, 39, 71, 81	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	48	PRO	7.7
1	D	48	PRO	7.6
1	C	21	ILE	7.1
1	A	138	PRO	6.2
1	D	49	GLY	5.5
1	C	46	LEU	5.4
1	D	46	LEU	5.3
1	B	138	PRO	5.0
1	D	45	HIS	5.0
1	D	138	PRO	4.8
1	B	48	PRO	4.2
1	D	51	THR	4.2
1	D	50	GLY	4.2
1	C	13	TYR	4.1
1	D	47	ARG	4.0
1	C	49	GLY	3.9
1	C	5	PHE	3.9
1	C	50	GLY	3.9
1	D	74	GLY	3.8
1	D	21	ILE	3.7
1	D	13	TYR	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	47	ARG	3.6
1	C	77	ALA	3.5
1	C	45	HIS	3.4
1	D	76	GLU	3.3
1	D	73	LEU	3.2
1	C	4	VAL	3.2
1	D	18	PHE	3.1
1	A	-5	ASN	3.0
1	C	2	GLU	2.9
1	C	17	VAL	2.9
1	D	42	GLN	2.8
1	B	-5	ASN	2.8
1	D	16	GLU	2.8
1	A	76	GLU	2.8
1	D	43	GLU	2.8
1	D	77	ALA	2.7
1	D	19	PRO	2.7
1	C	43	GLU	2.7
1	C	20	GLN	2.6
1	B	137	PRO	2.6
1	A	48	PRO	2.5
1	C	74	GLY	2.5
1	C	44	ARG	2.4
1	C	138	PRO	2.4
1	C	16	GLU	2.3
1	D	4	VAL	2.1
1	D	12	ALA	2.1
1	B	50	GLY	2.1
1	D	17	VAL	2.0
1	D	44	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	C	141	4/4	0.72	0.36	61,61,63,63	0
2	ACT	D	141	4/4	0.81	0.16	66,67,67,67	0
3	GOL	A	143	6/6	0.86	0.18	40,61,63,65	0
2	ACT	C	142	4/4	0.86	0.17	69,70,70,71	0
3	GOL	A	142	6/6	0.87	0.14	41,52,57,65	0
3	GOL	B	142	6/6	0.88	0.14	49,57,66,71	0
3	GOL	D	142	6/6	0.90	0.09	59,61,64,70	0
3	GOL	D	143	6/6	0.91	0.36	42,48,61,62	0
2	ACT	A	141	4/4	0.91	0.17	41,51,53,53	0
2	ACT	B	141	4/4	0.91	0.38	64,65,65,65	0
3	GOL	C	143	6/6	0.93	0.13	49,55,60,73	0
3	GOL	B	143	6/6	0.94	0.14	45,54,66,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.