



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 12:25 pm GMT

PDB ID : 3QW4  
Title : Structure of Leishmania donovani UMP synthase  
Authors : French, J.B.; Ealick, S.E.  
Deposited on : 2011-02-26  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1851 (3.00-3.00)
Clashscore	122126	2167 (3.00-3.00)
Ramachandran outliers	120053	2101 (3.00-3.00)
Sidechain outliers	120020	2104 (3.00-3.00)
RSRZ outliers	108989	1751 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	453	 84% 14% •
1	C	453	 82% 15% ••

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UMP synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	445	Total	C	N	O	S	0	1	0
			3290	2102	576	599	13			
1	C	447	Total	C	N	O	S	0	0	0
			3228	2056	569	591	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	THR	-	EXPRESSION TAG	UNP E9BCQ9
C	0	THR	-	EXPRESSION TAG	UNP E9BCQ9

- Molecule 2 is URIDINE-5'-MONOPHOSPHATE (three-letter code: U5P) (formula:  $C_9H_{13}N_2O_9P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			21	9	2	9	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	C	1	Total	C	N	O	P	0	0
			21	9	2	9	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	22	Total	O	0	0
			22	22		
4	C	19	Total	O	0	0
			19	19		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: UMP synthase



#### • Molecule 1: UMP synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.72Å 64.72Å 477.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.96 – 3.00 46.96 – 3.00	Depositor EDS
% Data completeness (in resolution range)	91.7 (46.96-3.00) 91.7 (46.96-3.00)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.80 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.236 , 0.304 0.231 , 0.294	Depositor DCC
$R_{free}$ test set	996 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	66.8	Xtriage
Anisotropy	0.230	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 39.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6623	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: U5P, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	0.48	0/3356	0.57	0/4565
1	C	0.44	0/3291	0.53	0/4492
All	All	0.46	0/6647	0.55	0/9057

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3290	0	3216	37	0
1	C	3228	0	3071	36	0
2	B	21	0	11	1	0
2	C	42	0	22	0	0
3	C	1	0	0	0	0
4	B	22	0	0	0	0
4	C	19	0	0	0	0
All	All	6623	0	6320	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:431:TRP:HB2	1:B:436:ALA:HB3	1.46	0.98
1:B:296:TYR:HB3	1:B:299:ILE:HD13	1.65	0.78
1:C:308:ALA:HA	1:C:337:MET:HE1	1.65	0.78
1:B:201:ILE:HD13	1:B:225:ILE:HG23	1.65	0.77
1:B:318:ARG:NH1	1:B:365:ASP:OD2	2.19	0.75
1:B:17:CYS:HB2	1:B:224:LEU:HD22	1.71	0.72
1:B:437:ILE:HG12	1:B:441:GLN:HB3	1.76	0.68
1:C:318:ARG:HG2	1:C:340:PRO:HB2	1.77	0.66
1:C:292:ARG:O	1:C:295:THR:HG22	1.98	0.63
1:C:255:LYS:N	1:C:256:GLY:HA2	2.13	0.63
1:C:161:ARG:HA	1:C:165:PRO:HD2	1.81	0.62
1:B:438:THR:OG1	1:B:441:GLN:HB2	2.00	0.61
1:C:373:LEU:HA	1:C:399:VAL:HG13	1.83	0.61
1:B:137[A]:SER:HB3	2:B:7485:U5P:HN3	1.66	0.59
1:B:56:GLU:HG2	1:C:63:TRP:CZ2	2.37	0.59
1:C:375:SER:HB3	1:C:403:ARG:HD3	1.85	0.58
1:C:116:SER:HB3	1:C:166:TRP:NE1	2.19	0.57
1:C:308:ALA:HA	1:C:337:MET:CE	2.35	0.56
1:C:311:LEU:HD21	1:C:368:VAL:HG11	1.86	0.56
1:B:327:LEU:HB3	1:B:328:PRO:HD3	1.88	0.56
1:B:136:THR:HG21	1:C:114:MET:HG2	1.87	0.56
1:B:4:PHE:HE2	1:B:171:ASN:HA	1.71	0.55
1:B:248:ILE:O	1:B:252:ARG:HG3	2.07	0.55
1:C:28:ALA:O	1:C:32:GLU:HG2	2.07	0.54
1:B:293:LEU:HD21	1:B:299:ILE:HG22	1.89	0.54
1:B:116:SER:HB3	1:B:166:TRP:NE1	2.23	0.53
1:B:249:ASN:HA	1:B:252:ARG:HD3	1.91	0.53
1:C:249:ASN:HA	1:C:252:ARG:HG2	1.90	0.52
1:B:152:ARG:HD2	1:B:190:ARG:HH12	1.73	0.51
1:C:51:ASN:HD22	1:C:82:ASP:HB3	1.75	0.51
1:C:318:ARG:NE	1:C:360:GLU:O	2.44	0.51
1:B:271:VAL:HG12	1:B:288:ILE:HG12	1.92	0.50
1:C:322:LEU:O	1:C:326:ALA:HB3	2.11	0.49
1:C:373:LEU:HA	1:C:399:VAL:CG1	2.42	0.49
1:C:290:LEU:HD12	1:C:372:ASP:HB2	1.93	0.49
1:C:318:ARG:HB2	1:C:367:VAL:HG12	1.95	0.49
1:B:135:LYS:HA	1:B:145:GLN:OE1	2.14	0.47
1:B:431:TRP:CD1	1:B:437:ILE:HG21	2.49	0.47
1:B:134:CYS:HB2	1:B:174:LEU:HD22	1.95	0.47
1:C:17:CYS:HB2	1:C:224:LEU:HD22	1.95	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:ILE:HG22	1:C:138:ASN:HB3	1.98	0.46
1:C:150:GLY:O	1:C:152:ARG:NH1	2.49	0.46
1:B:290:LEU:HD12	1:B:372:ASP:HB2	1.98	0.45
1:B:441:GLN:O	1:B:445:VAL:HG23	2.16	0.45
1:C:375:SER:HB2	1:C:405:MET:SD	2.57	0.45
1:B:161:ARG:HA	1:B:165:PRO:HD2	1.99	0.45
1:C:10:ARG:NH2	1:C:42:HIS:O	2.50	0.44
1:B:300:MET:CE	1:B:328:PRO:HB2	2.47	0.44
1:B:92:ALA:HB3	1:B:121:PRO:HG2	2.00	0.44
1:C:135:LYS:NZ	1:C:141:SER:OG	2.48	0.43
1:B:160:GLU:OE2	1:B:190:ARG:NH1	2.51	0.43
1:B:17:CYS:HB3	1:B:226:ASN:HA	2.01	0.43
1:C:271:VAL:HG11	1:C:431:TRP:CH2	2.54	0.43
1:C:404:ASP:OD2	1:C:425:HIS:ND1	2.53	0.42
1:C:428:LEU:HB2	1:C:429:PRO:HD3	2.01	0.42
1:B:197:LEU:O	1:B:199:PRO:HD3	2.19	0.42
1:B:201:ILE:HD13	1:B:225:ILE:CG2	2.44	0.42
1:B:224:LEU:HA	1:B:224:LEU:HD23	1.94	0.41
1:C:176:VAL:HG11	1:C:184:LEU:HA	2.02	0.41
1:C:322:LEU:HA	1:C:323:PRO:HD2	1.93	0.41
1:B:198:VAL:HG21	1:B:223:MET:HE2	2.02	0.41
1:B:178:ALA:HB2	1:B:198:VAL:HG12	2.03	0.41
1:B:315:LYS:O	1:B:366:ARG:NH2	2.53	0.41
1:C:134:CYS:HB2	1:C:174:LEU:HD22	2.03	0.40
1:B:115:GLY:HA3	1:C:140:GLY:O	2.20	0.40
1:C:375:SER:HA	1:C:403:ARG:HB2	2.03	0.40
1:B:198:VAL:HG21	1:B:223:MET:CE	2.51	0.40
1:C:327:LEU:HB3	1:C:328:PRO:HD3	2.02	0.40
1:C:320:ALA:HA	1:C:342:ILE:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	440/453 (97%)	423 (96%)	17 (4%)	0	100	100
1	C	441/453 (97%)	407 (92%)	30 (7%)	4 (1%)	19	59
All	All	881/906 (97%)	830 (94%)	47 (5%)	4 (0%)	31	71

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	22	PRO
1	C	279	LYS
1	C	76	GLY
1	C	150	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	311/352 (88%)	297 (96%)	14 (4%)	30	69
1	C	295/352 (84%)	279 (95%)	16 (5%)	24	62
All	All	606/704 (86%)	576 (95%)	30 (5%)	27	65

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	23	ARG
1	B	87	ASP
1	B	95	TYR
1	B	163	GLU
1	B	255	LYS
1	B	293	LEU
1	B	295	THR
1	B	302	LEU
1	B	341	LEU
1	B	379	THR
1	B	400	LEU
1	B	403	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	417	ASP
1	B	431	TRP
1	C	13	ARG
1	C	43	GLU
1	C	144	LEU
1	C	152	ARG
1	C	184	LEU
1	C	260	GLU
1	C	263	LYS
1	C	265	LEU
1	C	299	ILE
1	C	305	ARG
1	C	366	ARG
1	C	400	LEU
1	C	403	ARG
1	C	418	PHE
1	C	424	LEU
1	C	440	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	51	ASN
1	B	102	HIS
1	B	204	GLN
1	C	51	ASN
1	C	412	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	U5P	B	7485	-	17,22,22	1.63	4 (23%)	21,33,33	2.17	4 (19%)
2	U5P	C	454	-	17,22,22	1.60	4 (23%)	21,33,33	1.74	2 (9%)
2	U5P	C	7485	-	17,22,22	1.59	3 (17%)	21,33,33	2.09	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U5P	B	7485	-	-	0/6/26/26	0/2/2/2
2	U5P	C	454	-	-	0/6/26/26	0/2/2/2
2	U5P	C	7485	-	-	0/6/26/26	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	7485	U5P	O4'-C1'	2.05	1.44	1.41
2	B	7485	U5P	P-O2P	2.12	1.63	1.54
2	C	454	U5P	P-O3P	2.31	1.64	1.54
2	C	454	U5P	O4'-C1'	2.36	1.44	1.41
2	C	7485	U5P	O4'-C1'	2.76	1.45	1.41
2	C	454	U5P	C4-N3	3.42	1.39	1.33
2	C	454	U5P	C6-N1	3.63	1.40	1.35
2	B	7485	U5P	C6-N1	3.66	1.40	1.35
2	C	7485	U5P	C4-N3	3.73	1.39	1.33
2	B	7485	U5P	C4-N3	3.76	1.39	1.33
2	C	7485	U5P	C6-N1	3.78	1.40	1.35

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	7485	U5P	C5'-C4'-C3'	-4.77	97.35	115.29
2	B	7485	U5P	C5'-C4'-C3'	-4.18	99.57	115.29
2	B	7485	U5P	C4'-O4'-C1'	-3.59	106.08	109.83
2	C	454	U5P	O5'-P-O1P	2.50	113.49	106.47
2	C	7485	U5P	O5'-P-O1P	2.87	114.53	106.47
2	B	7485	U5P	O4'-C1'-N1	3.30	114.60	108.05
2	C	454	U5P	C4-N3-C2	6.83	120.02	114.14
2	B	7485	U5P	C4-N3-C2	6.83	120.02	114.14
2	C	7485	U5P	C4-N3-C2	6.98	120.14	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	7485	U5P	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	B	445/453 (98%)	-0.43	0	100 100	44, 56, 76, 92	0
1	C	447/453 (98%)	-0.19	4 (0%)	84 62	53, 87, 103, 116	0
All	All	892/906 (98%)	-0.31	4 (0%)	92 78	44, 71, 100, 116	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	354	LYS	2.4
1	C	16	LEU	2.2
1	C	48	TYR	2.1
1	C	12	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
3	CL	C	453	1/1	0.92	0.24	73,73,73,73	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	U5P	C	454	21/21	0.94	0.17	73,79,87,89	0
2	U5P	C	7485	21/21	0.94	0.12	85,88,91,94	0
2	U5P	B	7485	21/21	0.97	0.15	55,57,59,60	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.