



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 06:25 pm GMT

PDB ID : 2QXL  
Title : Crystal Structure Analysis of Sse1, a yeast Hsp110  
Authors : Hendrickson, W.A.; Liu, Q.  
Deposited on : 2007-08-12  
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4090 (2.44-2.40)
Clashscore	122126	4587 (2.44-2.40)
Ramachandran outliers	120053	4522 (2.44-2.40)
Sidechain outliers	120020	4523 (2.44-2.40)
RSRZ outliers	108989	3987 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	658	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>6%</div> </div> </div>
1	B	658	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>10%</div> <div>5%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock protein homolog SSE1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	617	Total	C	N	O	S	0	0	0
			4802	3037	810	943	12			
1	B	625	Total	C	N	O	S	0	0	0
			4837	3058	818	949	12			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	K	0	0
			1	1		
3	A	1	Total	K	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
4	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	126	Total	O	0	0
			126	126		
5	B	229	Total	O	0	0
			229	229		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Heat shock protein homolog SSE1



- Molecule 1: Heat shock protein homolog SSE1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	218.04Å 125.33Å 74.71Å 90.00° 98.67° 90.00°	Depositor
Resolution (Å)	29.49 – 2.41 29.49 – 2.41	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.49-2.41) 99.0 (29.49-2.41)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	37.00 (at 2.42Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.195 , 0.235 0.193 , 0.233	Depositor DCC
$R_{free}$ test set	3795 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.6	Xtriage
Anisotropy	0.483	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 46.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10060	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, K, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	4/4887 (0.1%)	0.58	2/6610 (0.0%)
1	B	0.71	2/4924 (0.0%)	0.61	4/6667 (0.1%)
All	All	0.63	6/9811 (0.1%)	0.60	6/13277 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	435	ARG	CZ-NH1	33.19	1.76	1.33
1	A	435	ARG	CZ-NH1	13.21	1.50	1.33
1	A	500	ASP	C-O	10.65	1.43	1.23
1	B	409	GLN	CD-NE2	5.65	1.47	1.32
1	A	435	ARG	CZ-NH2	5.36	1.40	1.33
1	A	409	GLN	CD-NE2	5.24	1.46	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	435	ARG	NE-CZ-NH2	-15.38	112.61	120.30
1	B	435	ARG	NE-CZ-NH2	-7.60	116.50	120.30
1	A	435	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	B	385	PRO	N-CA-CB	5.93	110.42	103.30
1	B	470	PRO	N-CA-CB	5.88	110.36	103.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	141	VAL	CB-CA-C	-5.79	100.39	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	414	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4802	0	4732	38	0
1	B	4837	0	4730	36	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	31	0	12	0	0
4	B	31	0	12	0	0
5	A	126	0	0	1	0
5	B	229	0	0	1	0
All	All	10060	0	9486	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:435:ARG:CZ	1:B:435:ARG:NH1	1.76	1.47
1:A:47:ARG:HD2	1:A:121:MET:CE	1.96	0.95
1:A:47:ARG:HD2	1:A:121:MET:HE3	1.53	0.89
1:B:61:ILE:O	1:B:89:THR:HG23	1.80	0.80
1:A:141:VAL:HG21	1:A:166:LEU:HD13	1.63	0.79

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:238:ASP:O	1:B:242:THR:HG23	1.84	0.78
1:A:238:ASP:O	1:A:242:THR:HG23	1.85	0.76
1:B:242:THR:HG21	1:B:268:LEU:HA	1.71	0.73
1:A:47:ARG:HD2	1:A:121:MET:HE1	1.69	0.73
1:A:645:LYS:O	1:A:649:LYS:HG2	1.90	0.72
1:A:61:ILE:O	1:A:89:THR:HG23	1.89	0.70
1:B:582:ARG:HG2	1:B:605:LEU:HD13	1.74	0.69
1:A:576:GLU:O	1:A:580:THR:HG23	1.93	0.68
1:B:257:ILE:HG22	1:B:292:MET:HE2	1.75	0.68
1:A:308:GLU:HA	1:A:311:LYS:HE2	1.76	0.67
1:B:350:LEU:O	1:B:354:ILE:HG23	1.97	0.64
1:B:433:LEU:HB3	1:B:435:ARG:HD3	1.81	0.63
1:B:432:THR:HG22	1:B:479:LYS:HG2	1.82	0.62
1:A:141:VAL:CG2	1:A:166:LEU:HD13	2.30	0.61
1:B:5:PHE:CD2	1:B:141:VAL:HG13	2.35	0.60
1:B:257:ILE:HG22	1:B:292:MET:CE	2.32	0.60
1:B:247:ASP:OD1	1:B:258:ARG:NH2	2.34	0.58
1:A:478:VAL:HG12	1:A:497:THR:HG22	1.86	0.58
1:B:335:VAL:O	1:B:360:LYS:HE2	2.03	0.57
1:A:11:ASN:HD21	1:A:67:ASN:HA	1.68	0.57
1:B:11:ASN:HD21	1:B:67:ASN:HA	1.70	0.56
1:A:242:THR:HG21	1:A:268:LEU:HA	1.87	0.56
1:B:311:LYS:HB3	1:B:312:PRO:HD3	1.88	0.55
1:B:11:ASN:ND2	1:B:67:ASN:HA	2.21	0.55
1:A:11:ASN:ND2	1:A:67:ASN:HA	2.23	0.54
1:A:379:ILE:HA	1:A:382:ILE:HD12	1.91	0.53
1:B:242:THR:HG22	1:B:271:ALA:CB	2.40	0.52
1:B:604:MET:HE1	1:B:630:GLU:HG2	1.92	0.51
1:A:432:THR:HG22	1:A:479:LYS:HG2	1.92	0.51
1:A:141:VAL:HG23	1:A:166:LEU:HB3	1.92	0.51
1:B:131:LYS:HG2	1:B:138:ILE:HG12	1.94	0.49
1:B:202:VAL:HG21	1:B:354:ILE:HD13	1.94	0.48
1:A:78:HIS:ND1	1:A:79:PRO:HD2	2.29	0.48
1:A:174:ASP:HB2	1:A:372:ILE:HD13	1.96	0.48
1:B:141:VAL:HG22	1:B:166:LEU:HB3	1.95	0.48
1:B:12:ASN:HB2	1:B:206:HIS:CG	2.49	0.47
1:A:238:ASP:O	1:A:242:THR:CG2	2.60	0.47
1:B:569:ASP:OD2	1:B:570:ARG:NH1	2.48	0.47
1:A:485:ASP:HB2	1:A:486:PRO:CD	2.46	0.45
1:A:12:ASN:HB2	1:A:206:HIS:CG	2.51	0.45
1:B:11:ASN:ND2	1:B:70:ARG:HG2	2.31	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:582:ARG:HG2	1:A:605:LEU:HD13	1.99	0.45
1:B:47:ARG:NH1	1:B:121:MET:HG2	2.32	0.45
1:B:570:ARG:HG3	1:B:621:LYS:HG3	1.97	0.45
1:A:589:TYR:HB3	1:A:638:ILE:HG21	1.99	0.45
1:A:242:THR:HB	1:A:267:ILE:HG22	1.98	0.44
1:A:498:ILE:CG2	1:A:499:GLU:N	2.80	0.44
1:B:414:ASP:OD1	1:B:415:HIS:HB2	2.18	0.44
1:A:67:ASN:HB2	1:A:88:PHE:CZ	2.53	0.43
1:A:69:LYS:HE2	5:A:1022:HOH:O	2.18	0.43
1:A:40:VAL:HG13	1:A:121:MET:HE2	2.00	0.43
1:A:220:GLN:HA	1:A:393:LYS:O	2.18	0.43
1:B:275:LYS:HB3	1:B:275:LYS:HE2	1.85	0.43
1:A:453:PRO:O	1:A:454:ASN:HB2	2.18	0.43
1:A:242:THR:HG21	1:A:268:LEU:HD12	2.00	0.42
1:B:587:GLU:C	5:B:1222:HOH:O	2.57	0.42
1:B:338:VAL:HG11	1:B:354:ILE:HD11	2.00	0.42
1:B:476:VAL:HA	1:B:477:PRO:HD3	1.89	0.42
1:A:476:VAL:HA	1:A:477:PRO:HD3	1.82	0.42
1:B:97:ASP:O	1:B:98:LYS:HB2	2.20	0.42
1:B:139:THR:HG23	1:B:140:ASP:OD2	2.20	0.41
1:B:581:LEU:HD23	1:B:605:LEU:HD21	2.02	0.41
1:A:543:ASP:OD1	1:A:546:LYS:HB2	2.20	0.41
1:A:11:ASN:ND2	1:A:70:ARG:HG2	2.36	0.41
1:A:379:ILE:O	1:A:382:ILE:HB	2.21	0.41
1:A:448:ILE:HD13	1:A:457:GLU:HA	2.03	0.41
1:A:217:LYS:HB2	1:A:220:GLN:HG2	2.02	0.41
1:B:582:ARG:HH22	1:B:609:GLU:CD	2.24	0.41
1:B:379:ILE:O	1:B:382:ILE:HG12	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	609/658 (93%)	594 (98%)	15 (2%)	0	100	100
1	B	619/658 (94%)	599 (97%)	20 (3%)	0	100	100
All	All	1228/1316 (93%)	1193 (97%)	35 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	514/567 (91%)	497 (97%)	17 (3%)	41	60
1	B	512/567 (90%)	493 (96%)	19 (4%)	37	55
All	All	1026/1134 (90%)	990 (96%)	36 (4%)	39	57

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	61	ILE
1	A	89	THR
1	A	105	ARG
1	A	149	TYR
1	A	195	LYS
1	A	209	TYR
1	A	242	THR
1	A	252	LYS
1	A	298	SER
1	A	321	VAL
1	A	354	ILE
1	A	360	LYS
1	A	433	LEU
1	A	482	LEU
1	A	531	ASP
1	A	580	THR
1	B	9	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	75	ASP
1	B	89	THR
1	B	105	ARG
1	B	141	VAL
1	B	149	TYR
1	B	195	LYS
1	B	209	TYR
1	B	230	LYS
1	B	242	THR
1	B	335	VAL
1	B	354	ILE
1	B	389	VAL
1	B	433	LEU
1	B	468	GLN
1	B	588	GLU
1	B	602	GLN
1	B	634	LEU
1	B	643	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	B	11	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ATP	A	1002	3,2	27,33,33	0.91	1 (3%)	27,52,52	1.99	3 (11%)
4	ATP	B	1001	3,2	27,33,33	1.02	2 (7%)	27,52,52	2.26	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	1002	3,2	-	0/18/38/38	0/3/3/3
4	ATP	B	1001	3,2	-	0/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1001	ATP	PG-O3B	2.61	1.64	1.60
4	A	1002	ATP	C5-C4	2.77	1.46	1.40
4	B	1001	ATP	C5-C4	2.90	1.47	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1001	ATP	N3-C2-N1	-10.06	120.25	128.86
4	A	1002	ATP	N3-C2-N1	-8.40	121.67	128.86
4	A	1002	ATP	PA-O3A-PB	-2.23	125.14	132.63
4	B	1001	ATP	C1'-N9-C4	-2.16	122.90	126.64
4	B	1001	ATP	O3G-PG-O2G	2.14	116.06	107.59
4	A	1002	ATP	C2-N1-C6	2.44	122.89	118.75
4	B	1001	ATP	C2-N1-C6	3.32	124.40	118.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	617/658 (93%)	0.33	31 (5%) 29 26	46, 58, 67, 74	0
1	B	625/658 (94%)	0.11	30 (4%) 30 28	36, 46, 61, 72	0
All	All	1242/1316 (94%)	0.22	61 (4%) 29 27	36, 53, 66, 74	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	472	GLY	6.1
1	B	527	THR	5.6
1	B	498	ILE	4.8
1	A	22	ARG	4.7
1	B	500	ASP	4.5
1	B	385	PRO	4.5
1	A	596	ALA	4.4
1	B	501	ILE	3.5
1	B	650	LYS	3.5
1	A	7	LEU	3.4
1	B	22	ARG	3.4
1	A	500	ASP	3.3
1	B	471	GLU	3.3
1	A	597	GLU	3.2
1	B	470	PRO	3.2
1	A	528	VAL	3.2
1	A	498	ILE	3.1
1	B	387	LEU	3.0
1	A	410	VAL	2.9
1	B	386	THR	2.9
1	A	195	LYS	2.8
1	B	145	VAL	2.8
1	A	527	THR	2.7
1	A	145	VAL	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	382	ILE	2.7
1	B	474	ASP	2.7
1	A	193	GLU	2.6
1	B	502	GLU	2.6
1	A	412	ASP	2.6
1	B	454	ASN	2.6
1	A	391	PRO	2.5
1	B	146	PRO	2.5
1	B	7	LEU	2.5
1	A	390	ARG	2.5
1	A	526	LYS	2.4
1	A	383	HIS	2.4
1	B	143	ILE	2.4
1	A	595	ASP	2.4
1	B	651	GLN	2.4
1	A	118	LEU	2.3
1	A	329	LYS	2.3
1	A	333	GLU	2.3
1	B	123	ILE	2.3
1	B	383	HIS	2.3
1	A	332	ALA	2.3
1	A	543	ASP	2.3
1	B	384	SER	2.2
1	A	227	ALA	2.2
1	B	652	ALA	2.1
1	A	331	SER	2.1
1	A	384	SER	2.1
1	B	388	ARG	2.1
1	B	144	ALA	2.1
1	A	21	ASN	2.1
1	A	147	PRO	2.1
1	B	121	MET	2.0
1	A	446	THR	2.0
1	B	119	ALA	2.0
1	B	147	PRO	2.0
1	A	146	PRO	2.0
1	B	210	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	1004	1/1	0.93	0.19	45,45,45,45	0
2	MG	B	1003	1/1	0.94	0.06	28,28,28,28	0
4	ATP	A	1002	31/31	0.98	0.07	35,38,43,44	0
3	K	B	1006	1/1	0.99	0.03	36,36,36,36	0
4	ATP	B	1001	31/31	0.99	0.07	25,28,30,30	0
3	K	A	1005	1/1	0.99	0.11	48,48,48,48	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.