



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 4, 2019 – 01:52 PM EDT

PDB ID : 6R2C  
Title : Crystal structure of the SucA domain of Mycobacterium smegmatis KGD after soaking with succinylphosphonate phosphonoethyl ester (PESP)  
Authors : Wagner, T.; Alzari, P.M.; Bellinzoni, M.  
Deposited on : 2019-03-15  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.4  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	868	<div> <div>3%</div> <div>90%</div> <div>6%</div> </div>
1	B	868	<div> <div>4%</div> <div>91%</div> <div>6%</div> </div>
1	C	868	<div> <div>5%</div> <div>90%</div> <div>6%</div> </div>
1	D	868	<div> <div>3%</div> <div>91%</div> <div>6%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 26262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Multifunctional 2-oxoglutarate metabolism enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	814	Total	C	N	O	S	0	1	0
			6331	3987	1121	1199	24			
1	B	815	Total	C	N	O	S	0	0	0
			6288	3965	1118	1181	24			
1	C	813	Total	C	N	O	S	0	2	0
			6357	4003	1126	1205	23			
1	D	812	Total	C	N	O	S	0	0	0
			6275	3954	1107	1191	23			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	360	GLY	-	expression tag	UNP A0R2B1
B	360	GLY	-	expression tag	UNP A0R2B1
C	360	GLY	-	expression tag	UNP A0R2B1
D	360	GLY	-	expression tag	UNP A0R2B1

- Molecule 2 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O<sub>7</sub>P<sub>2</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
2	B	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
2	C	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0
2	D	1	Total 26	C 12	N 4	O 7	P 2	S 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ca 1 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	D	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		

- Molecule 5 is 4-[ethoxy(oxidanyl)phosphoryl]-4-oxidanylidene-butanoic acid (three-letter code: JQ5) (formula: C<sub>6</sub>H<sub>11</sub>O<sub>6</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	P	0	0
			13	6	6	1		
5	B	1	Total	C	O	P	0	0
			13	6	6	1		
5	C	1	Total	C	O	P	0	0
			13	6	6	1		
5	D	1	Total	C	O	P	0	0
			13	6	6	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	219	Total	O	0	0
			219	219		
6	B	214	Total	O	0	0
			214	214		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	192	Total 192	O 192	0	0
6	D	222	Total 222	O 222	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme




- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme





- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme





● Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.39Å 83.96Å 160.68Å 99.54° 98.78° 100.89°	Depositor
Resolution (Å)	21.07 – 2.09 78.47 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.4 (21.07-2.09) 97.4 (78.47-2.09)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.48 (at 2.08Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.232 , 0.246 0.234 , 0.251	Depositor DCC
$R_{free}$ test set	11720 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.1	Xtriage
Anisotropy	0.644	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	26262	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.93 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8946e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, JQ5, MG, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/6463	0.58	0/8761
1	B	0.43	0/6417	0.58	0/8705
1	C	0.43	0/6492	0.58	0/8796
1	D	0.43	0/6403	0.58	0/8688
All	All	0.44	0/25775	0.58	0/34950

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6331	0	6125	14	0
1	B	6288	0	6072	13	0
1	C	6357	0	6172	12	0
1	D	6275	0	6050	9	0
2	A	26	0	16	3	0
2	B	26	0	16	3	0
2	C	26	0	16	4	0
2	D	26	0	16	4	0
3	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	13	0	0	3	0
5	B	13	0	0	2	0
5	C	13	0	0	4	0
5	D	13	0	0	4	0
6	A	219	0	0	0	0
6	B	214	0	0	1	0
6	C	192	0	0	0	0
6	D	222	0	0	0	0
All	All	26262	0	24483	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:2001:TPP:H2	5:C:2004:JQ5:C6	1.83	1.09
2:A:2001:TPP:H2	5:A:2004:JQ5:C6	1.82	1.08
2:D:2001:TPP:H2	5:D:2004:JQ5:C6	1.83	1.07
2:B:2001:TPP:H2	5:B:2004:JQ5:C6	1.96	0.95
2:A:2001:TPP:H2	5:A:2004:JQ5:C5	2.29	0.63
2:C:2001:TPP:H2	5:C:2004:JQ5:C5	2.28	0.63
2:D:2001:TPP:H2	5:D:2004:JQ5:C5	2.28	0.62
2:B:2001:TPP:H2	5:B:2004:JQ5:C5	2.31	0.60
1:A:778:LEU:HB3	1:A:784:ILE:HG12	1.89	0.54
1:C:848:LEU:HD11	1:C:865:GLU:HA	1.93	0.50
1:D:744:ARG:HH12	1:D:767:LYS:C	2.15	0.50
1:B:744:ARG:HH12	1:B:767:LYS:C	2.15	0.49
1:C:705:PRO:HG2	1:C:735:VAL:HG13	1.94	0.48
1:C:1112:LEU:HD21	1:C:1155:LEU:HD22	1.96	0.48
1:A:898:LEU:O	1:A:945:VAL:HA	2.14	0.47
2:A:2001:TPP:H2	5:A:2004:JQ5:O7	2.12	0.47
1:B:778:LEU:HB3	1:B:784:ILE:HG12	1.96	0.47
1:A:976:GLN:OE1	2:B:2001:TPP:HM43	2.14	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:778:LEU:HB3	1:C:784:ILE:HG12	1.96	0.47
2:C:2001:TPP:C2	5:C:2004:JQ5:C6	2.76	0.47
1:B:1119:ILE:HD12	1:B:1169:GLU:HG2	1.99	0.45
1:C:1177:TRP:CD1	1:C:1197:ARG:HD3	2.52	0.45
1:D:1177:TRP:CD1	1:D:1197:ARG:HD3	2.52	0.45
1:C:1112:LEU:CD2	1:C:1155:LEU:HD22	2.47	0.45
2:D:2001:TPP:C2	5:D:2004:JQ5:C6	2.75	0.45
1:B:1002:LEU:HB3	1:B:1061:GLN:HB2	2.00	0.44
1:B:1177:TRP:CD1	1:B:1197:ARG:HD3	2.52	0.44
1:A:1177:TRP:CD1	1:A:1197:ARG:HD3	2.52	0.44
1:D:1155:LEU:HD13	1:D:1164:LYS:HE2	2.00	0.44
1:D:1002:LEU:HB3	1:D:1061:GLN:HB2	2.00	0.44
1:A:530:LEU:HD22	1:A:636:SER:HA	2.00	0.44
1:B:1155:LEU:HD13	1:B:1164:LYS:HE2	2.00	0.43
2:D:2001:TPP:H2	5:D:2004:JQ5:O7	2.13	0.43
1:C:898:LEU:O	1:C:945:VAL:HA	2.18	0.43
1:B:486:LEU:HD11	1:B:714:PRO:HG3	2.01	0.43
1:D:898:LEU:O	1:D:945:VAL:HA	2.19	0.42
1:C:1002:LEU:HB3	1:C:1061:GLN:HB2	2.00	0.42
1:A:1155:LEU:HD13	1:A:1164:LYS:HE2	2.00	0.42
1:C:1155:LEU:HD13	1:C:1164:LYS:HE2	2.00	0.42
1:A:705:PRO:HG2	1:A:735:VAL:HG13	2.01	0.42
1:B:898:LEU:O	1:B:945:VAL:HA	2.20	0.42
1:B:541:GLY:HA2	6:B:2125:HOH:O	2.20	0.42
1:A:981:VAL:HG22	1:A:988:ILE:HD11	2.02	0.42
1:D:1091:ARG:HD2	1:D:1150:ARG:NH2	2.35	0.41
1:A:1002:LEU:HB3	1:A:1061:GLN:HB2	2.02	0.41
1:A:1112:LEU:HD12	1:A:1161:VAL:HG21	2.03	0.41
1:D:1069:PRO:HB2	1:D:1072:MET:HB3	2.02	0.41
1:B:1155:LEU:HD11	1:B:1192:PHE:CZ	2.56	0.41
1:C:1155:LEU:HD11	1:C:1192:PHE:CZ	2.56	0.41
1:C:981:VAL:HG22	1:C:988:ILE:HD11	2.02	0.41
1:A:626:THR:HG21	1:A:636:SER:OG	2.21	0.41
1:B:1112:LEU:HD13	1:B:1155:LEU:HD22	2.02	0.41
1:C:486:LEU:HD11	1:C:714:PRO:HG3	2.02	0.41
1:B:981:VAL:HG22	1:B:988:ILE:HD11	2.03	0.41
1:A:1155:LEU:HD11	1:A:1192:PHE:CZ	2.56	0.41
1:A:1119:ILE:HD12	1:A:1169:GLU:HG2	2.03	0.40
1:D:1119:ILE:HD12	1:D:1169:GLU:HG2	2.04	0.40
2:C:2001:TPP:H2	5:C:2004:JQ5:O7	2.14	0.40
1:D:1155:LEU:HD11	1:D:1192:PHE:CZ	2.56	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1069:PRO:HB2	1:A:1072:MET:HB3	2.03	0.40
1:B:1069:PRO:HB2	1:B:1072:MET:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	805/868 (93%)	791 (98%)	14 (2%)	0	100	100
1	B	805/868 (93%)	790 (98%)	15 (2%)	0	100	100
1	C	805/868 (93%)	787 (98%)	18 (2%)	0	100	100
1	D	802/868 (92%)	787 (98%)	15 (2%)	0	100	100
All	All	3217/3472 (93%)	3155 (98%)	62 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	654/726 (90%)	646 (99%)	8 (1%)	74	80
1	B	643/726 (89%)	638 (99%)	5 (1%)	83	88
1	C	660/726 (91%)	651 (99%)	9 (1%)	69	76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	645/726 (89%)	640 (99%)	5 (1%)	83	88
All	All	2602/2904 (90%)	2575 (99%)	27 (1%)	78	83

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	367	ASN
1	A	464	GLN
1	A	539	HIS
1	A	682	PHE
1	A	868	ARG
1	A	953	PHE
1	A	976	GLN
1	A	1112	LEU
1	B	367	ASN
1	B	539	HIS
1	B	682	PHE
1	B	953	PHE
1	B	976	GLN
1	C	367	ASN
1	C	384	MET
1	C	464	GLN
1	C	539	HIS
1	C	682	PHE
1	C	831	LEU
1	C	848	LEU
1	C	953	PHE
1	C	976	GLN
1	D	539	HIS
1	D	682	PHE
1	D	953	PHE
1	D	976	GLN
1	D	1112	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	TPP	A	2001	3	22,27,27	0.59	0	26,40,40	0.74	0
5	JQ5	A	2004	-	5,12,12	4.63	1 (20%)	7,16,16	2.51	3 (42%)
2	TPP	B	2001	3	22,27,27	0.63	0	26,40,40	0.77	0
5	JQ5	B	2004	-	5,12,12	4.55	1 (20%)	7,16,16	2.49	3 (42%)
2	TPP	C	2001	3	22,27,27	0.44	0	26,40,40	0.77	0
5	JQ5	C	2004	-	5,12,12	4.72	1 (20%)	7,16,16	2.56	3 (42%)
2	TPP	D	2001	3	22,27,27	0.56	0	26,40,40	0.76	0
5	JQ5	D	2004	-	5,12,12	4.82	1 (20%)	7,16,16	2.35	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPP	A	2001	3	-	0/16/17/17	0/2/2/2
5	JQ5	A	2004	-	-	2/6/15/15	-
2	TPP	B	2001	3	-	1/16/17/17	0/2/2/2
5	JQ5	B	2004	-	-	2/6/15/15	-
2	TPP	C	2001	3	-	1/16/17/17	0/2/2/2

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	JQ5	C	2004	-	-	2/6/15/15	-
2	TPP	D	2001	3	-	0/16/17/17	0/2/2/2
5	JQ5	D	2004	-	-	2/6/15/15	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	2004	JQ5	P10-O12	10.60	1.71	1.57
5	C	2004	JQ5	P10-O12	10.38	1.71	1.57
5	A	2004	JQ5	P10-O12	10.15	1.71	1.57
5	B	2004	JQ5	P10-O12	9.99	1.71	1.57

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	2004	JQ5	O12-P10-O15	-4.27	103.06	114.93
5	A	2004	JQ5	O12-P10-O15	-4.17	103.32	114.93
5	B	2004	JQ5	O12-P10-O15	-4.07	103.61	114.93
5	A	2004	JQ5	O11-P10-O15	3.89	124.35	110.50
5	B	2004	JQ5	O11-P10-O15	3.86	124.27	110.50
5	D	2004	JQ5	O11-P10-O15	3.77	123.94	110.50
5	C	2004	JQ5	O11-P10-O15	3.75	123.84	110.50
5	C	2004	JQ5	C4-C5-C6	3.42	117.81	113.70
5	D	2004	JQ5	C4-C5-C6	3.39	117.77	113.70
5	D	2004	JQ5	O12-P10-O15	-3.31	105.72	114.93
5	B	2004	JQ5	C4-C5-C6	3.26	117.62	113.70
5	A	2004	JQ5	C4-C5-C6	3.13	117.47	113.70

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	2004	JQ5	C13-O12-P10-O11
5	D	2004	JQ5	C13-O12-P10-O15
5	C	2004	JQ5	C13-O12-P10-O15
5	C	2004	JQ5	C14-C13-O12-P10
5	A	2004	JQ5	C14-C13-O12-P10
2	B	2001	TPP	PB-O3A-PA-O7
5	D	2004	JQ5	C14-C13-O12-P10
5	B	2004	JQ5	C14-C13-O12-P10
5	A	2004	JQ5	C13-O12-P10-O11

*Continued on next page...*



*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	C	2001	TPP	PB-O3A-PA-O7

There are no ring outliers.

8 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	TPP	3	0
5	A	2004	JQ5	3	0
2	B	2001	TPP	3	0
5	B	2004	JQ5	2	0
2	C	2001	TPP	4	0
5	C	2004	JQ5	4	0
2	D	2001	TPP	4	0
5	D	2004	JQ5	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	814/868 (93%)	0.37	26 (3%)	47	55	22, 39, 68, 103	0
1	B	815/868 (93%)	0.35	31 (3%)	40	47	22, 39, 74, 103	0
1	C	813/868 (93%)	0.43	41 (5%)	29	35	24, 41, 73, 103	0
1	D	812/868 (93%)	0.35	29 (3%)	42	49	24, 40, 70, 95	0
All	All	3254/3472 (93%)	0.38	127 (3%)	39	46	22, 40, 72, 103	0

All (127) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	561	PHE	6.9
1	B	394	ASN	6.1
1	C	394	ASN	6.0
1	D	413	TRP	5.7
1	D	501	VAL	5.2
1	B	472	ASP	5.0
1	A	1083	ARG	5.0
1	C	779	ILE	4.8
1	D	831	LEU	4.7
1	A	561	PHE	4.6
1	B	813	HIS	4.6
1	A	411	THR	4.4
1	C	630	GLY	4.3
1	C	472	ASP	4.3
1	B	473	LYS	4.2
1	A	399	SER	4.2
1	B	371	ILE	4.2
1	A	394	ASN	4.2
1	B	419	PHE	4.1
1	B	1102	GLY	4.1
1	C	808	ARG	4.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	1083	ARG	4.1
1	B	501	VAL	4.0
1	C	626	THR	4.0
1	B	810	LEU	3.9
1	C	413	TRP	3.8
1	C	633	ASN	3.8
1	A	413	TRP	3.8
1	D	810	LEU	3.8
1	B	1101	ASP	3.7
1	D	793	LEU	3.6
1	B	413	TRP	3.6
1	D	797	GLN	3.5
1	A	812	LYS	3.4
1	D	371	ILE	3.4
1	D	397	PHE	3.3
1	D	631	SER	3.3
1	C	471	HIS	3.3
1	C	792	ALA	3.2
1	A	1210	SER	3.2
1	B	793	LEU	3.2
1	D	394	ASN	3.1
1	D	817	PRO	3.1
1	D	630	GLY	3.1
1	C	805	ASN	3.1
1	C	1210	SER	3.0
1	C	506	PHE	3.0
1	B	368	ALA	3.0
1	A	472	ASP	3.0
1	C	831	LEU	3.0
1	A	371	ILE	3.0
1	C	634	ARG	2.9
1	A	750	GLY	2.9
1	C	368	ALA	2.9
1	C	500	TYR	2.9
1	D	808	ARG	2.8
1	C	371	ILE	2.8
1	A	785	SER	2.8
1	C	397	PHE	2.8
1	D	807	VAL	2.8
1	A	575	ASP	2.7
1	B	805	ASN	2.7
1	B	395	THR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	471	HIS	2.7
1	C	796	TYR	2.7
1	C	807	VAL	2.7
1	B	817	PRO	2.6
1	C	629	GLU	2.6
1	B	786	MET	2.6
1	B	745	ARG	2.6
1	D	589	MET	2.6
1	B	807	VAL	2.5
1	C	812	LYS	2.5
1	A	497	GLN	2.5
1	C	935	ASP	2.5
1	C	489	ALA	2.5
1	A	501	VAL	2.5
1	C	412	LEU	2.4
1	B	502	GLY	2.4
1	B	561	PHE	2.4
1	C	852	GLU	2.4
1	B	779	ILE	2.4
1	C	830	LYS	2.4
1	C	632	ASP	2.4
1	C	789	ALA	2.4
1	B	470	LYS	2.4
1	D	368	ALA	2.4
1	C	481	TYR	2.3
1	B	475	THR	2.3
1	A	857	HIS	2.3
1	B	796	TYR	2.3
1	B	804	PHE	2.3
1	D	813	HIS	2.3
1	D	749	GLU	2.3
1	C	750	GLY	2.3
1	A	920	THR	2.2
1	C	920	THR	2.2
1	D	395	THR	2.2
1	D	934	PRO	2.2
1	B	789	ALA	2.2
1	D	1101	ASP	2.2
1	D	756	THR	2.2
1	A	397	PHE	2.1
1	D	561	PHE	2.1
1	C	794	ARG	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	592	ASP	2.1
1	A	398	ARG	2.1
1	B	1098	MET	2.1
1	D	1220	GLU	2.1
1	A	500	TYR	2.1
1	B	777	ALA	2.1
1	A	779	ILE	2.1
1	A	740	LEU	2.1
1	D	471	HIS	2.1
1	D	502	GLY	2.1
1	A	803	VAL	2.1
1	D	745	ARG	2.1
1	A	796	TYR	2.1
1	C	497	GLN	2.0
1	C	589	MET	2.0
1	D	473	LYS	2.0
1	D	759	TYR	2.0
1	A	805	ASN	2.0
1	C	810	LEU	2.0
1	B	766	THR	2.0
1	C	775	THR	2.0
1	A	844	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	JQ5	D	2004	13/13	0.91	0.19	39,45,48,49	0

*Continued on next page...*


*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	JQ5	A	2004	13/13	0.91	0.20	38,42,46,48	0
5	JQ5	C	2004	13/13	0.91	0.21	39,44,54,55	0
3	MG	B	2002	1/1	0.95	0.09	27,27,27,27	0
5	JQ5	B	2004	13/13	0.95	0.19	43,48,52,52	0
2	TPP	B	2001	26/26	0.96	0.14	22,25,29,30	0
2	TPP	D	2001	26/26	0.96	0.13	22,26,30,30	0
2	TPP	C	2001	26/26	0.96	0.14	23,27,31,31	0
3	MG	D	2002	1/1	0.96	0.11	26,26,26,26	0
3	MG	C	2002	1/1	0.96	0.07	23,23,23,23	0
4	CA	A	2003	1/1	0.97	0.08	35,35,35,35	0
2	TPP	A	2001	26/26	0.97	0.13	19,26,29,31	0
4	CA	C	2003	1/1	0.98	0.05	37,37,37,37	0
3	MG	A	2002	1/1	0.98	0.10	24,24,24,24	0
4	CA	D	2003	1/1	0.99	0.11	36,36,36,36	0
4	CA	B	2003	1/1	0.99	0.10	39,39,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.