



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 17, 2019 – 09:22 PM EDT

PDB ID : 6RFC  
Title : Crystal structure of the potassium-pumping G263F mutant of the light-driven sodium pump KR2 in the monomeric form, pH 4.3  
Authors : Kovalev, K.; Polovinkin, V.; Gushchin, I.; Borshchevskiy, V.; Gordeliy, V.  
Deposited on : 2019-04-12  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031633  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031633


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	7193 (2.00-2.00)
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)
RSRZ outliers	108989	7011 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	<div> <div>3%</div> <div>81%</div> <div>11%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LFA	A	324	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LFA	A	335	-	-	X	-
5	GOL	A	331	-	-	-	X
5	GOL	A	333	-	-	X	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2573 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sodium pumping rhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	0	2	0
			2161	1452	325	375	9			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	263	PHE	GLY	engineered mutation	UNP N0DKS8
A	281	LEU	-	expression tag	UNP N0DKS8
A	282	GLU	-	expression tag	UNP N0DKS8
A	283	HIS	-	expression tag	UNP N0DKS8
A	284	HIS	-	expression tag	UNP N0DKS8
A	285	HIS	-	expression tag	UNP N0DKS8
A	286	HIS	-	expression tag	UNP N0DKS8
A	287	HIS	-	expression tag	UNP N0DKS8
A	288	HIS	-	expression tag	UNP N0DKS8

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is EICOSANE (three-letter code: LFA) (formula: C<sub>20</sub>H<sub>42</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 16 16	0	0
3	A	1	Total C 9 9	0	0
3	A	1	Total C 6 6	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 13 13	0	0
3	A	1	Total C 8 8	0	0
3	A	1	Total C 8 8	0	0
3	A	1	Total C 12 12	0	0
3	A	1	Total C 10 10	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 11 11	0	0
3	A	1	Total C 12 12	0	0
3	A	1	Total C 10 10	0	0

Continued on next page...

*Continued from previous page...*


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 6 6	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 6 6	0	0
3	A	1	Total C 6 6	0	0
3	A	1	Total C 10 10	0	0
3	A	1	Total C 7 7	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 11 11	0	0
3	A	1	Total C 4 4	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 10 10	0	0
3	A	1	Total C 12 12	0	0
3	A	1	Total C 14 14	0	0
3	A	1	Total C 18 18	0	0
3	A	1	Total C 8 8	0	0

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C<sub>21</sub>H<sub>40</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			25	21	4		
4	A	1	Total	C	O	0	0
			25	21	4		
4	A	1	Total	C	O	0	0
			12	8	4		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is RETINAL (three-letter code: RET) (formula: C<sub>20</sub>H<sub>28</sub>O).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	C	0	0
			20	20		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	54	Total	O	0	0
			54	54		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sodium pumping rhodopsin



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.77Å 81.96Å 232.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 40.50 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-2.00) 99.3 (40.50-2.00)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.89 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, $R_{free}$	0.177 , 0.228 0.191 , 0.238	Depositor DCC
$R_{free}$ test set	1306 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.3	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 65.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2573	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OLC, LFA, RET, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/2228	0.42	0/3030

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	109	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2161	0	2128	69	0
2	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	257	0	467	36	0
4	A	62	0	93	2	0
5	A	18	0	24	12	3
6	A	20	0	27	7	0
7	A	54	0	0	3	0
All	All	2573	0	2739	78	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179[A]:PHE:CE1	3:A:335:LFA:H142	1.61	1.32
1:A:183[B]:TRP:HE3	3:A:335:LFA:C13	1.42	1.31
1:A:183[B]:TRP:CE3	3:A:335:LFA:C13	2.17	1.27
1:A:179[A]:PHE:CZ	3:A:335:LFA:H142	1.71	1.24
1:A:183[B]:TRP:CE3	3:A:335:LFA:H132	1.77	1.14

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:333:GOL:O1	5:A:333:GOL:O1[3_555]	1.60	0.60
5:A:333:GOL:O2	5:A:333:GOL:O2[3_555]	1.87	0.33
5:A:333:GOL:C2	5:A:333:GOL:O2[3_555]	2.08	0.12

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/288 (94%)	263 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/249 (90%)	217 (97%)	7 (3%)	43	42

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	233	PHE
1	A	272	ASN
1	A	270	SER
1	A	22	GLU
1	A	271	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	106	ASN
1	A	112	ASN
1	A	157	GLN
1	A	206	ASN
1	A	272	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 37 ligands modelled in this entry, 1 is monoatomic - leaving 36 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	LFA	A	302	-	15,15,19	0.28	0	14,14,18	0.46	0
3	LFA	A	303	-	8,8,19	0.29	0	7,7,18	0.43	0
3	LFA	A	304	-	5,5,19	0.27	0	4,4,18	0.35	0
3	LFA	A	305	-	4,4,19	0.29	0	3,3,18	0.36	0
3	LFA	A	306	-	12,12,19	0.22	0	11,11,18	0.60	0
3	LFA	A	307	-	7,7,19	0.24	0	6,6,18	0.48	0
3	LFA	A	308	-	7,7,19	0.26	0	6,6,18	0.46	0
3	LFA	A	309	-	11,11,19	0.29	0	10,10,18	0.43	0
3	LFA	A	310	-	9,9,19	0.28	0	8,8,18	0.44	0
3	LFA	A	311	-	4,4,19	0.29	0	3,3,18	0.38	0
3	LFA	A	312	-	4,4,19	0.31	0	3,3,18	0.35	0
3	LFA	A	313	-	10,10,19	0.30	0	9,9,18	0.46	0
3	LFA	A	314	-	11,11,19	0.27	0	10,10,18	0.48	0
3	LFA	A	315	-	9,9,19	0.26	0	8,8,18	0.49	0
3	LFA	A	316	-	5,5,19	0.26	0	4,4,18	0.37	0
3	LFA	A	317	-	4,4,19	0.26	0	3,3,18	0.36	0
3	LFA	A	318	-	5,5,19	0.28	0	4,4,18	0.32	0
3	LFA	A	319	-	5,5,19	0.24	0	4,4,18	0.41	0
3	LFA	A	320	-	9,9,19	0.24	0	8,8,18	0.54	0
3	LFA	A	321	-	6,6,19	0.27	0	5,5,18	0.38	0
3	LFA	A	322	-	4,4,19	0.29	0	3,3,18	0.39	0
3	LFA	A	323	-	10,10,19	0.25	0	9,9,18	0.49	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	LFA	A	324	-	3,3,19	0.36	0	2,2,18	0.62	0
3	LFA	A	325	-	4,4,19	0.32	0	3,3,18	0.34	0
3	LFA	A	326	-	9,9,19	0.29	0	8,8,18	0.43	0
3	LFA	A	327	-	11,11,19	0.25	0	10,10,18	0.52	0
4	OLC	A	328	-	24,24,24	0.90	1 (4%)	25,25,25	0.77	1 (4%)
4	OLC	A	329	-	24,24,24	0.94	1 (4%)	25,25,25	0.87	1 (4%)
4	OLC	A	330	-	11,11,24	1.37	1 (9%)	12,12,25	1.11	1 (8%)
5	GOL	A	331	-	5,5,5	0.28	0	5,5,5	0.22	0
5	GOL	A	332	-	5,5,5	0.28	0	5,5,5	0.23	0
5	GOL	A	333	-	5,5,5	0.29	0	5,5,5	0.22	0
3	LFA	A	334	-	13,13,19	0.26	0	12,12,18	0.52	0
3	LFA	A	335	-	17,17,19	0.28	0	16,16,18	0.51	0
3	LFA	A	336	-	7,7,19	0.24	0	6,6,18	0.45	0
6	RET	A	337	1	20,20,21	0.63	0	27,27,28	1.63	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LFA	A	302	-	-	0/13/13/17	0/0/0/0
3	LFA	A	303	-	-	0/6/6/17	0/0/0/0
3	LFA	A	304	-	-	0/3/3/17	0/0/0/0
3	LFA	A	305	-	-	0/2/2/17	0/0/0/0
3	LFA	A	306	-	-	0/10/10/17	0/0/0/0
3	LFA	A	307	-	-	0/5/5/17	0/0/0/0
3	LFA	A	308	-	-	0/5/5/17	0/0/0/0
3	LFA	A	309	-	-	0/9/9/17	0/0/0/0
3	LFA	A	310	-	-	0/7/7/17	0/0/0/0
3	LFA	A	311	-	-	0/2/2/17	0/0/0/0
3	LFA	A	312	-	-	0/2/2/17	0/0/0/0
3	LFA	A	313	-	-	0/8/8/17	0/0/0/0
3	LFA	A	314	-	-	0/9/9/17	0/0/0/0
3	LFA	A	315	-	-	0/7/7/17	0/0/0/0
3	LFA	A	316	-	-	0/3/3/17	0/0/0/0
3	LFA	A	317	-	-	0/2/2/17	0/0/0/0
3	LFA	A	318	-	-	0/3/3/17	0/0/0/0
3	LFA	A	319	-	-	0/3/3/17	0/0/0/0
3	LFA	A	320	-	-	0/7/7/17	0/0/0/0
3	LFA	A	321	-	-	0/4/4/17	0/0/0/0
3	LFA	A	322	-	-	0/2/2/17	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LFA	A	323	-	-	0/8/8/17	0/0/0/0
3	LFA	A	324	-	-	0/1/1/17	0/0/0/0
3	LFA	A	325	-	-	0/2/2/17	0/0/0/0
3	LFA	A	326	-	-	0/7/7/17	0/0/0/0
3	LFA	A	327	-	-	0/9/9/17	0/0/0/0
4	OLC	A	328	-	-	0/24/24/24	0/0/0/0
4	OLC	A	329	-	-	0/24/24/24	0/0/0/0
4	OLC	A	330	-	-	0/11/11/24	0/0/0/0
5	GOL	A	331	-	-	0/4/4/4	0/0/0/0
5	GOL	A	332	-	-	0/4/4/4	0/0/0/0
5	GOL	A	333	-	-	0/4/4/4	0/0/0/0
3	LFA	A	334	-	-	0/11/11/17	0/0/0/0
3	LFA	A	335	-	-	0/15/15/17	0/0/0/0
3	LFA	A	336	-	-	0/5/5/17	0/0/0/0
6	RET	A	337	1	-	0/13/30/31	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	328	OLC	O20-C1	4.12	1.45	1.33
4	A	329	OLC	O20-C1	4.33	1.46	1.33
4	A	330	OLC	O20-C1	4.38	1.46	1.33

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	337	RET	C18-C5-C6	-3.76	120.32	124.51
6	A	337	RET	C7-C8-C9	-3.14	121.50	126.21
6	A	337	RET	C10-C11-C12	-3.08	114.03	123.31
6	A	337	RET	C11-C10-C9	-2.84	123.26	127.31
4	A	328	OLC	O20-C1-C2	2.22	119.07	111.93

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 60 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	307	LFA	1	0
3	A	314	LFA	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	316	LFA	2	0
3	A	319	LFA	1	0
4	A	329	OLC	1	0
4	A	330	OLC	1	0
5	A	331	GOL	2	0
5	A	333	GOL	10	3
3	A	334	LFA	1	0
3	A	335	LFA	31	0
6	A	337	RET	7	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	271/288 (94%)	-0.16	8 (2%) 50 49	23, 36, 66, 105	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	LYS	6.0
1	A	233	PHE	4.5
1	A	230	VAL	3.7
1	A	232	GLY	3.6
1	A	231	ASP	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	LFA	A	310	10/20	0.33	0.21	74,81,89,91	0
4	OLC	A	330	12/25	0.52	0.37	79,102,118,134	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	LFA	A	335	18/20	0.52	0.25	62,87,124,126	0
3	LFA	A	314	12/20	0.56	0.20	70,80,91,92	0
3	LFA	A	327	12/20	0.62	0.24	63,79,80,84	0
3	LFA	A	321	7/20	0.63	0.22	75,86,91,95	0
3	LFA	A	324	4/20	0.63	0.58	75,79,83,85	0
3	LFA	A	326	10/20	0.66	0.25	78,85,91,92	0
5	GOL	A	332	6/6	0.68	0.30	89,99,99,99	0
4	OLC	A	329	25/25	0.69	0.31	52,86,103,115	0
3	LFA	A	315	10/20	0.71	0.21	53,75,105,105	0
3	LFA	A	303	9/20	0.71	0.15	64,67,73,74	0
3	LFA	A	334	14/20	0.74	0.27	49,56,77,78	14
3	LFA	A	312	5/20	0.76	0.17	59,63,76,84	0
3	LFA	A	308	8/20	0.76	0.32	67,71,83,86	0
3	LFA	A	309	12/20	0.77	0.21	53,77,88,97	0
3	LFA	A	313	11/20	0.78	0.13	73,89,105,113	0
3	LFA	A	325	5/20	0.78	0.33	70,72,73,80	0
3	LFA	A	319	6/20	0.78	0.19	52,74,77,83	0
3	LFA	A	306	13/20	0.79	0.24	45,60,75,80	0
5	GOL	A	331	6/6	0.79	0.43	66,77,82,83	0
3	LFA	A	305	5/20	0.79	0.20	70,78,79,79	0
3	LFA	A	323	11/20	0.80	0.22	73,83,91,94	0
3	LFA	A	320	10/20	0.81	0.29	81,83,87,92	0
3	LFA	A	317	5/20	0.81	0.21	78,78,85,86	0
5	GOL	A	333	6/6	0.82	1.02	87,162,193,213	1
3	LFA	A	318	6/20	0.82	0.12	66,74,76,82	0
3	LFA	A	322	5/20	0.82	0.19	72,73,78,84	0
3	LFA	A	302	16/20	0.83	0.15	46,60,76,80	0
4	OLC	A	328	25/25	0.84	0.18	54,73,90,94	0
3	LFA	A	316	6/20	0.88	0.16	69,72,80,81	0
3	LFA	A	304	6/20	0.89	0.33	60,67,74,76	0
3	LFA	A	307	8/20	0.90	0.14	59,69,75,78	0
3	LFA	A	311	5/20	0.91	0.16	71,73,81,82	0
3	LFA	A	336	8/20	0.92	0.10	53,64,66,67	0
6	RET	A	337	20/21	0.94	0.15	25,30,37,41	0
2	NA	A	301	1/1	0.96	0.23	42,42,42,42	0

## 6.5 Other polymers

There are no such residues in this entry.