



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 02:20 pm GMT

PDB ID : 1S2N
Title : Crystal structure of a cold adapted subtilisin-like serine proteinase
Authors : Arnorsdottir, J.; Kristjansson, M.M.; Ficner, R.
Deposited on : 2004-01-09
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1296 (2.46-2.42)
Clashscore	122126	1359 (2.46-2.42)
Ramachandran outliers	120053	1350 (2.46-2.42)
Sidechain outliers	120020	1350 (2.46-2.42)
RSRZ outliers	108989	1278 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>..</div> </div> </div>
1	B	284	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>14%</div> <div>..</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called extracellular subtilisin-like serine proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	4	5	0
			2043	1239	365	430	9			
1	B	273	Total	C	N	O	S	0	2	0
			1973	1200	351	415	7			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Ca	0	0
			3	3		
2	A	3	Total	Ca	0	0
			3	3		

- Molecule 3 is phenylmethanesulfonic acid (three-letter code: PMS) (formula: C₇H₈O₃S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			10	7	2	1		
3	B	1	Total	C	O	S	6	0
			10	7	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	194	Total	O	0	0
			194	194		
4	B	184	Total	O	0	0
			184	184		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: extracellular subtilisin-like serine proteinase



- Molecule 1: extracellular subtilisin-like serine proteinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.42Å 36.84Å 143.06Å 90.00° 93.88° 90.00°	Depositor
Resolution (Å)	29.11 – 2.44 29.13 – 2.44	Depositor EDS
% Data completeness (in resolution range)	97.7 (29.11-2.44) 97.7 (29.13-2.44)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.177 , 0.247 0.185 , 0.255	Depositor DCC
R_{free} test set	1754 reflections (10.17%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.196	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 31.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4420	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/2098 (0.0%)	0.87	15/2855 (0.5%)
1	B	0.38	0/2014	0.77	13/2741 (0.5%)
All	All	0.42	1/4112 (0.0%)	0.82	28/5596 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	20	ARG	CD-NE	-12.66	1.25	1.46

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	20	ARG	CD-NE-CZ	16.44	146.62	123.60
1	A	20	ARG	CG-CD-NE	12.01	137.01	111.80
1	A	19	ASP	CB-CG-OD2	7.37	124.94	118.30
1	B	260	ASP	CB-CG-OD2	7.10	124.69	118.30
1	A	27	ASP	CB-CG-OD2	6.91	124.52	118.30
1	B	19	ASP	CB-CG-OD2	6.78	124.40	118.30
1	B	56	ASP	CB-CG-OD2	6.65	124.28	118.30
1	A	210	ASP	CB-CG-OD2	6.58	124.22	118.30
1	B	9	ASP	CB-CG-OD2	6.57	124.21	118.30
1	B	63	ASP	CB-CG-OD2	6.30	123.97	118.30
1	A	56	ASP	CB-CG-OD2	6.11	123.80	118.30
1	A	9	ASP	CB-CG-OD2	6.10	123.79	118.30
1	A	113	ASP	CB-CG-OD2	6.02	123.72	118.30
1	A	12	ASP	CB-CG-OD2	5.92	123.62	118.30
1	B	210	ASP	CB-CG-OD2	5.92	123.62	118.30
1	B	61	ASP	CB-CG-OD2	5.90	123.61	118.30
1	A	260	ASP	CB-CG-OD2	5.79	123.51	118.30
1	B	27	ASP	CB-CG-OD2	5.77	123.49	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	61	ASP	CB-CG-OD2	5.63	123.37	118.30
1	A	63	ASP	CB-CG-OD2	5.62	123.36	118.30
1	B	196	ASP	CB-CG-OD2	5.56	123.31	118.30
1	A	37	ASP	CB-CG-OD2	5.38	123.14	118.30
1	B	113	ASP	CB-CG-OD2	5.33	123.10	118.30
1	A	274	ASP	CB-CG-OD2	5.23	123.00	118.30
1	A	59	ASP	CB-CG-OD2	5.15	122.94	118.30
1	B	59	ASP	CB-CG-OD2	5.12	122.91	118.30
1	B	274	ASP	CB-CG-OD2	5.04	122.83	118.30
1	B	37	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2043	0	1911	15	0
1	B	1973	0	1854	14	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	10	0	7	1	0
3	B	10	0	7	1	0
4	A	194	0	0	1	0
4	B	184	0	0	1	0
All	All	4420	0	3779	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:SER:OG	3:B:2284:PMS:S	2.40	0.78
1:B:258:VAL:H	1:B:266:ASN:HD21	1.34	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:VAL:H	1:A:266:ASN:HD21	1.35	0.71
1:A:106:SER:HB2	4:A:1452:HOH:O	2.01	0.60
1:B:223:THR:N	1:B:224:PRO:HD2	2.23	0.54
1:A:1:GLN:O	1:A:21:ASN:HA	2.09	0.53
1:B:273:ALA:O	1:B:274:ASP:C	2.48	0.52
1:A:223:THR:N	1:A:224:PRO:HD2	2.25	0.52
1:A:220:SER:CB	3:A:1284:PMS:S	2.99	0.50
1:A:86:ALA:HB1	1:A:89:VAL:HG23	1.96	0.48
1:B:229:VAL:HA	1:B:232:LEU:HD22	1.95	0.47
1:A:271:SER:OG	1:A:272:LEU:HD12	2.16	0.46
1:A:229:VAL:HA	1:A:232:LEU:HD22	1.96	0.46
1:B:94:VAL:O	1:B:94:VAL:HG12	2.16	0.46
1:B:99:CYS:HB2	4:B:2421:HOH:O	2.16	0.46
1:B:86:ALA:HB1	1:B:89:VAL:HG23	1.98	0.45
1:B:271:SER:OG	1:B:272:LEU:HD12	2.17	0.44
1:A:265[B]:THR:HG22	1:A:267:LYS:HG3	2.00	0.43
1:A:94:VAL:HG12	1:A:94:VAL:O	2.19	0.43
1:A:68:ASN:OD1	1:A:70[A]:HIS:HD2	2.01	0.42
1:B:158:SER:H	1:B:190:ASN:HD21	1.69	0.41
1:B:69:GLY:HA2	1:B:209:TYR:HB3	2.02	0.41
1:A:185:ARG:NH2	1:A:262[B]:ARG:HD2	2.36	0.41
1:B:86:ALA:HB2	1:B:232:LEU:HD13	2.02	0.41
1:A:274:ASP:HB3	1:A:279:PRO:HD3	2.03	0.40
1:A:69:GLY:HA2	1:A:209:TYR:HB3	2.02	0.40
1:B:200:PRO:O	1:B:222:ALA:HA	2.21	0.40
1:B:70:HIS:O	1:B:74:VAL:HG13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/284 (100%)	274 (96%)	10 (4%)	0	100	100
1	B	273/284 (96%)	265 (97%)	8 (3%)	0	100	100
All	All	557/568 (98%)	539 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/220 (102%)	205 (92%)	19 (8%)	12	13
1	B	214/220 (97%)	199 (93%)	15 (7%)	16	21
All	All	438/440 (100%)	404 (92%)	34 (8%)	14	17

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	GLN
1	A	40	VAL
1	A	41	ASN
1	A	74	VAL
1	A	215	THR
1	A	221	MET
1	A	229	VAL
1	A	232	LEU
1	A	234	LEU
1	A	240	LEU
1	A	243	LEU
1	A	245	LEU
1	A	252	ARG
1	A	257	LYS
1	A	262[A]	ARG
1	A	262[B]	ARG
1	A	266	ASN
1	A	269	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	278	GLU
1	B	40	VAL
1	B	41	ASN
1	B	74	VAL
1	B	215	THR
1	B	221	MET
1	B	229	VAL
1	B	232	LEU
1	B	234	LEU
1	B	240	LEU
1	B	243	LEU
1	B	245	LEU
1	B	252	ARG
1	B	257	LYS
1	B	266	ASN
1	B	269	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	42	ASN
1	A	43	ASN
1	A	118	ASN
1	A	126	ASN
1	A	190	ASN
1	A	256	ASN
1	A	266	ASN
1	B	41	ASN
1	B	42	ASN
1	B	43	ASN
1	B	126	ASN
1	B	133	GLN
1	B	190	ASN
1	B	256	ASN
1	B	266	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PMS	A	1284	1	7,10,11	0.71	0	11,12,15	2.35	3 (27%)
3	PMS	B	2284	-	7,10,11	30.73	1 (14%)	11,12,15	18.74	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PMS	A	1284	1	-	0/4/4/5	0/1/1/1
3	PMS	B	2284	-	-	0/4/4/5	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2284	PMS	C-C1	81.27	2.86	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2284	PMS	C-C1-C2	-46.62	68.20	120.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2284	PMS	C1-C-S	-22.03	52.28	110.72
3	A	1284	PMS	C-C1-C2	-2.40	117.87	120.56
3	B	2284	PMS	O2S-S-C	-2.04	102.17	105.55
3	A	1284	PMS	C-C1-C6	2.11	122.94	120.56
3	B	2284	PMS	O1S-S-C	6.30	115.97	105.55
3	A	1284	PMS	O1S-S-C	6.90	116.98	105.55
3	B	2284	PMS	C-C1-C6	34.00	158.75	120.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1284	PMS	1	0
3	B	2284	PMS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/284 (98%)	-0.25	7 (2%) 57 53	17, 18, 21, 38	1 (0%)
1	B	273/284 (96%)	-0.26	4 (1%) 73 71	17, 18, 21, 32	1 (0%)
All	All	554/568 (97%)	-0.26	11 (1%) 65 62	17, 18, 21, 38	2 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	281	CYS	4.3
1	A	278	GLU	4.1
1	B	2	SER	3.5
1	A	2	SER	3.4
1	A	133	GLN	2.8
1	A	1	GLN	2.7
1	A	279	PRO	2.7
1	B	274	ASP	2.6
1	A	280	ASP	2.6
1	B	263	GLY	2.4
1	B	3	ASN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PMS	A	1284	10/11	0.84	0.23	46,47,47,47	0
3	PMS	B	2284	10/11	0.85	0.19	40,40,60,60	6
2	CA	A	1290	1/1	0.94	0.05	28,28,28,28	0
2	CA	A	1291	1/1	0.98	0.05	30,30,30,30	0
2	CA	B	2291	1/1	0.98	0.02	27,27,27,27	0
2	CA	B	2290	1/1	0.98	0.02	22,22,22,22	0
2	CA	B	2292	1/1	0.99	0.05	29,29,29,29	0
2	CA	A	1292	1/1	0.99	0.02	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.