



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 03:34 pm GMT

PDB ID : 1S3T  
Title : BORATE INHIBITED BACILLUS PASTEURII UREASE CRYSTAL  
STRUCTURE  
Authors : Benini, S.; Rypniewski, W.R.; Wilson, K.S.; Ciurli, S.; Mangani, S.  
Deposited on : 2004-01-14  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	<div> <div>%</div> <div> <div></div> <div>96%</div> <div>•</div> </div> </div>
2	B	126	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>5% • •</div> </div> </div>
3	C	570	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>8% •</div> </div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6376 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urease gamma subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			772	487	129	150	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	CXM	MET	MODIFIED RESIDUE	UNP P41022

- Molecule 2 is a protein called Urease beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	0	0	0
			913	571	163	178	1			

- Molecule 3 is a protein called Urease alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	570	Total	C	N	O	S	2	0	0
			4274	2687	735	829	23			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLU	ARG	CONFLICT	UNP P41020
C	28	TRP	GLY	CONFLICT	UNP P41020
C	29	ILE	-	INSERTION	UNP P41020
C	36	THR	TYR	CONFLICT	UNP P41020
C	37	THR	TYR	CONFLICT	UNP P41020
C	38	TYR	LEU	CONFLICT	UNP P41020
C	220	KCX	LYS	MODIFIED RESIDUE	UNP P41020
C	263	LEU	VAL	CONFLICT	UNP P41020
C	420	ILE	MET	CONFLICT	UNP P41020

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ni	0	0
			2	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is BORIC ACID (three-letter code: BO3) (formula: BH<sub>3</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	B	O	0	0
			4	1	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	53	Total	O	0	0
			53	53		
7	B	72	Total	O	0	0
			72	72		
7	C	281	Total	O	0	0
			281	281		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Urease gamma subunit



- Molecule 2: Urease beta subunit



- Molecule 3: Urease alpha subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	130.91Å 130.91Å 189.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.77 – 2.10 24.76 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.77-2.10) 99.5 (24.76-2.10)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	0.13	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.177 , 0.205 0.188 , 0.208	Depositor DCC
$R_{free}$ test set	1142 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.1	Xtriage
Anisotropy	0.479	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6376	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, NI, BO3, CXM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/772	0.72	0/1041
2	B	1.56	2/924 (0.2%)	0.80	1/1246 (0.1%)
3	C	0.47	0/4343	0.82	7/5895 (0.1%)
All	All	0.75	2/6039 (0.0%)	0.81	8/8182 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	13	ARG	CZ-NH1	36.88	1.80	1.33
2	B	13	ARG	CZ-NH2	26.54	1.67	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	328	ASN	CA-CB-CG	-14.77	80.91	113.40
2	B	13	ARG	NH1-CZ-NH2	-14.10	103.89	119.40
3	C	399	ASP	CB-CG-OD2	6.97	124.58	118.30
3	C	144	ASP	CB-CG-OD2	6.29	123.96	118.30
3	C	328	ASN	CB-CG-OD1	6.04	133.68	121.60
3	C	538	ASP	CB-CG-OD2	5.77	123.50	118.30
3	C	382	ASP	CB-CG-OD2	5.24	123.01	118.30
3	C	527	ASP	CB-CG-OD2	5.23	123.01	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	772	0	776	2	0
2	B	913	0	875	5	0
3	C	4274	0	4199	27	0
4	C	2	0	0	0	0
5	C	5	0	0	0	0
6	C	4	0	1	0	0
7	A	53	0	0	0	0
7	B	72	0	0	0	0
7	C	281	0	0	1	1
All	All	6376	0	5851	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:13:ARG:NH2	2:B:13:ARG:CZ	1.67	1.52
2:B:13:ARG:NH1	2:B:13:ARG:CZ	1.81	1.43
3:C:201:HIS:HE1	3:C:222:HIS:H	1.42	0.65
2:B:125:VAL:O	2:B:126:GLU:HB2	2.02	0.60
3:C:201:HIS:CE1	3:C:222:HIS:H	2.20	0.59
3:C:300:SER:OG	3:C:352:HIS:HE1	1.85	0.59
3:C:158:PHE:CE2	3:C:418:GLN:HG3	2.40	0.56
2:B:22:ASN:HD21	2:B:66:ARG:HH22	1.53	0.55
3:C:70:LEU:HD11	3:C:86:ASP:HB3	1.88	0.55
3:C:383:LYS:NZ	3:C:387:GLN:HE22	2.04	0.55
3:C:181:GLU:OE2	3:C:507:LYS:NZ	2.40	0.51
3:C:249:HIS:CE1	3:C:281:GLY:HA3	2.46	0.51
3:C:383:LYS:HZ3	3:C:387:GLN:HE22	1.59	0.51
3:C:201:HIS:CE1	3:C:221:ILE:HA	2.48	0.48
2:B:22:ASN:ND2	2:B:66:ARG:HH22	2.12	0.48
3:C:274:PHE:O	3:C:275:HIS:C	2.54	0.46
3:C:201:HIS:HE1	3:C:221:ILE:HA	1.80	0.46
3:C:302:ASN:O	3:C:376:ARG:HD3	2.17	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:220:KCX:CX	3:C:222:HIS:HD2	2.30	0.45
3:C:300:SER:OG	3:C:352:HIS:CE1	2.69	0.44
3:C:470:THR:N	3:C:471:PRO:CD	2.80	0.44
3:C:302:ASN:N	3:C:303:PRO:CD	2.80	0.44
3:C:362:THR:O	3:C:368:GLY:HA3	2.17	0.44
3:C:384:MET:HA	3:C:384:MET:HE2	2.00	0.43
1:A:73:VAL:N	1:A:74:PRO:CD	2.81	0.43
1:A:30:LEU:HD13	1:A:38:ILE:HD12	2.01	0.43
3:C:132:GLY:HA3	3:C:155:THR:OG1	2.20	0.42
3:C:358:SER:HA	3:C:531:ASN:HB3	2.02	0.41
3:C:7:GLN:CG	7:C:868:HOH:O	2.67	0.41
3:C:76:ILE:O	3:C:82:ILE:HA	2.20	0.41
3:C:303:PRO:HG3	3:C:367:MET:O	2.21	0.41
3:C:302:ASN:N	3:C:303:PRO:HD2	2.36	0.41
3:C:284:ALA:HA	3:C:285:PRO:HA	1.84	0.41
3:C:316:LEU:O	3:C:320:MET:HG2	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:C:659:HOH:O	7:C:882:HOH:O[12_565]	1.66	0.54

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
2	B	120/126 (95%)	116 (97%)	3 (2%)	1 (1%)	21	16
3	C	567/570 (100%)	546 (96%)	20 (4%)	1 (0%)	49	51
All	All	785/796 (99%)	758 (97%)	25 (3%)	2 (0%)	43	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	99	ILE
3	C	275	HIS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/84 (96%)	81 (100%)	0	100	100
2	B	90/105 (86%)	89 (99%)	1 (1%)	76	81
3	C	445/460 (97%)	439 (99%)	6 (1%)	71	78
All	All	616/649 (95%)	609 (99%)	7 (1%)	76	81

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	5	ASN
3	C	1	MET
3	C	77	LEU
3	C	158	PHE
3	C	253	LEU
3	C	323	HIS
3	C	418	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
2	B	22	ASN
3	C	201	HIS
3	C	267	ASN
3	C	352	HIS
3	C	387	GLN
3	C	472	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CXM	A	1	1	7,10,11	0.70	0	5,11,13	1.38	2 (40%)
3	KCX	C	220	3,4	8,11,12	1.11	1 (12%)	6,12,14	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	A	1	1	-	0/6/10/12	0/0/0/0
3	KCX	C	220	3,4	-	0/6/10/12	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	220	KCX	CE-NZ	2.05	1.50	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	CXM	O-C-CA	-2.16	120.06	125.09
1	A	1	CXM	CB-CA-C	-2.11	107.82	111.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	220	KCX	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	BO3	C	602	4	3,3,3	0.46	0	3,3,3	0.72	0
5	SO4	C	603	-	4,4,4	0.24	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BO3	C	602	4	-	0/0/0/0	0/0/0/0
5	SO4	C	603	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	99/100 (99%)	-0.40	1 (1%) 82 85	15, 21, 26, 34	0
2	B	122/126 (96%)	-0.17	2 (1%) 72 76	18, 23, 30, 43	0
3	C	569/570 (99%)	-0.45	6 (1%) 80 84	15, 19, 32, 47	2 (0%)
All	All	790/796 (99%)	-0.40	9 (1%) 80 84	15, 20, 30, 47	2 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	126	GLU	6.3
3	C	396	ASN	4.3
1	A	100	SER	4.0
3	C	324	HIS	2.7
3	C	390	PRO	2.5
3	C	549	ASP	2.3
2	B	5	ASN	2.1
3	C	395	LYS	2.0
3	C	322	CYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CXM	A	1	11/12	0.96	0.09	19,21,28,28	0
3	KCX	C	220	12/13	0.97	0.07	16,17,19,19	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	BO3	C	602	4/4	0.93	0.12	23,23,24,25	0
5	SO4	C	603	5/5	0.96	0.12	39,39,40,41	0
4	NI	C	600	1/1	1.00	0.02	21,21,21,21	0
4	NI	C	601	1/1	1.00	0.02	20,20,20,20	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.