



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 02:11 pm GMT

PDB ID : 1SO6  
Title : Crystal structure of E112Q/H136A double mutant of 3-keto-L-gulonate 6-phosphate decarboxylase with bound L-threonohydroxamate 4-phosphate  
Authors : Wise, E.L.; Yew, W.S.; Gerlt, J.A.; Rayment, I.  
Deposited on : 2004-03-12  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	5502 (1.90-1.90)
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>.</div> </div> </div>
1	B	216	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TX4	A	2301	X	-	-	-
3	TX4	B	1301	X	-	-	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3735 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-keto-L-gulonate 6-phosphate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	0	0	0
			1642	1042	285	307	8			
1	B	215	Total	C	N	O	S	0	0	0
			1625	1031	285	301	8			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	112	GLN	GLU	ENGINEERED	UNP P39304
A	136	ALA	HIS	ENGINEERED	UNP P39304
B	112	GLN	GLU	ENGINEERED	UNP P39304
B	136	ALA	HIS	ENGINEERED	UNP P39304

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is L-THREONOHYDROXAMATE 4-PHOSPHATE (three-letter code: TX4) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			14	4	1	8	1		
3	B	1	Total	C	N	O	P	0	0
			14	4	1	8	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	226	Total	O	0	0
			226	226		
4	B	212	Total	O	0	0
			212	212		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-keto-L-gulonate 6-phosphate decarboxylase



- Molecule 1: 3-keto-L-gulonate 6-phosphate decarboxylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.77Å 41.81Å 90.94Å 90.00° 97.22° 90.00°	Depositor
Resolution (Å)	91.29 – 1.90 47.77 – 1.90	Depositor EDS
% Data completeness (in resolution range)	95.0 (91.29-1.90) 95.0 (47.77-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.18 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, $R_{free}$	0.158 , 0.208 0.169 , 0.215	Depositor DCC
$R_{free}$ test set	1736 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.9	Xtriage
Anisotropy	1.038	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 52.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3735	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TX4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.07	3/1670 (0.2%)	0.90	6/2267 (0.3%)
1	B	0.84	1/1652 (0.1%)	0.99	12/2243 (0.5%)
All	All	0.96	4/3322 (0.1%)	0.95	18/4510 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	215	TRP	C-N	23.67	1.75	1.33
1	A	2	SER	C-N	20.40	1.80	1.34
1	B	115	GLY	C-N	14.79	1.68	1.34
1	A	136	ALA	C-N	5.12	1.45	1.34

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	113	LEU	O-C-N	-12.00	103.50	122.70
1	B	115	GLY	O-C-N	8.78	136.75	122.70
1	B	115	GLY	CA-C-N	-7.78	100.09	117.20
1	A	11	ASP	CB-CG-OD2	7.41	124.97	118.30
1	A	140	ASP	CB-CG-OD2	7.16	124.75	118.30
1	A	136	ALA	O-C-N	-6.94	111.59	122.70
1	B	82	ASP	CB-CG-OD2	6.84	124.45	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	11	ASP	CB-CG-OD2	6.71	124.34	118.30
1	B	30	ASP	CB-CG-OD2	6.60	124.24	118.30
1	B	152	ASP	CB-CG-OD2	6.59	124.23	118.30
1	B	196	ASP	CB-CG-OD2	6.41	124.07	118.30
1	A	100	ASP	CB-CG-OD2	6.31	123.98	118.30
1	B	91	ASP	CB-CG-OD2	5.95	123.66	118.30
1	B	115	GLY	C-N-CA	5.79	136.16	121.70
1	B	176	ASP	CB-CG-OD2	5.74	123.47	118.30
1	B	113	LEU	CA-C-N	5.71	129.76	117.20
1	A	215	TRP	O-C-N	-5.30	114.19	123.20
1	A	176	ASP	CB-CG-OD2	5.22	123.00	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	113	LEU	Mainchain
1	B	114	THR	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1642	0	1645	17	0
1	B	1625	0	1627	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	14	0	7	0	0
3	B	14	0	7	0	0
4	A	226	0	0	4	1
4	B	212	0	0	1	1
All	All	3735	0	3286	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:115:GLY:C	1:B:116:TYR:N	1.68	1.46
1:A:215:TRP:C	1:A:216:GLY:N	1.75	1.39
1:A:2:SER:C	1:A:3:LEU:N	1.81	1.34
1:B:115:GLY:CA	1:B:116:TYR:N	2.45	0.79
1:B:113:LEU:O	1:B:114:THR:CB	2.36	0.74
1:B:60:LEU:HD11	1:B:110:GLN:NE2	2.04	0.73
1:A:100:ASP:HB2	4:A:2522:HOH:O	1.96	0.65
1:A:89:CYS:O	1:B:115:GLY:HA3	1.99	0.62
1:A:122:ALA:HA	1:A:125:TRP:CE3	2.36	0.61
1:A:2:SER:N	4:A:2397:HOH:O	2.35	0.58
1:B:122:ALA:HA	1:B:125:TRP:CE3	2.40	0.55
1:A:209:ARG:HD2	4:A:2464:HOH:O	2.09	0.52
1:A:2:SER:CA	1:A:3:LEU:N	2.70	0.52
1:A:158:ARG:O	1:A:162:MET:HG3	2.11	0.50
1:A:89:CYS:O	1:B:115:GLY:CA	2.60	0.49
1:B:60:LEU:HD11	1:B:110:GLN:HE22	1.76	0.48
1:A:2:SER:HA	1:A:3:LEU:HA	1.94	0.48
1:B:23:ARG:HD3	4:B:1408:HOH:O	2.13	0.48
1:A:126:ARG:HG2	1:A:164:PHE:CZ	2.51	0.46
1:A:176:ASP:OD2	4:A:2491:HOH:O	2.21	0.46
1:B:153:ILE:HG23	1:B:183:ILE:HD11	1.99	0.45
1:B:60:LEU:HD12	1:B:83:TRP:HE3	1.82	0.45
1:B:25:ILE:HD12	1:B:25:ILE:C	2.38	0.44
1:B:69:GLY:O	1:B:73:SER:HB2	2.18	0.43
1:A:177:LEU:N	1:A:178:PRO:CD	2.82	0.42
1:A:202:GLU:O	1:A:206:GLN:HG3	2.18	0.42
1:A:126:ARG:HD2	1:A:126:ARG:HA	1.92	0.41
1:A:25:ILE:C	1:A:25:ILE:HD12	2.41	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2311:HOH:O	4:B:1512:HOH:O[2_655]	2.06	0.14

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/216 (99%)	210 (99%)	3 (1%)	0	100	100
1	B	213/216 (99%)	209 (98%)	4 (2%)	0	100	100
All	All	426/432 (99%)	419 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/168 (99%)	165 (99%)	1 (1%)	87	88
1	B	162/168 (96%)	161 (99%)	1 (1%)	87	88
All	All	328/336 (98%)	326 (99%)	2 (1%)	87	88

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	SER
1	B	112	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	80	ASN
1	A	186	HIS
1	A	206	GLN
1	B	7	GLN
1	B	13	GLN
1	B	110	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	112	GLN
1	B	206	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	TX4	A	2301	2	11,13,13	1.63	1 (9%)	14,18,18	1.54	2 (14%)
3	TX4	B	1301	2	11,13,13	1.25	1 (9%)	14,18,18	1.81	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TX4	A	2301	2	1/1/4/5	0/14/16/16	0/0/0/0
3	TX4	B	1301	2	1/1/4/5	0/14/16/16	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2301	TX4	O1-C1	-5.14	1.31	1.40
3	B	1301	TX4	O1-C1	-3.41	1.34	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1301	TX4	O6-P-O5	2.22	116.37	107.59
3	A	2301	TX4	O6-P-O5	2.43	117.20	107.59
3	A	2301	TX4	O1-C1-C2	3.81	118.53	109.46
3	B	1301	TX4	O1-C1-C2	5.59	122.77	109.46

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	1301	TX4	C1
3	A	2301	TX4	C1

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	2:SER	C	3:LEU	N	1.81
1	A	215:TRP	C	216:GLY	N	1.75

*Continued on next page...*

*Continued from previous page...*

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	115:GLY	C	116:TYR	N	1.68

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/216 (99%)	-0.53	2 (0%) 84 86	10, 16, 25, 43	0
1	B	215/216 (99%)	-0.52	4 (1%) 66 70	10, 15, 27, 43	0
All	All	430/432 (99%)	-0.52	6 (1%) 75 78	10, 16, 26, 43	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	SER	4.1
1	B	115	GLY	2.8
1	B	114	THR	2.6
1	B	116	TYR	2.4
1	A	216	GLY	2.2
1	B	151	ALA	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TX4	B	1301	14/14	0.99	0.06	7,10,16,18	0
2	MG	B	1300	1/1	0.99	0.04	11,11,11,11	0
2	MG	A	2300	1/1	0.99	0.03	11,11,11,11	0
3	TX4	A	2301	14/14	0.99	0.06	10,12,18,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.